

Research paper

Predictive models for first-onset and persistence of depression and anxiety among university students[☆]



Laura Ballester^{a,b,c}, Itxaso Alayo^{a,c}, Gemma Vilagut^{a,c}, Philippe Mortier^{a,c}, José Almenara^d, Ana Isabel Cebrià^{e,f}, Enrique Echeburúa^g, Andrea Gabilondo^h, Margalida Giliⁱ, Carolina Lagares^d, José Antonio Piqueras^j, Miquel Rocaⁱ, Victoria Soto-Sanz^j, Maria Jesús Blasco^{a,c}, Pere Castellví^k, Andrea Miranda-Mendizabal^k, Ronny Bruffaerts^l, Randy P. Auerbach^m, Matthew K. Nockⁿ, Ronald C. Kessler^o, Jordi Alonso^{a,c,p,*}, on behalf of the on behalf of the UNIVERSAL study group

^a Health Services Research Group, IMIM (Institut Hospital del Mar d'Investigacions Mèdiques), Barcelona, Spain

^b Girona University (UdG), Girona, Spain

^c CIBER Epidemiología y Salud Pública (CIBERESP), Spain

^d University of Cádiz (UCA), Spain

^e Department of Mental Health, Corporació Sanitària Parc Taulí, Sabadell, Spain

^f CIBER Salud Mental (CIBERSAM), Madrid, Spain

^g University of the Basque Country (UPV-EHU), San Sebastián, Spain

^h BioDonostia Health Research Institute, Osakidetza, San Sebastián, Spain

ⁱ Institut Universitari d'Investigació en Ciències de la Salut (IUNICS-IDISBA), Rediapp, University of Balearic Islands (UIB), Palma de Mallorca, Spain

^j Miguel Hernández University of Elche (UMH), Spain

^k International University of Catalonia (UIC), Barcelona, Spain

^l Universitair Psychiatrisch Centrum (UPC-KUL), Center for Public Health Psychiatry, KU Leuven, Leuven, Belgium

^m Department of Psychiatry, Columbia University, New York, United States

ⁿ Department of Psychology, Harvard University, Boston, MA, United States

^o Department of Health Care Policy, Harvard Medical School, Boston, MA, United States

^p Department of Medicine and Life Sciences, Universitat Pompeu Fabra, Barcelona, Spain

ARTICLE INFO

Keywords:

Depression

Anxiety

Epidemiology

Universities

Prediction

High risk

ABSTRACT

Background: Depression and anxiety are both prevalent among university students. They frequently co-occur and share risk factors. Yet few studies have focused on identifying students at highest risk of first-onset and persistence of either of these conditions.

Methods: Multicenter cohort study among Spanish first-year university students. At baseline, students were assessed for lifetime and 12-month Major Depressive Episode and/or Generalized Anxiety Disorder (MDE-GAD), other mental disorders, childhood-adolescent adversities, stressful life events, social support, socio-demographics, and psychological factors using web-based surveys; 12-month MDE-GAD was again assessed at 12-month follow-up.

Abbreviations: aOR, Adjusted Odds Ratio; ASRS, Adult ADHD Self-Report Scales; AIC, Akaike's information criterion; AUDIT, Alcohol Use Disorders Identification Test; AUC, Area under the Curve; ADHD, Attention-deficit/hyperactivity disorder; UIB, Balearic Islands University; UPV-EHU, Basque Country University; UCA, Cádiz University; CIDI-SC, Composite International Diagnostic Interview Screening Scales; CI, Confidence Interval; GAD, Generalized anxiety disorder; ITC, International Test Commission; IPW, Inverse-probability weighting; MDE, Major Depressive Episode; MDE-GAD, Major Depressive Episode and/or Generalized Anxiety Disorder; UMH, Miguel Hernández University; MI, Multiple Imputation; OR, Odds Ratio; UPF, Pompeu Fabra University; SDS, Sheehan Disability Scale; SE, Standard Error; SD, Standard Deviation; WMH-ICS, World Mental Health International College Student initiative.

[☆] The UNIVERSAL study group is formed by: Itxaso Alayo, Jordi Alonso, José Almenara, Laura Ballester, Gabriela Barbaglia, Maria Jesús Blasco, Pere Castellví, Ana Isabel Cebrià, Enrique Echeburúa, Andrea Gabilondo, Carlos G. Forero, Margalida Gili, Álvaro Iruin, Carolina Lagares, David Leiva, Andrea Miranda-Mendizabal, Oleguer Parès-Badell, María Teresa Pérez-Vázquez, José Antonio Piqueras, Miquel Roca, Jesús Rodríguez-Marín, Albert Sesé, Victoria Soto-Sanz, Gemma Vilagut and Margalida Vives.

* Corresponding author at: Health Services Research Group, Barcelona Biomedical Research Park, Dr. Aiguader, 88 08003 Barcelona, Spain.

E-mail address: jalonso@imim.es (J. Alonso).

<https://doi.org/10.1016/j.jad.2021.10.135>

Received 13 September 2020; Received in revised form 21 October 2021; Accepted 25 October 2021

Available online 6 April 2022

0165-0327/© 2022 Elsevier B.V. All rights reserved.

Results: A total of 1253 students participated in both surveys (59.2% of baseline respondents; mean age = 18.7 (SD = 1.3); 56.0% female). First-onset of MDE-GAD at follow-up was 13.3%. Also 46.7% of those with baseline MDE-GAD showed persistence at follow-up. Childhood/Adolescence emotional abuse or neglect (OR= 4.33), prior bipolar spectrum disorder (OR= 4.34), prior suicidal ideation (OR=4.85) and prior lifetime symptoms of MDE (ORs=2.33-3.63) and GAD (ORs=2.15-3.75) were strongest predictors of first-onset MDE-GAD. Prior suicidal ideation (OR=3.17) and prior lifetime GAD symptoms (ORs=2.38-4.02) were strongest predictors of MDE-GAD persistence. Multivariable predictions from baseline showed AUCs of 0.76 for first-onset and 0.81 for persistence. 74.9% of first-onset MDE-GAD cases occurred among 30% students with highest predicted risk at baseline.

Limitations: Self-report data were used; external validation of the multivariable prediction models is needed.

Conclusion: MDE-GAD among university students is frequent, suggesting the need to implement web-based screening at university entrance that identify those students with highest risk.

1. Introduction

The transition to university may be stressful and increase the risk for onset of mental disorders (Arnett, 2000; Auerbach et al., 2018). Major depressive episode (MDE) and generalized anxiety disorder (GAD) are the most prevalent mental disorders among university students (Auerbach et al., 2018, 2016; Ibrahim et al., 2013; Vázquez et al., 2011). But less is known about their first-onset during the university period and the extent to which any of these disorders persist. Results from longitudinal studies among young adults and university students reported an incidence of MDE at 7% and of GAD at 4% (Beesdo et al., 2010; Ebert et al., 2019). Among adolescents, a longitudinal study found a 8-year incidence of MDE around 13% and GAD of 2% (Benjet et al., 2016). Also, depression and anxiety have been found to be persistent among college students (Zivin et al., 2009). Longitudinal studies are necessary to understand the course of these disorders to guide preventive interventions on campus.

Depression and GAD co-occur frequently (Garber and Weersing, 2010; Gorman, 1996; Kessler et al., 2008; Moffitt et al., 2007), have overlapping risk factors (Dozois et al., 2009), share similar symptoms (Goldberg et al., 2009) and non-specific cognitive and behavioral processes (Kotov, 2011). Additionally, similar efficacy of transdiagnostic interventions to disorder-specific treatments has been found (McEvoy et al., 2009; Newby et al., 2015), suggesting potential benefits of a single transdiagnostic protocol (Barlow et al., 2016). Notwithstanding the commonalities among these disorders, there are no studies focused on the broad occurrence of Major Depressive Episode and/or Generalized Anxiety Disorder (as from now, MDE-GAD) among university students.

There is robust evidence about a number of risk factors associated with depression and anxiety such as childhood adversities (Kessler et al., 2008), stressful life events (Spinoven et al., 2010), prior history of mental disorders (Kessler et al., 2002) and previous symptomatology (Horwath, 1992; Rueter et al., 1999), as well as protective factors such as social support (Galatzer-Levy et al., 2012) and psychological factors (Crockett et al., 2007; Riskind et al., 1987). These known risk and protective factors could allow the development of prediction algorithms to early identify students at risk for the onset or persistence of MDE-GAD. In the mental health field, some efforts have been made to detect at-risk populations early, but most of them are directed to clinical or general adult populations (King et al., 2011; Wang et al., 2014). Developing an algorithm for both MDE-GAD may allow to deploy subsequent preventive interventions that may address their co-occurrence and common underlying factors (Nolen-Hoeksema and Watkins, 2011; Titov et al., 2011).

The present study aims to: (1) estimate 12-month incidence and persistence of MDE-GAD; (2) assess a broad range of risk and protective factors; and (3) develop and validate prediction models for the risk of first-onset and persistence of MDE-GAD among university students.

2. Methods

2.1. Study design

Longitudinal data were obtained from the UNIVERSAL project, a multi-center observational cohort study of university students in five Spanish universities. The UNIVERSAL project is part of the World Mental Health International College Student initiative (WHM-ICS, 2015). Information on the rationale and methods of the UNIVERSAL project has been published elsewhere (Blasco et al., 2016).

2.2. Participants and setting

The study was carried out in five public universities from different Autonomous Regions of Spain: Balearic Islands (UIB), Basque Country (UPV-EHU), Andalusia (UCA), Valencia (UMH), and Catalonia (UPF). These universities represented around 8% of the total number of students in public universities of Spain in the year 2014–15, and their distribution in terms of gender, nationality and academic field was similar to that of the overall population of students in public universities of Spain (Ballester et al., 2020).

In the academic year 2014–2015, all incoming first-year students at the participating universities aged 18 to 24 years old enrolled for the first time in a university were invited to participate in the baseline survey ($N = 16,332$). The baseline survey was carried out over the subsequent 12 months (October 2014–October 2015). Students were recruited in two stages. In the first stage of recruitment, all eligible students were invited to participate (i.e., census sampling). In order to increase participation, a second stage was carried out where a random subsample of eligible students who had not responded to the baseline survey (ranging from 6% to 11% of non-respondents depending on the University) was contacted by e-mail offering a direct economic incentive to complete the baseline survey. At UPV-EHU university, only the first stage was carried out. Baseline respondents were invited by e-mail with a personalized link to complete the follow-up survey twelve months after baseline evaluation. The data collection platform followed the international recommendations and guidelines for computerized assessment (International Test Commission -ITC-, 2005) (International Test Commission, 2019). Ethical approval was provided by the Parc de Salut Mar-Clinical Research Ethics Committee (Reference: 2013/5252/I).

2.3. Variables

Sociodemographic and university-related variables were asked at the beginning of the survey and included: gender, age, country of birth, parental education, urbanicity, religion, sexual orientation, university, academic field and first-term living location during the university period.

Mental disorders. Items taken from Composite International Diagnostic Interview Screening Scales (CIDI-SC) (Kessler et al., 2013a; Kessler and Üstün, 2004) were used to screen for lifetime and 12-month

DSM-IV Major Depressive Episode (MDE) and Generalized Anxiety Disorder (GAD) at baseline, and 12-month disorders at follow-up. CIDI-SC have shown good psychometric properties (Kessler et al., 2013b) and good concordance with blinded clinical diagnoses (Ballester et al., 2019; Kessler et al., 2013b). The outcome variable in this study was a positive screen for 12-month MDE and/or 12-month GAD at follow-up (i.e. 12-month MDE-GAD from now on). We distinguished between: a) first-onset MDE-GAD: 12-month MDE-GAD at follow-up among those without a baseline history of MDE and/or GAD (neither lifetime nor 12-month at baseline); and b) persistent MDE-GAD: 12-month MDE-GAD at follow-up among students with a baseline history of MDE and/or GAD (with or without 12-month diagnostic at baseline).

Bipolar spectrum disorder, panic disorder and drug use disorder were assessed with CIDI-SC. Alcohol use disorder was screened using a modified version of the Alcohol Use Disorders Identification Test (AUDIT) (Saunders et al., 1993). Adult Attention-deficit hyperactivity disorder (Adult ADHD) was assessed with the Adult ADHD Self-Report Scales (ASRS) referring to the previous 6 months (Kessler et al., 2005). Suicidal thoughts and behaviors were assessed with modified versions of the Self-Injurious Thoughts and Behaviors Interview (SITBI) (Nock et al., 2007) and the Columbia-Suicide Severity Rating Scale (C-SSRS) (Posner et al., 2011). All of them were screened for lifetime and 12-month at baseline.

Childhood and adolescent adversities prior to the age of 17 were assessed using 20 items adapted from the CIDI-3.0 (Kessler and Üstün, 2004), the Adverse Childhood Experience Scale (Felitti et al., 1998), and the Bully Survey (Swearer and Cary, 2003) and included: breakdown of family structure (parental death, separation or divorce), family maladaptation (parental psychopathology or household dysfunction), childhood maltreatment (emotional abuse or neglect and physical or sexual abuse) (Blasco et al., 2019), bully victimization and dating violence. The presence of any specific adversity was considered when the response to the corresponding item was different than “never,” except for bully victimization, where the cut-point was “sometimes” or more.

Childhood and adolescent perceived support prior to the age of 17 was assessed using 13 items adapted from the CIDI-3.0 (Kessler and Üstün, 2004), the Psychological Sense of School Membership Scale (Goodenow, 1993), the Adverse Childhood Experience Scale (Felitti et al., 1998), and the Childhood Trauma Questionnaire (Bernstein et al., 1997). Three constructs were considered: relationships within school, family, and peers/others. Confirmatory factor analysis showed good fit with the data (Comparative Fit Index = 0.986; Tucker-Lewis Index = 0.983; RMSEA = 0.055). Scores for the constructs were categorized into tertiles.

Psychological factors included: personality, positive coping strategies and student expectations. Personality was assessed using the Ten Item Personality Inventory (TIPI: Gosling et al., 2003). Scores were calculated as the mean of the two items in each dimension (Gosling et al., 2003). *Positive coping strategies* were assessed using 5 items adapted from the Hurricane Katrina Community Advisory Group Survey (Kessler et al., 2006), and the Army STARRS survey (Ursano, 2012). The five items were added together to give a total coping score. *Student expectations* were assessed using 2 items adapted from the Survey of Student Attitudes, Experiences and Expectations (Round, 2005). The items asked about expectations on academic demands and social life at university on a 7-point Likert scale. The score was obtained as the mean of the item responses.

12-month recent stressful events items were adapted from the Life Events Questionnaire (Brugha and Cragg, 1990), the Deployment Risk and Resilience Inventory (Vogt et al., 2008), and the Department of Defense Survey of Health-Related Behaviors (Bray et al., 2009). *Current severe stress* was assessed using 8 items adapted from the CIDI-3.0 (Kessler and Üstün, 2004) asking about how much stress the students currently had in different areas of their life. The presence of stress in any area of life was considered when the response to the corresponding item was “severe” or more.

Severe role impairment was assessed with an adapted version of the

Sheehan Disability Scale (Leon et al., 1997). This scale is composed of four role domains: home management, college-related and other work, close personal relationships, and social. Severe self-reported role impairment was defined as having a 7–10 rating.

2.4. Analysis

Missing item-level data among respondents to both the baseline and follow-up assessment were imputed using multiple imputation (MI) by chained equations (van Buuren, 2012) with 43 imputed datasets, equivalent to the percentage of incomplete subjects (White et al., 2011), and 10 iterations per imputation.

Out of the 2118 baseline participants, we analyze here those that completed at least 40% of the follow-up survey, which include MDE and GAD sections (1253) (see Supplementary Table 1 for the characteristics of the 12-month follow-up sample). To correct the bias caused by lost to follow-up, inverse-probability weighting (IPW) (Seaman et al., 2012; Seaman and White, 2013) was calculated based on baseline covariates. Additionally, post-stratification weights were used to restore population distribution of gender, country of birth, and academic field within each university and across universities (further details available upon request).

For first-onset MDE-GAD, analyses were restricted to students reporting no history MDE-GAD at baseline ($n = 794$); for persistence, analyses were restricted to those with lifetime MDE-GAD at baseline ($n = 459$). Descriptive statistics are reported as weighted percentages and SE. Bivariate logistic regression analyses were performed to examine the association between hypothesized risk and protective factors and the outcome variable. Crude Odds Ratios (ORs) and MI-based 95% confidence intervals (CIs) were calculated, and statistical significance was assessed with F-test based on MI.

Multivariable logistic regression models were estimated for first-onset and persistence of MDE-GAD in the 12 months prior to follow-up. We identified the subset of variables that best explain first-onset and persistence of MDE-GAD using group lasso penalty (Tibshirani, 1996; Yuan and Lin, 2006) in the stacked imputed dataset (Zhao and Long, 2017), with gender forced into the model. We chose group lasso because this method has shown a good performance in terms of selection consistency (Bach and Orlitzky, 2008). Akaike information criterion (AIC) with effective degrees of freedom was used to select the regularization parameter λ (Harrell, 2001). The AIC criterion has shown to be an efficient selection method (Flynn et al., 2013) with lower computational cost than cross-validation (Gelman et al., 2014). Adjusted odds ratios (aORs) and MI-based CIs were obtained. Statistical significance was evaluated with two-side F test based on MI and α level of significance of 0.05. Firth's penalized likelihood estimation was applied to avoid overfitting due to data sparseness (Heinze, 2006). The area under the curve (AUC) was estimated to assess discriminant capacity of the models. To correct the over-estimation of prediction accuracy we used cross-validation with leave-one-out method (Efron and Gong, 1983).

Predicted probabilities were discretized into deciles and cross-classified with observed incident and persistent cases. Sensitivity was defined as the proportion of first-onset and persistent cases found among predefined proportions of respondents with the highest predicted probabilities as defined by the deciles. Positive predictive value was defined as the probability of first-onset MDE-GAD and of persistent MDE-GAD when estimated among respondents with the highest predicted probabilities based on each decile. MI was carried out using mice package from R (van Buuren and Groothuis-Oudshoorn, 2011). Analyses were performed using R v3.64 (RStudio Inc., 2017), SAS v9.4 (SAS Institute Inc., 2014) and Mplus v7.11 (Muthén and Muthén, 2015).

3. Results

3.1. 12-month first-onset and persistence of MDE-GAD

Fig. 1 shows the overall 12-month MDE-GAD prevalence, first-onset and persistence of MDE-GAD. A total of 24.9% of students had MDE-GAD at 12-month follow-up. Of students without a history of MDE-GAD at baseline, 13.3% (95% CI 10.6–15.9) had first-onset of MDE-GAD at 12-month follow-up. Of the students with MDE-GAD at baseline, 46.7% (95% CI 41.3–52.1) reported persistence at 12-month follow-up. No statistically significant differences by gender were found either in first-onset or persistence.

3.2. Risk and protective factors associated with the onset and persistence of MDE-GAD

Bivariate analyses revealed that most factors were statistically significantly associated with the onset and persistence of MDE-GAD, with the magnitude of associations being generally higher with the onset of MDE-GAD.

For first-onset of MDE-GAD, the strongest associations were childhood–adolescent emotional abuse or neglect (OR = 4.33; 95% CI 2.67–7.01), any childhood–adolescent adversity (OR = 4.08; 95% CI 2.40–6.94), current severe stress in any area of life (OR = 3.90; 95% CI 2.32–6.57) (Table 1); 12-month bipolar spectrum disorder (OR = 4.34; 95% CI 1.21–15.5), 12-month suicidal ideation (OR = 4.85; 95% CI 2.12–11.1) (Table 2); lifetime MDE symptoms (OR range = 2.33–3.63) and lifetime GAD symptoms (OR range = 2.15–3.75) (Table 3).

For persistence of MDE-GAD, the most important predictors were any 12-month stressful event (OR = 2.55; 95% CI 1.30–5.0) (Table 1); 12-month suicidal ideation (OR = 3.17; 95% CI 1.92–5.21) (Table 2); and lifetime GAD symptoms (OR range = 2.38–4.02) (Table 3).

Regarding protective factors examined, significant negative associations were found for perceived support with the onset of MDE-GAD (family: OR = 0.45; 95% CI 0.24–0.81, highest tertile and peers/others: OR = 0.30, 95% CI 0.16–0.58, highest tertile) and with persistence (peers/others: OR = 0.49; 95% CI 0.30–0.80, middle tertile and school: OR = 0.47, 95% CI 0.27–0.82, highest tertile) (Table 1). Also, positive coping strategies (OR = 0.94, 95% CI 0.89–0.99; OR = 0.93, 95% CI 0.89–0.97 for first-onset and persistence, respectively) and positive expectations (OR = 0.71, 95% CI 0.57–0.90; OR = 0.66, 95% CI 0.54–0.80 for first-onset and persistence, respectively) were found associated (Table 2).

As shown in Supplementary Table 2, sociodemographic and university-related variables did not significantly predict the onset of MDE-GAD. For persistence of MDE-GAD, only sexual orientation was associated.

3.3. Prediction models for the onset and persistence of MDE-GAD

The multivariable prediction models after lasso selection had a reasonable performance with a cross-validated AUC of 0.76 for predicting onset of MDE-GAD and 0.81 for predicting persistence. Table 4 shows cross-validated sensitivity and PPV estimates for different proportions of students at highest predicted risk based on the final baseline models. For instance, considering the 30% of students at highest baseline predicted risk for first-onset of MDE-GAD would include 74.9% (SE = 6.34) of all observed cases, being the probability of MDE-GAD onset of 33.4% (SE = 5.12) within this 30% of respondents. On the other hand, the 30% of students at highest baseline predicted risk of persistent MDE-GAD would include 54.6% (SE = 6.23) of all observed cases, with a probability of persistent MDE-GAD in this group of 85.1% (SE = 4.08).

The final prediction algorithm for first-onset of MDE-GAD (Supplementary Table 3) contained 21 variables in addition to gender. Independent predictors were: age of 18 years old (aOR = 0.5; 95% CI 0.3–1.0), childhood–adolescent emotional abuse or neglect (aOR = 3.2; 95% CI 1.7–5.7), current severe stress in any area of life (aOR = 3.0; 95% CI 1.7–5.3), a lifetime GAD symptom: worried about number of things or about everything (aOR = 2.4; 95% CI 1.4–4.0), a lifetime MDE symptom: sadness or depressed (aOR = 1.9; 95% CI 1.0–3.7), and neuroticism (aOR = 1.7; 95% CI 1.3–2.3).

The final prediction algorithm for persistence of MDE-GAD (Supplementary Table 4) contained 23 variables, in addition to gender, where the independent predictors were: family support (aOR = 2.8; 95% CI 1.3–5.9, highest tertile), a lifetime MDE symptom: diminished interest or pleasure (aOR = 2.1; 95% CI 1.2–3.9), 12-month suicidal ideation (aOR = 2.0; 95% CI 1.0–3.8), 3 lifetime GAD symptoms: more anxious, nervous or worried than other people (aOR = 1.9; 95% CI 1.1–3.6), worry or anxiety about things than other people wouldn't worry (aOR = 1.9; 95% CI 1.1–3.5) and restlessness or feeling keyed up or on edge (aOR = 2.2; 95% CI 1.2–4.2), neuroticism (aOR = 1.5; 95% CI 1.1–2.1) and openness to new experiences (aOR = 1.4; 95% CI 1.0–1.8).




Fig. 1. First-onset and persistence of MDE-GAD at 12-month follow-up by gender. (% weighted).

Table 1

Bivariate associations of childhood-adolescent experiences and recent stressful events with first-onset and persistence of MDE-GAD at 12-month follow-up.

	First-onset				Persistence			
			Bivariate model				Bivariate model	
	%	SE	OR	95%CI	%	SE	OR	95%CI
Childhood-Adolescent experiences								
Adversities								
Breakdown of family structure	16.5	1.6	1.16	(0.65–2.05)	23.7	2.3	1.28	(0.77–2.11)
Family maladaptation	30.9	1.9	2.14	(1.33–3.45)	46.6	2.9	1.74	(1.15–2.63)
Physical or sexual abuse	10.1	1.2	2.38	(1.26–4.50)	17.3	2.1	1.36	(0.78–2.38)
Emotional abuse or neglect	20.8	1.7	4.33	(2.67–7.01)	33.4	2.5	1.47	(0.95–2.28)
Dating violence	2.2	0.6	1.01	(0.24–4.32)	5.9	1.2	1.60	(0.67–3.86)
Bully victimization	28.0	1.9	1.55	(0.95–2.56)	50.8	2.6	2.06	(1.35–3.16)
Any adversity	54.6	2.0	4.08	(2.40–6.94)	75.2	2.5	1.61	(0.97–2.69)
Perceived support ^a								
Family (ref = Low)								
High	38.1	1.9	0.45	(0.24–0.81)	29.0	2.3	1.11	(0.67–1.85)
Middle	32.4	1.8	1.12	(0.66–1.90)	34.8	2.6	1.19	(0.72–1.98)
Peers/others (ref = Low)								
High	26.6	1.7	0.30	(0.16–0.58)	23.2	2.2	0.64	(0.38–1.07)
Middle	40.3	2.0	0.51	(0.30–0.84)	33.6	2.4	0.49	(0.30–0.80)
School (ref = Low)								
High	31.6	1.8	0.90	(0.50–1.59)	19.8	2.0	0.47	(0.27–0.82)
Middle	40.8	1.9	0.65	(0.36–1.17)	34.3	2.4	0.49	(0.30–0.79)
Recent stressful events								
12-month stressful events								
Death illness injury or accident	50.8	2.0	1.77	(1.12–2.79)	57.9	2.6	1.38	(0.90–2.10)
Breakup or betrayal arguments	49.5	2.0	1.62	(1.01–2.62)	69.7	2.4	1.85	(1.14–3.01)
Seriously physically or sexually assaulted	5.6	0.9	2.37	(1.10–5.12)	3.7	0.9	1.77	(0.61–5.14)
Trouble with the police or serious legal problem	6.2	1.0	1.19	(0.44–3.19)	3.4	1.1	3.80	(0.86–16.8)
Another stressful event	5.1	0.9	1.18	(0.46–3.01)	11.3	1.7	2.18	(1.08–4.39)
Any stressful event	77.7	1.7	1.52	(0.85–2.72)	87.0	1.8	2.55	(1.30–5.00)
Current stress								
Severe stress in any area of life	17.3	1.5	3.90	(2.32–6.57)	46.4	2.6	1.45	(0.95–2.21)

Significant odds ratios are shown in bold (Raw *p*-value statistically significant after adjustment for multiple comparisons using Benjamini-Hochberg procedure with false discovery rate 0.05). *P*-value obtained from F-test to evaluate significant difference in estimates based on multiple imputations.

% weighted. SE: Standard Error; OR: odds ratio; CI: confidence interval; ref.: Reference category.

^a Family: lowest tertile [1–3.75], middle tertile [3.75–4.5], highest tertile [4.5–5.0]; Peers/others: lowest tertile [1–2.75], middle tertile (2.75–3.5], highest tertile (3.5–5.0]; School: lowest tertile [1–3.33], middle tertile [3.33–4.17], highest tertile [4.17–5.0].

Table 2

Bivariate associations of psychological factors, impairment and mental health problems with first-onset and persistence of MDE-GAD at 12-month follow-up.

	First-onset				Persistence			
			Bivariate model				Bivariate model	
	Mean	SD	OR	95%CI	Mean	SD	OR	95%CI
Psychological factors								
Personality								
Extraversion	4.40	1.7	0.91	(0.77–1.07)	4.06	1.74	0.9	(0.79–1.02)
Agreeableness	4.34	1.0	0.77	(0.61–0.98)	4.27	0.94	1.09	(0.85–1.40)
Conscientiousness	5.30	1.4	0.89	(0.74–1.07)	5.02	1.18	0.77	(0.64–0.93)
Neuroticism	3.21	1.4	1.65	(1.37–1.98)	4.40	1.49	1.74	(1.45–2.07)
Openness to new experiences	5.23	1.3	1.11	(0.90–1.36)	5.11	1.41	1.31	(1.10–1.55)
Coping strategies								
Positive coping strategies	26.8	5.3	0.94	(0.89–0.99)	24.16	5.59	0.93	(0.89–0.97)
Student expectations								
Positive expectations	5.79	1.0	0.71	(0.57–0.90)	5.34	1.15	0.66	(0.54–0.80)
Impairment and mental health problems	%	SE			%	SE		
12-month role impairment								
Any severe role impairment	11.0	1.3	2.63	(1.44–4.81)	34.4	2.5	2.08	(1.34–3.23)
12-month mental health problems								
Bipolar spectrum disorder	2.1	0.6	4.34	(1.21–15.5)	7.1	1.4	2.26	(0.99–5.18)
Panic disorder	0.7	0.4	1.25	(0.11–14.7)	6.2	1.2	1.46	(0.63–3.38)
Alcohol abuse or dependence	5.0	0.9	0.38	(0.09–1.59)	3.9	1	0.67	(0.24–1.88)
Substance abuse or dependence	2.0	0.5	2.58	(0.71–9.37)	3.1	0.8	0.58	(0.17–1.98)
Adult Attention deficit hyperactivity disorder	7.7	1.1	2.31	(1.16–4.58)	23.7	2.2	1.92	(1.14–3.21)
Suicidal ideation	3.8	0.8	4.85	(2.12–11.1)	23.8	2.1	3.17	(1.92–5.21)

Significant odds ratios are shown in bold (Raw *p*-value statistically significant after adjustment for multiple comparisons using Benjamini-Hochberg procedure with false discovery rate 0.05). *P*-value obtained from F-test to evaluate significant difference in estimates based on multiple imputations.

% weighted; SD: Standard Deviation; SE: Standard Error; OR: odds ratio; CI: confidence interval; ref.: Reference category.

Table 3

Bivariate associations of lifetime symptoms of MDE and GAD with first-onset and persistence of MDE-GAD at 12-month follow-up.

	First-onset				Persistence			
			Bivariate model				Bivariate model	
	%	SE	OR	95%CI	%	SE	OR	95%CI
Lifetime symptoms								
Lifetime MDE symptoms								
Sadness or depressed	21.4	1.6	3.63	(2.24–5.88)	81.6	2.0	1.36	(0.80–2.32)
Discouraged	24.5	1.7	3.19	(2.00–5.10)	84.7	1.9	1.73	(0.96–3.13)
Diminished interest or pleasure	13.1	1.3	2.49	(1.39–4.49)	72.1	2.4	1.73	(1.09–2.76)
Worthlessness, feel down on yourself	16.0	1.4	2.33	(1.40–3.85)	73.6	2.2	1.91	(1.18–3.08)
Thoughts of death	4.6	0.9	1.65	(0.65–4.21)	35	2.5	1.58	(1.00–2.48)
Diminished concentration	6.8	1.0	1.92	(0.93–3.99)	50.8	2.7	1.78	(1.18–2.69)
Number of symptoms (ref = 0)								
1	16.5	1.4	2.53	(1.31–4.89)	3.5	1.0	0.34	(0.06–2.03)
2	13.7	1.3	5.13	(2.72–9.67)	6.4	1.3	1.89	(0.49–7.30)
3	11.2	1.3	6.19	(3.22–11.9)	17.2	2.0	1.10	(0.35–3.47)
4	2.3	0.6	4.08	(1.19–14.1)	28.3	2.3	1.16	(0.38–3.50)
5	0.0	0.0	n.a.	n.a.	26.3	2.2	2.04	(0.67–6.22)
6	0.0	0.0	n.a.	n.a.	14.3	1.8	4.24	(1.28–14.0)
Lifetime GAD symptoms								
Anxiety or nervousness	23.4	1.6	2.68	(1.67–4.32)	67.9	2.4	2.38	(1.49–3.80)
Worry about a number of different things or about everything	29.2	1.8	3.75	(2.37–5.95)	82	1.9	4.02	(2.18–7.43)
More anxious, nervous or worried than other people	7.4	1.0	1.12	(0.52–2.42)	61.5	2.5	3.13	(2.00–4.90)
Worry or anxiety about things other people wouldn't worry about	10.9	1.3	3.23	(1.81–5.76)	57.6	2.5	3.48	(2.23–5.43)
Difficulty to control the worry or anxiety	11.5	1.2	3.22	(1.84–5.62)	61.6	2.5	3.66	(2.29–5.87)
Restlessness or feeling keyed up or on edge	14.6	1.4	2.98	(1.76–5.06)	60.9	2.5	3.02	(1.91–4.76)
Irritability	17.2	1.5	2.15	(1.29–3.59)	68.4	2.5	1.36	(0.87–2.14)
Number of symptoms (ref = 0)								
1	14.4	1.4	9.63	(4.73–19.6)	4.9	1.1	2.13	(0.28–16.4)
2	11.3	1.3	3.28	(1.39–7.73)	6	1.2	2.59	(0.38–17.8)
3	11.6	1.2	4.97	(2.25–11.0)	7.7	1.4	5.0	(0.83–30.2)
4	6.7	1.0	11.7	(5.25–26.2)	10.8	1.9	10.9	(1.81–65.6)
5	3.3	0.7	22.4	(8.38–59.9)	15.2	1.9	8.22	(1.50–45.1)
6	1.4	0.5	14.8	(3.70–59.5)	16.6	1.9	14.4	(2.58–80.2)
7	0.0	0.0	n.a.	n.a.	18.7	2.0	13.3	(2.50–71.1)
8	0.0	0.0	n.a.	n.a.	14.7	1.9	31.3	(5.38–182.5)

Significant odds ratios are shown in bold (Raw p-value statistically significant after adjustment for multiple comparisons using Benjamini-Hochberg procedure with false discovery rate 0.05). P-value obtained from F-test to evaluate significant difference in estimates based on multiple imputations.

% weighted; SE: Standard Error; OR: odds ratio; CI: confidence interval; ref.: Reference category.

Table 4

Concentration of risk of MDE-GAD first-onset and persistence at 12-month follow-up in different proportions of university students.

% at highest predicted risk	First-onset				Persistence			
	Sensitivity ^a		PPV ^b		Sensitivity ^a		PPV ^b	
	%	SE	%	SE	%	SE	%	SE
100	100.0	0.00	13.3	2.01	100.0	0.00	46.7	3.73
90	99.6	0.58	14.8	2.19	99.1	0.70	51.4	3.92
80	97.8	1.45	16.3	2.44	97.8	1.08	57.1	4.06
70	94.7	2.28	18.1	2.71	94.3	2.08	62.9	4.19
60	91.2	3.32	20.3	3.10	86.9	3.01	67.6	4.38
50	86.4	4.49	23.1	3.51	80.5	3.75	75.3	4.28
40	81.8	5.22	27.4	4.24	69.9	4.70	81.7	4.04
30	74.9	6.34	33.4	5.12	54.6	6.23	85.1	4.08
20	64.9	7.57	43.4	6.59	37.5	6.31	87.6	4.73
10	45.5	9.04	60.7	8.58	19.2	6.33	89.5	6.84

SE: Standard error.

^a Sensitivity = proportion of first-onset or persistent MDE-GAD cases found among respondents at highest predicted risk, based on cross-validated predicted probabilities.

^b Positive Predictive Value (PPV) = probability of first-onset or persistent MDE-GAD when estimated among respondents at highest predicted risk, based on cross-validated predicted probabilities.

4. Discussion

4.1. Main findings

This study provides the first prospective data on first-onset and

persistence of MDE-GAD among Spanish university students during their first academic year. These disorders are common during university, 13.3% of students had a first-onset of MDE and/or GAD during their first year in university, and 46.7% of those with a history of these mental disorders before entering college had MDE and/or GAD at 12-month follow-up. We found a higher than expected incidence of MDE-GAD in our study than the 6.9% previously reported for MDE in Belgian university students (Ebert et al., 2019) and for GAD among young adults in Germany of 4.3% (Beesdo et al., 2010). Also our prevalence estimates were higher than reported from a sample of adolescents from Mexico of 8-year incidence of 12.9% and 1.6%, for MDE and GAD respectively (Benjet et al., 2016). Regarding persistence, these were also somewhat higher than reported in prior studies of university students (MDE: 26.9%; anxiety: 30.4%) but slightly lower than those found among adolescents (MDE: 70.2%, GAD: 51.2%) (Kessler et al., 2012; Zivin et al., 2009). However, these results are not directly comparable due to the aggregation of MDE and GAD in our study. In general, our findings suggest that these disorders are common during university period, which had been previously reported in cross-sectional studies (Auerbach et al., 2018; Ballester et al., 2020; Eisenberg et al., 2007), highlighting the importance of early detection of young people at risk of common mental disorders.

We also found that, while most of the childhood adversities included in the study were associated with first-onset of MDE-GAD, fewer were associated with persistence. This is consistent with previous findings of a reduced impact of stressful events on mental disorders according to the increase of the number of previous episodes (Kessler et al., 2008; McLaughlin et al., 2010). It could be explained through some kind of kindling phenomenon, understood as a progressive neurobiological

sensitization (Post, 2007). Also consistent with previous research, lifetime MDE and GAD symptoms were predictors of first-onset (Fava and Tossani, 2007; Horwath, 1992; Liu et al., 2019). These symptoms may correspond to prodromal stages or subthreshold disorders (Murphy et al., 1989; Wolitzky-Taylor et al., 2014). Thus, early detection of students with subclinical symptoms could facilitate preventive interventions before the onset of a mental disorder. Otherwise, we found that GAD symptoms predicted persistence of MDE-GAD more strongly than did MDE symptoms. This finding is in line with Kessler et al. (2008), reporting that while GAD predicted onset and persistence of MDE, MDE did not predict persistence of GAD. One possible explanation could be to interpret GAD as a prodrome or severity marker of MDE when they are considered together (Kessler et al., 2008; Wittchen et al., 2001). Nevertheless, Moffitt et al. (2007) found that the reverse pattern also occurs with almost the same frequency (Moffitt et al., 2007). Finally, protective factors were associated with the onset and persistence of MDE-GAD. Consistent with previous studies, family relationships, peer support and connection to school were protective (Galatzer-Levy et al., 2012; van Voorhees et al., 2008) as well as positive coping strategies (Crockett et al., 2007; McLafferty et al., 2019) and student expectations (Brackney and Karabenick, 1995; Riskind et al., 1987). These findings suggest that the identification of modifiable protective factors could aid prevention efforts, beyond a focus on the reduction of potential risk factors and which could contribute to resilience (Steinhardt and Dolbier, 2008).

Our findings show that prediction models applied to freshmen could be useful in identifying students at high risk of onset and persistence of Major Depressive Episode (MDE) and/or Generalized Anxiety Disorder (GAD) over their first year of school. Also, in comparison to previous integrative models of MDE or GAD (Ebert et al., 2019; King et al., 2011), this model jointly adds personality traits, protective factors and previous symptomatology to compose the model. The multivariate algorithm for onset had good performance ($AUC = 0.76$), similar than AUCs obtained previously for MDE among university students, general population and primary care ($AUC = 0.71$ – 0.82) (Bellón et al., 2011; Ebert et al., 2019; Nigatu et al., 2016); and slightly higher than AUC for GAD among general population and primary care ($AUC = 0.62$ – 0.75) (King et al., 2011; Nigatu and Wang, 2019). Less evidence exists on multivariate models for persistence of MDE and GAD (King et al., 2011; van Loo et al., 2015, 2014). Based in the models, may be defined 30% or greater as “high risk”, where over than 70% of MDE-GAD onset cases will occur and 55% of MDE-GAD persistent cases (an estimated of 79 students and 117 students respectively), which could be offered evidence-based interventions (Harrer et al., 2019; Rith-Najarian et al., 2019). However, the sensitivity could be increased, considering the prevalence of the persistence (46.7%), by targeting the top 50% at risk. Also on the other hand, the identification of false positives could be reduced by targeting the top 10% at risk (an estimate of 31 students for onset and 5 students for persistence). Notwithstanding, students at lower risk could benefit from awareness campaigns or universal prevention programs (Conley et al., 2015). Moreover, there is not yet evidence to support these threshold values to be used as intervention rules or the effectiveness of specific cut-off interventions, so a cost-effectiveness research should be conducted. Addressing these issues in future research will be important to enhance prediction tools on mental health in university settings.

4.2. Limitations

Several limitations should be taken into account when interpreting our results. First, a low response rate at baseline might have caused a biased representation of the total population of university students. Also, losses at 12-month follow-up were experienced in our study. These limitations were minimized by applying population-based adjustments through post stratification with a specific end-game strategy that resulted in the use of inverse-probability weights (Brick, 2013). Second, a convenience sample of universities was used; however, the

characteristics of the included universities were very similar to all Spanish universities. However, as described in the methods section, the characteristics of the included universities were very similar to all public Spanish universities, and therefore we are confident that the results are generalizable within this context. Third, the assessment of mental disorders was based on self-reports. However, a clinical reappraisal was carried out showing good concordance with blinded clinical diagnoses (Ballester et al., 2019). Fourth, our study does not differentiate between pure MDE and GAD cases or comorbid cases at baseline, which limits the analyses of specific effects in terms of onset and persistence of both conditions at follow-up. This is the first study that analyses the conjunction of MDE-GAD and future research in this sample will be necessary to study these groups more in-depth. Also, the analysis of MDE-GAD aggregated in our study limits the comparability with the incidence and persistence rates previously found in the literature. However, the use of MDE-GAD to develop a single algorithm could be a potential tool in the university setting for the early detection of a large population with symptoms of these common mental disorders and the subsequent implementation of preventive interventions addressing common underlying factors (Titov et al., 2011). Fifth, we divided the sample in two groups (first-onset and persistence) based on whether the students had previous history of MDE-GAD or not. We collapsed all cases with a baseline history of MDE-GAD as “persistent cases”, thus including a small proportion of recurrent cases (65 students had 12-month MDE-GAD at baseline). However, our definition of persistence is consistent with previous longitudinal studies among university students (Blasco et al., 2019; Mortier et al., 2017). Finally, the models were not validated with external samples. Leave-one-out method was used for cross-validation. Other methods, such as leave-one-site-out validation, could provide better evidence of the generalizability of the results. Because this method does not guarantee transportability to distinct samples, additional studies in other countries are necessary.

5. Conclusion

The current study provides relevant information on the high proportion of MDE-GAD among Spanish university students, highlighting the need to develop an active strategy for the detection and prevention of common mental disorders. University authorities should consider risk screening at university entrance to provide timely interventions or prevention strategies. The risk algorithms for MDE-GAD compared favorably with previous algorithms for prediction psychiatric outcomes and may be useful as prediction tools in university campuses.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jad.2021.10.135>.

Funding

This work was supported by Fondo de Investigación Sanitaria, Instituto de Salud Carlos III FEDER (J.A., grant number PI13/00343); Ministerio de Sanidad, Servicios Sociales e Igualdad, Plan Nacional Sobre Drogas PNSD (J.A., grant number exp.2015I015); and from the DIUE of the Generalitat de Catalunya (J.A., grant number 2017SGR452). Ministerio de Educación, Cultura y Deporte (L. B., FPU grant number FPU15/05728); and Instituto de Salud Carlos III (M. J. B., Río Hortega grant number CM14/00125; P. C., Sara Borrell grant number CD12/00440; P. M. Sara Borrell grant number CD18/00049).

Data availability statement

The data that support the findings of this study are available from the corresponding author, J.A., upon reasonable request.

CRediT authorship contribution statement

Jordi Alonso had full access to all of the data in this study and takes

responsibility for the integrity of the data, and the accuracy of the data analysis.

Study concept and design: Ballester, L., Alayo, I., Vilagut, G., Mortier, P., and Alonso, J.

Acquisition of data: Ballester, L., Vilagut, G., Alayo I., Almenara, J., Blasco, M.J., Barbaglia, G., Castellví, P., Cebrià, A.I., Echeburúa, E., Gabilondo, A., Gili, M., Iruin, A., Lagares, C., Miranda-Mendizabal, A., Parès-Badell, O., Pérez-Vázquez, M.T., Piqueras, J.A., Roca, M., Rodríguez-Marín, J., Sesé, A., Soto-Sanz, V., Vives, M., and Alonso, J.

Statistical analysis and interpretation of data: Vilagut, G., Alayo, I., Mortier, P., Ballester, L., Alonso, J.

Draft of the manuscript: Ballester, L.

Critical revision of the manuscript for important intellectual content: Vilagut, G., Alayo, I., Mortier, P., Almenara, J., Cebrià, A.I., Echeburúa, E., Gabilondo, A., Gili, M., Lagares, C., Piqueras, J.A., Roca, M., Soto-Sanz, V., Blasco, M.J., Castellví, P., Miranda-Mendizabal, A., Bruffaerts, R., Auerbach, R.P., Nock, M.K., Kessler, R.C., and Alonso, J.

Conflict of interest

In the past 3 years, Dr. Kessler was a consultant for Datastat, Inc., Holmusk, RallyPoint Networks, Inc., and Sage Therapeutics. Dr. Roca received research funds from Lundbeck and Janssen.”

Acknowledgements

We thank Roser Busquets and Mercedes Bayo for their continuous managerial assistance to this project, Gerbert Oliver for technological support and Arantxa Urdangarin for technical support.

References

- Arnett, J.J., 2000. Emerging adulthood: a theory of development from the late teens through the twenties. *Am. Psychol.* 55, 469–480. <https://doi.org/10.1037/0003-066X.55.5.469>.
- Auerbach, R.P., Alonso, J., Axinn, W.G., Cuijpers, P., Ebert, D., Green, J.G., Hwang, I., Kessler, R.C., Liu, H., Mortier, P., Nock, M.K., Pinder-Amaker, S., Sampson, N.A., Aguilar-Gaxiola, S., Al-Hamzawi, A., Andrade, L.H., Benjet, C., Caldas-de-Almeida, J. M., Demyttenaere, K., Florescu, S., de Girolamo, G., Gureje, O., Haro, J.M., Karam, E. G., Kiejna, A., Kovess-Masfety, V., Lee, S., McGrath, J.J., O'Neill, S., Pennell, B.-E., Scott, K., ten Have, M., Torres, Y., Zaslavsky, A.M., Zarkov, Z., Bruffaerts, R., 2016. Mental disorders among college students in the World Health Organization World Mental Health Surveys. *Psychol. Med.* 46, 2955–2970. <https://doi.org/10.1017/S0033291716001665>.
- Auerbach, R.P., Mortier, P., Bruffaerts, R., Alonso, J., Benjet, C., Cuijpers, P., Demyttenaere, K., Ebert, D.D., Green, J.G., Hasking, P., Murray, E., Nock, M.K., Pinder-Amaker, S., Sampson, N.A., Stein, D.J., Vilagut, G., Zaslavsky, A.M., Kessler, R.C., Collaborators, W.-I., 2018. The WHO World Mental Health Surveys International College Student project: prevalence and distribution of mental disorders. *J. Abnorm. Psychol.* 127, 623. <https://doi.org/10.1037/abn0000362>.
- Bach, F.R., Org, F.B., 2008. Consistency of the group lasso and multiple kernel learning. *J. Mach. Learn. Res.* 9, 1179–1225.
- Ballester, L., Alayo, I., Vilagut, G., Almenara, J., Cebrià, A.I., Echeburúa, E., Gabilondo, A., Gili, M., Lagares, C., Piqueras, J.A., Roca, M., Soto-Sanz, V., Blasco, M.J., Castellví, P., Forero, C.G., Bruffaerts, R., Mortier, P., Auerbach, R.P., Nock, M.K., Sampson, N., Kessler, R.C., Alonso, J., 2019. Accuracy of online survey assessment of mental disorders and suicidal thoughts and behaviors in Spanish university students. Results of the WHO world mental health- international college student initiative. *PLoS One* 14, e0221529. <https://doi.org/10.1371/journal.pone.0221529>.
- Ballester, L., Alayo, I., Vilagut, G., Almenara, J., Cebrià, A.I., Echeburúa, E., Gabilondo, A., Gili, M., Lagares, C., Piqueras, J.A., Roca, M., Soto-Sanz, V., Blasco, M.J., Castellví, P., Mortier, P., Bruffaerts, R., Auerbach, R.P., Nock, M.K., Kessler, R.C., Jordi, A., 2020. Mental disorders in Spanish university students: prevalence, age-of-onset, severe role impairment and mental health treatment. *J. Affect. Disord.* 273, 604–613. <https://doi.org/10.1016/j.jad.2020.04.050>.
- Barlow, D.H., Allen, L.B., Choate, M.L., 2016. Toward a unified treatment for emotional disorders – republished article. *Behav. Ther.* 47, 838–853. <https://doi.org/10.1016/j.beth.2016.11.005>.
- Beesdo, K., Pine, D.S., Lieb, R., Wittchen, H.U., 2010. Incidence and risk patterns of anxiety and depressive disorders and categorization of generalized anxiety disorder. *Arch. Gen. Psychiatry* 67, 47–57. <https://doi.org/10.1001/archgenpsychiatry.2009.177>.
- Bellón, J.A., de Dios Luna, J., King, M., Moreno-Küstner, B., Nazareth, I., Montón-Franco, C., GildeGómez-Barragán, M.J., Sánchez-Celaya, M., Díaz-Barreiros, M.A., Vicens, C., Cervilla, J.A., Švab, I., Maaroos, H.-I., Xavier, M., Geerlings, M.I., Saldívia, S., Gutiérrez, B., Motrico, E., Martínez-Cañavate, M.T., Oliván-Blázquez, B., Sánchez-Artiaga, M.S., March, S., del Mar Muñoz-García, M., Vázquez-Medrano, A., Moreno-Peral, P., Torres-González, F., 2011. Predicting the onset of major depression in primary care: international validation of a risk prediction algorithm from Spain. *Psychol. Med.* 41, 2075–2088. <https://doi.org/10.1017/S0033291711000468>.
- Benjet, C., Borges, G., Méndez, E., Albor, Y., Casanova, L., Orozco, R., Curiel, T., Fleiz, C., Medina-Mora, M.E., 2016. Eight-year incidence of psychiatric disorders and service use from adolescence to early adulthood: longitudinal follow-up of the Mexican Adolescent Mental Health Survey. *Eur. Child Adolesc. Psychiatry* 25, 163–173. <https://doi.org/10.1007/s00787-015-0721-5>.
- Bernstein, D.P., Ahluwalia, T., Pogge, D., Handelsman, L., 1997. Validity of the childhood trauma questionnaire in an adolescent psychiatric population. *J. Am. Acad. Child Adolesc. Psychiatry* 36, 340–348. <https://doi.org/10.1097/00004583-199703000-00012>.
- Blasco, M.J., Castellví, P., Almenara, J., Lagares, C., Roca, M., Sesé, A., Piqueras, J.A., Soto-Sanz, V., Rodríguez-Marín, J., Echeburúa, E., Gabilondo, A., Cebrià, A.I., Miranda-Mendizabal, A., Vilagut, G., Bruffaerts, R., Auerbach, R.P., Kessler, R.C., Alonso, J., 2016. Predictive models for suicidal thoughts and behaviors among Spanish university students: rationale and methods of the UNIVERSAL (University & mental health) project. *BMC Psychiatry* 16, 122. <https://doi.org/10.1186/s12888-016-0820-y>.
- Blasco, M.J., Vilagut, G., Alayo, I., Almenara, J., Cebrià, A.I., Echeburúa, E., Gabilondo, A., Gili, M., Lagares, C., Piqueras, J.A., Roca, M., Soto-Sanz, V., Ballester, L., Urdangarin, A., Bruffaerts, R., Mortier, P., Auerbach, R.P., Nock, M.K., Kessler, R.C., Alonso, J., 2019. First-onset and persistence of suicidal ideation in university students: a one-year follow-up study. *J. Affect. Disord.* 256, 192–204. <https://doi.org/10.1016/j.jad.2019.05.035>.
- Brackney, B.E., Karabenick, S.A., 1995. Psychopathology and academic performance: the role of motivation and learning strategies. *J. Couns. Psychol.* 42, 456–465. <https://doi.org/10.1037/0022-0167.42.4.456>.
- Bray, R.M., Pemberton, M.R., Hourani, L.L., Witt, M., Olmsted, K.L., Brown, J.M., Weimer, B., Lance, M.E., Marsden, M.E., Scheffler, S., 2009. Department of Defense Survey of Health Related Behaviors Among Active Duty Military Personnel. Research Triangle Institute, Research Triangle Park, North Carolina.
- Brick, J.M., 2013. Unit nonresponse and weighting adjustments: a critical review. *J. Off. Stat.* 29, 329–353. <https://doi.org/10.2478/jos-2013-0026>.
- Brugha, T.S., Cragg, D., 1990. The List of Threatening Experiences: the reliability and validity of a brief life events questionnaire. *Acta Psychiatr. Scand.* 82, 77–81. <https://doi.org/10.1111/j.1600-0447.1990.tb01360.x>.
- Conley, C.S., Durlak, J.A., Kirsch, A.C., 2015. In: A Meta-analysis of Universal Mental Health Prevention Programs for Higher Education Students, pp. 487–507. <https://doi.org/10.1007/s11211-015-0543-1>.
- Crockett, L.J., Iturbide, M.I., Torres Stone, R.A., McGinley, M., Raffaelli, M., Carlo, G., 2007. Acculturative stress, social support, and coping: relations to psychological adjustment among Mexican American college students. *Cult. Divers. Ethn. Minor. Psychol.* 13, 347–355. <https://doi.org/10.1037/1099-9809.13.4.347>.
- Dozois, D.J.A., Seeds, P.M., Collins, K.A., 2009. Transdiagnostic approaches to the prevention of depression and anxiety. *J. Cogn. Psychother.* 23, 44–59. <https://doi.org/10.1891/0889-8391.23.1.44>.
- Ebert, D.D., Buntrock, C., Mortier, P., Auerbach, R., Weisel, K.K., Kessler, R.C., Cuijpers, P., Green, J.G., Kiekens, G., Nock, M.K., Demyttenaere, K., Bruffaerts, R., 2019. Prediction of major depressive disorder onset in college students. *Depress. Anxiety* 36, 294–304. <https://doi.org/10.1002/da.22867>.
- Efron, B., Gong, G., 1983. A leisurely look at the bootstrap, the jackknife, and cross-validation. *Am. Stat.* 37, 36–48.
- Eisenberg, D., Gollust, S.E., Golberstein, E., Hefner, J.L., 2007. Prevalence and correlates of depression, anxiety, and suicidality among university students. *Am. J. Orthopsychiatry* 77, 534–542. <https://doi.org/10.1037/0002-9432.77.4.534>.
- Fava, G.A., Tossani, E., 2007. Prodromal stage of major depression. *Early Interv. Psychiatry* 1, 9–18. <https://doi.org/10.1111/j.1751-7893.2007.00005.x>.
- Felitti, V.J., Anda, R.F., Nordenberg, D., Williamson, D.F., Spitz, A.M., Edwards, V., Koss, M.P., Marks, J.S., 1998. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. *Am. J. Prev. Med.* 14, 245–258. [https://doi.org/10.1016/S0749-3797\(98\)00017-8](https://doi.org/10.1016/S0749-3797(98)00017-8).
- Flynn, C., Hurvich, C., Simonoff, J., 2013. Efficiency for regularization parameter selection in penalized likelihood estimation of misspecified models on JSTOR. *J. Am. Stat. Assoc.* 108, 1031–1043.
- Galatzer-Levy, I.R., Burton, C.L., Bonanno, G.A., 2012. Coping flexibility, potentially traumatic life events, and resilience: a prospective study of college student adjustment. *J. Soc. Clin. Psychol.* 31, 542–567. <https://doi.org/10.1521/jscp.2012.31.6.542>.
- Garber, J., Weersing, V.R., 2010. Comorbidity of anxiety and depression in youth: implications for treatment and prevention. *Clin. Psychol. Sci. Pract.* 17, 293–306. <https://doi.org/10.1111/j.1468-2850.2010.01221.x>.
- Gelman, A., Hwang, J., Vehtari, A., 2014. Understanding predictive information criteria for Bayesian models. *Stat. Comput.* 24, 997–1016. <https://doi.org/10.1007/s11222-013-9416-2>.
- Goldberg, D.P., Krueger, R.F., Andrews, G., Hobbs, M.J., 2009. Emotional disorders: Cluster 4 of the proposed meta-structure for DSM-V and ICD-11. *Psychol. Med.* 39, 2043–2059. <https://doi.org/10.1017/S0033291709990298>.
- Goodenow, C., 1993. The psychological sense of school membership among adolescents: scale development and educational correlates. *Psychol. Sch.* 30, 79–90. [https://doi.org/10.1002/1520-6807\(199301\)30:1<79::AID-PITS2310300113>3.0.CO;2-X](https://doi.org/10.1002/1520-6807(199301)30:1<79::AID-PITS2310300113>3.0.CO;2-X).
- Gorman, J., 1996. Comorbid anxiety and depression spectrum disorders. *Depress. Anxiety* 4, 160–168.

- Gosling, S.D., Rentfrow, P.J., Swann, W.B., 2003. A very brief measure of the Big-Five personality domains. *J. Res. Pers.* 37, 504–528. [https://doi.org/10.1016/S0092-6566\(03\)00046-1](https://doi.org/10.1016/S0092-6566(03)00046-1).
- Harrell, F.E., 2001. *Regression Modeling Strategies: With Applications to Linear Models, Logistic Regression, And Survival Analysis, Graduate Texts in Mathematics*. Springer.
- Harrer, M., Adam, S.H., Baumeister, H., Cuijpers, P., Karyotaki, E., Auerbach, R.P., Kessler, R.C., Bruffaerts, R., Berkling, M., Ebert, D.D., 2019. Internet interventions for mental health in university students: a systematic review and meta-analysis. *Int. J. Methods Psychiatr. Res.* 28, 1–18. <https://doi.org/10.1002/MP.1759>.
- Heinze, G., 2006. A comparative investigation of methods for logistic regression with separated or nearly separated data. *Stat. Med.* 25, 4216–4226.
- Horwath, E., 1992. Depressive symptoms as relative and attributable risk factors for first-onset major depression. *Arch. Gen. Psychiatry* 49, 817. <https://doi.org/10.1001/archpsyc.1992.01820100061011>.
- Ibrahim, A.K., Kelly, S.J., Adams, C.E., Glazebrook, C., 2013. A systematic review of studies of depression prevalence in university students. *J. Psychiatr. Res.* 47, 391–400. <https://doi.org/10.1016/j.jpsychires.2012.11.015>.
- International Test Commission, 2019. International Test Commission Guidelines [WWW Document]. <https://www.intestcom.org/page/5>.
- Kessler, R.C., Üstün, T.B., 2004. The World Mental Health (WMH) Survey Initiative version of the World Health Organization (WHO) Composite International Diagnostic Interview (CIDI). *Int. J. Methods Psychiatr. Res.* 13, 93–121. <https://doi.org/10.1002/MP.168>.
- Kessler, R.C., Andrade, L.H., Bijl, R.V., Offord, D.R., Demler, O.V., Stein, D.J., 2002. The effects of co-morbidity on the onset and persistence of generalized anxiety disorder in the ICPE surveys. *Psychol. Med.* 32, 1213–1225. <https://doi.org/10.1017/S0033291702006104>.
- Kessler, R.C., Adler, L., Ames, M., Demler, O., Faraone, S., Hiripi, E., Howes, M.J., Jin, R., Secnik, K., Spencer, T., Ustun, T.B., Walters, E.E., 2005. The World Health Organization adult ADHD self-report scale (ASRS): a short screening scale for use in the general population. *Psychol. Med.* 35, 245–256.
- Kessler, R.C., Galea, S., Jones, R.T., Parker, H.A., Hurricane Katrina Community Advisory Group, 2006. Mental illness and suicidality after Hurricane Katrina. *Bull. World Health Organ.* 84, 930–939. <https://doi.org/10.2471/BLT.06.033019>.
- Kessler, R.C., Gruber, M., Hettema, J.M., Hwang, I., Sampson, N., Yonkers, K.A., 2008. Co-morbid major depression and generalized anxiety disorders in the National Comorbidity Survey follow-up. *Psychol. Med.* 38, 365–374. <https://doi.org/10.1017/S0033291707002012>.
- Kessler, R.C., Avenevoli, S., Costello, E.J., Georgiades, K., Green, J.G., Gruber, M.J., He, J.P., Koretz, D., McLaughlin, K.A., Petukhova, M.V., Sampson, N.A., Zaslavsky, A.M., Merikangas, K.R., Costello, J.E., Georgiades, K., Greif Green, J., Gruber, M.J., He, J.P., Koretz, D., McLaughlin, K.A., Petukhova, M.V., Sampson, N.A., Zaslavsky, A.M., Merikangas, K.R., 2012. Prevalence, persistence, and sociodemographic correlates of DSM-IV disorders in the National Comorbidity Survey Replication Adolescent Supplement. *Arch. Gen. Psychiatry* 69, 372. <https://doi.org/10.1001/archgenpsychiatry.2011.160>.
- Kessler, R.C., Calabrese, J.R., Farley, P.A., Gruber, M.J., Jewell, M.A., Katon, W., Keck, P. E., Nierenberg, A.A., Sampson, N.A., Shear, M.K., Shillington, A.C., Stein, M.B., Thase, M.E., Wittchen, H.-U., 2013a. Composite International Diagnostic Interview screening scales for DSM-IV anxiety and mood disorders. *Psychol. Med.* 43, 1625–1637. <https://doi.org/10.1017/S0033291712002334>.
- Kessler, R.C., Santiago, P.N., Colpe, L.J., Dempsey, C.L., First, M.B., Heeringa, S.G., Stein, M.B., Fullerton, C.S., Gruber, M.J., Naifeh, J.A., Nock, M.K., Sampson, N.A., Schoenbaum, M., Zaslavsky, A.M., Ursano, R.J., 2013b. Clinical reappraisal of the Composite International Diagnostic Interview Screening Scales (CIDI-SC) in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS). *Int. J. Methods Psychiatr. Res.* 22, 303–321. <https://doi.org/10.1002/MP.1398>.
- King, M., Bottomley, C., Bellón-Saamen, J.A., Torres-Gonzalez, F., Švab, I., Rifel, J., Maaros, H.I., Aluoja, A., Geerlings, M.I., Xavier, M., Carraa, I., Vicente, B., Saldivia, S., Nazareth, I., 2011. An international risk prediction algorithm for the onset of generalized anxiety and panic syndromes in general practice attendees: PredictA. *Psychol. Med.* 41, 1625–1639. <https://doi.org/10.1017/S0033291710002400>.
- Kotov, R., 2011. New dimensions in the quantitative classification of mental illness. *Arch. Gen. Psychiatry* 68, 1003. <https://doi.org/10.1001/archgenpsychiatry.2011.107>.
- Leon, A.C., Olsson, M., Portera, L., Farber, L., Sheehan, D.V., 1997. Assessing psychiatric impairment in primary care with the Sheehan Disability Scale. *Int. J. Psychiatry Med.* 27, 93–105. <https://doi.org/10.2190/t8em-c8yh-373n-1uwd>.
- Liu, Y., Zhang, N., Bao, G., Huang, Y., Ji, B., Wu, Y., Liu, C., Li, G., 2019. Predictors of depressive symptoms in college students: a systematic review and meta-analysis of cohort studies. *J. Affect. Disord.* 244, 196–208. <https://doi.org/10.1016/j.jad.2018.10.084>.
- McEvoy, P.M., Nathan, P., Norton, P.J., 2009. Efficacy of transdiagnostic treatments: a review of published outcome studies and future research directions. *J. Cogn. Psychother.* 23, 20–33. <https://doi.org/10.1891/0889-8391.23.1.20>.
- McLafferty, M., Armour, C., Bunting, B., Ennis, E., Lapsley, C., Murray, E., O'Neill, S., 2019. Coping, stress, and negative childhood experiences: the link to psychopathology, self-harm, and suicidal behavior. *Psych J.* 8, 293–306. <https://doi.org/10.1002/pchj.301>.
- McLaughlin, K.A., Green, J.G., Gruber, M.J., Sampson, N.A., Zaslavsky, A.M., Kessler, R. C., 2010. Childhood adversities and adult psychiatric disorders in the National Comorbidity Survey Replication II. *Arch. Gen. Psychiatry* 67, 124. <https://doi.org/10.1001/archgenpsychiatry.2009.187>.
- Moffitt, T.E., Harrington, H., Caspi, A., Kim-Cohen, J., Goldberg, D., Gregory, A.M., Poulton, R., 2007. Depression and generalized anxiety disorder. *Arch. Gen. Psychiatry* 64, 651. <https://doi.org/10.1001/archpsyc.64.6.651>.
- Mortier, P., Kiekens, G., Auerbach, R.P., Cuijpers, P., Demyttenaere, K., Green, J.G., Kessler, R.C., Nock, M.K., Zaslavsky, A.M., Bruffaerts, R., 2017. A risk algorithm for the persistence of suicidal thoughts and behaviors during college. *J. Clin. Psychiatry* 78, e828–e836. <https://doi.org/10.4088/JCP.17m11485>.
- Murphy, J.M., Sobol, A.M., Olivier, D.C., Monson, R.R., Leighton, A.H., Pratt, L.A., 1989. Prodromes of depression and anxiety. The Stirling County study. *Br. J. Psychiatry* 155, 490–495. <https://doi.org/10.1192/bjp.155.4.490>.
- Muthén, L., Muthén, B., 2015. *Mplus, Seventh edition*. Muthén Muthén, Los Angeles, CA.
- Newby, J.M., McKinnon, A., Kuyken, W., Gilbody, S., Dalgleish, T., 2015. Systematic review and meta-analysis of transdiagnostic psychological treatments for anxiety and depressive disorders in adulthood. *Clin. Psychol. Rev.* 40, 91–110. <https://doi.org/10.1016/j.cpr.2015.06.002>.
- Nigatu, Y.T., Wang, J.L., 2019. External validation of the International Risk Prediction Algorithm for the onset of generalized anxiety and/or panic syndromes (The Predict A) in the US general population. *J. Anxiety Disord.* 64, 40–44. <https://doi.org/10.1016/j.janxdis.2019.03.004>.
- Nigatu, Y.T., Liu, Y., Wang, J.L., 2016. External validation of the international risk prediction algorithm for major depressive episode in the US general population: the PredictD-US study. *BMC Psychiatry* 16, 1–8. <https://doi.org/10.1186/s12888-016-0971-x>.
- Nock, M.K., Holmberg, E.B., Photos, V.I., Michel, B.D., 2007. Self-Injurious Thoughts and Behaviors Interview: development, reliability, and validity in an adolescent sample. *Psychol. Assess.* 19, 309–317. <https://doi.org/10.1037/1040-3590.19.3.309>.
- Nolen-Hoeksema, S., Watkins, E.R., 2011. A heuristic for developing transdiagnostic models of psychopathology: explaining multifinality and divergent trajectories. *Perspect. Psychol. Sci.* 6, 589–609. <https://doi.org/10.1177/1745691611419672>.
- Posner, K., Brown, G.K., Stanley, B., Brent, D.A., Yershova, K.V., Oquendo, M.A., Currier, G.W., Melvin, G.A., Greenhill, L., Shen, S., Mann, J.J., 2011. The Columbia-Suicide Severity Rating Scale: initial validity and internal consistency findings from three multisite studies with adolescents and adults. *Am. J. Psychiatry* 168, 1266–1277. <https://doi.org/10.1176/appi.ajp.2011.10111704>.
- Post, R.M., 2007. Kindling and sensitization as models for affective episode recurrence, cyclicity, and tolerance phenomena. *Neurosci. Biobehav. Rev.* 31, 858–873. <https://doi.org/10.1016/j.neubiorev.2007.04.003>.
- Riskind, J.H., Rholes, W.S., Brannon, A.M., Burdick, C.A., 1987. Attributions and expectations: a confluence of vulnerabilities in mild depression in a college student population. *J. Pers. Soc. Psychol.* 53, 349–354. <https://doi.org/10.1037/0022-3514.53.2.349>.
- Rith-Najarian, L.R., Boustani, M.M., Chorpita, B.F., 2019. A systematic review of prevention programs targeting depression, anxiety, and stress in university students. *J. Affect. Disord.* 257, 568–584. <https://doi.org/10.1016/j.jad.2019.06.035>.
- Round, A., 2005. *A Survey of Student Attitudes, Experiences And Expectations on Selected Vocational Courses at the University of Northumbria*. University of Northumbria.
- RStudio Inc., 2017. *RStudio Software 1.1.383*.
- Rueter, M.A., Scaramella, L., Wallace, L.E., Conger, R.D., 1999. First onset of depressive or anxiety disorders predicted by the longitudinal course of internalizing symptoms and parent-adolescent disagreements. *Arch. Gen. Psychiatry* 56, 726–732. <https://doi.org/10.1001/archpsyc.56.8.726>.
- SAS Institute Inc, 2014. *SAS Software 9.4*.
- Saunders, J.B., Aasland, O.G., Babor, T.F., De la Fuente, J.R., Grant, M., 1993. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption-II. *Addiction* 88, 791–804. <https://doi.org/10.1111/j.1360-0443.1993.tb02093.x>.
- Seaman, S.R., White, I.R., 2013. Review of inverse probability weighting for dealing with missing data. *Stat. Methods Med. Res.* 22, 278–295. <https://doi.org/10.1177/0962280210395740>.
- Seaman, S.R., White, I.R., Copas, A.J., Li, L., 2012. Combining multiple imputation and inverse-probability weighting. *Biometrics* 68, 129–137. <https://doi.org/10.1111/j.1541-0420.2011.01666.x>.
- Spinhoven, P., Elzinga, B.M., Hovens, J.G., Roelofs, K., Zitman, F.G., van Oppen, P., Penninx, B.W., 2010. The specificity of childhood adversities and negative life events across the life span to anxiety and depressive disorders. *J. Affect. Disord.* 126, 103–112. <https://doi.org/10.1016/j.jad.2010.02.132>.
- Steinhardt, M., Dolbier, C., 2008. Evaluation of a resilience intervention to enhance coping strategies and protective factors and decrease symptomatology. *J. Am. Coll. Health* 56, 445–453. <https://doi.org/10.3200/JACH.56.4.445-454>.
- Swearer, S.M., Cary, P.T., 2003. Perceptions and attitudes toward bullying in middle school youth. *J. Appl. Sch. Psychol.* 19, 63–79. <https://doi.org/10.1300/J008v19n02.05>.
- Tibshirani, R., 1996. Regression shrinkage and selection via the lasso. *J. R. Stat. Soc. Ser. B* 58, 267–288.
- Titov, N., Dear, B.F., Schwencke, G., Andrews, G., Johnston, L., Craske, M.G., McEvoy, P., 2011. Transdiagnostic internet treatment for anxiety and depression: a randomised controlled trial. *Behav. Res. Ther.* 49, 441–452. <https://doi.org/10.1016/j.brat.2011.03.007>.
- Ursano, R., 2012. *Modifiable Risk And Protective Factors for Suicidal Behaviors in the US Army*. National Institute of Mental Health, Bethesda, MD.
- van Buuren, S., 2012. Flexible Imputation of Missing Data. In: *CRC Interdisciplinary Statistics Series*. Chapman and Hall/CRC. <https://doi.org/10.1201/b11826>.
- van Buuren, S., Groothuis-Oudshoorn, K., 2011. MICE: multivariate imputation by chained equations in R. *J. Stat. Softw.* 45, 1–68.

- van Loo, H.M., Cai, T., Gruber, M.J., Li, J., de Jonge, P., Petukhova, M., Rose, S., Sampson, N.A., Schoevers, R.A., Wardenaar, K.J., Wilcox, M.A., Al-Hamzawi, A.O., Andrade, L.H., Bromet, E.J., Bunting, B., Fayyad, J., Florescu, S.E., Gureje, O., Hu, C., Huang, Y., Levinson, D., Medina-Mora, M.E., Nakane, Y., Posada-Villa, J., Scott, K.M., Xavier, M., Zarkov, Z., Kessler, R.C., 2014. Major depressive disorder subtypes to predict long-term course. *Depress. Anxiety* 31, 765–777. <https://doi.org/10.1002/da.22233>.
- van Loo, H.M., Aggen, S.H., Gardner, C.O., Kendler, K.S., 2015. Multiple risk factors predict recurrence of major depressive disorder in women. *J. Affect. Disord.* 180, 52–61. <https://doi.org/10.1016/j.jad.2015.03.045>.
- van Voorhees, B.W., Paunesku, D., Kuwabara, S.A., Basu, A., Gollan, J., Hankin, B.L., Melkonian, S., Reinecke, M., 2008. Protective and vulnerability factors predicting new-onset depressive episode in a representative of U.S. adolescents. *J. Adolesc. Health* 42, 605–616. <https://doi.org/10.1016/j.jadohealth.2007.11.135>.
- Vázquez, F.L., Torres, Á., Otero, P., Díaz, O., 2011. Prevalence, comorbidity, and correlates of DSM-IV axis I mental disorders among female university students. *J. Nerv. Ment. Dis.* 199, 379–383. <https://doi.org/10.1097/NMD.0b013e31821cd29c>.
- Vogt, D.S., Proctor, S.P., King, D.W., King, L.A., Vasterling, J.J., 2008. Validation of scales from the deployment risk and resilience inventory in a sample of operation Iraqi freedom veterans. *Assessment* 15, 391–403. <https://doi.org/10.1177/1073191108316030>.
- Wang, J.L., Sareen, J., Patten, S., Bolton, J., Schmitz, N., Birney, A., 2014. A prediction algorithm for first onset of major depression in the general population: development and validation. *J. Epidemiol. Community Health* 68, 418–424. <https://doi.org/10.1136/jech-2013-202845>.
- White, I.R., Royston, P., Wood, A.M., 2011. Multiple imputation using chained equations: issues and guidance for practice. *Stat. Med.* 30, 377–399. <https://doi.org/10.1002/sim.4067>.
- WHM-ICS, 2015. The WHO World Mental Health Surveys International College Student Project (WMH-ICS) [WWW Document]. URL: https://www.hcp.med.harvard.edu/wmh/college_student_survey.php (accessed 5.29.20).
- Wittchen, H.U., Hoyer, J., Friis, R., 2001. Generalized anxiety disorder - a risk factor for depression? *Int. J. Methods Psychiatr. Res.* 10, 52–57. <https://doi.org/10.1002/mpr.100>.
- Wolitzky-Taylor, K., Dour, H., Zinbarg, R., Mineka, S., Vrshek-Schallhorn, S., Epstein, A., Bobova, L., Griffith, J., Waters, A., Nazarian, M., Rose, R., Craske, M.G., 2014. Experiencing core symptoms of anxiety and unipolar mood disorders in late adolescence predicts disorder onset in early adulthood. *Depress. Anxiety* 31, 207–213. <https://doi.org/10.1002/da.22250>.
- Yuan, M., Lin, Y., 2006. Model selection and estimation in regression with grouped variables. *J. R. Stat. Soc. Ser. B (Stat. Methodol.)* 68, 49–67. <https://doi.org/10.1111/j.1467-9868.2005.00532.x>.
- Zhao, Y., Long, Q., 2017. Variable selection in the presence of missing data: imputation-based methods. *Wiley Interdiscip. Rev. Comput. Stat.* 9, e1402 <https://doi.org/10.1002/wics.1402>.
- Zivin, K., Eisenberg, D., Gollust, S.E., Golberstein, E., 2009. Persistence of mental health problems and needs in a college student population. *J. Affect. Disord.* 117, 180–185. <https://doi.org/10.1016/j.jad.2009.01.001>.