

Valorization of Wild Edible Plants as Food Ingredients and Their Economic Value

Jesús Clemente-Villalba ¹, Francisco Burló ¹, Francisca Hernández ²  and Ángel A. Carbonell-Barrachina ^{1,*} 

¹ Research Group “Food Quality and Safety”, Centro de Investigación e Innovación Agroalimentaria y Agroambiental (CIAGRO-UMH), Miguel Hernández University, 03312 Orihuela, Spain

² Grupo de Investigación en Fruticultura y Técnicas de Producción, Centro de Investigación e Innovación Agroalimentaria y Agroambiental (CIAGRO-UMH), Miguel Hernández University, 03312 Orihuela, Spain

* Correspondence: angel.carbonell@umh.es; Tel.: +34-96-674-97-54

Abstract: (1) Background: Wild Edible Plants (WEPs) are plants that grow without human help, by simply using the available resources. These types of plants are undervalued, because there is a lack of knowledge about their bioactive composition and nutritional/functional potential. (2) Scope and Approach: The main aim of this review is to fully identify the potential uses and importance of WEPs in certain regions based on (i) their sustainability, because they grow with their own resources, (ii) their content of bioactive compounds and consequently nutritional and functional value, (iii) their socio-economic relevance, and (iv) their ability to be useful in the agri-food industry in the short term. (3) Results: This review found evidence that a consumption of between 100 and 200 g of some of these WEPs can cover up to 50% of the recommended daily intake of proteins and fiber, being also a natural source of macro- and micro-minerals. Regarding their bioactive composition, most of these plants contain phenolic compounds and flavonoids, which determine their antioxidant capacity. (4) Conclusions: These reported results clearly demonstrate the high potential of the WEPs from a nutritional, economic and social point of view; although further studies are needed to gather deeper scientific information about their potential role in the socio-economic sustainability of specific groups of farmers worldwide.

Keywords: antioxidants; edible flowers; economic value; minerals; principal component analysis; sugars; total phenol content; flavonoids



Citation: Clemente-Villalba, J.; Burló, F.; Hernández, F.; Carbonell-Barrachina, Á.A. Valorization of Wild Edible Plants as Food Ingredients and Their Economic Value. *Foods* **2023**, *12*, 1012. <https://doi.org/10.3390/foods12051012>

Academic Editors: Luís Catarino, Maria Romeiras and Ângela Fernandes

Received: 7 February 2023
Revised: 20 February 2023
Accepted: 23 February 2023
Published: 27 February 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

“Wild Edibles” is a term used to describe both plants and animals consumed by humans. In 1999, the Food and Agriculture Organization of the United Nations (FAO) described the term wild plants as “those that grow spontaneously in self-maintaining populations in natural or semi-natural ecosystems and can exist independently of direct human action” [1]. The FAO estimated in 2016 that as much as ~100 million people consumed wild edible plants (WEPs) in Europe [2]; this figure highlights the potential of these plants even currently. The consumption of this type of plants dates back to the Bronze Age as shown by the remains found in a site in Peñalosa (Jaén, Spain); more than 50 species were identified in this site, including *Rumex* sp. and *Calendula* sp. The conclusion of this ethnobotanical study was that these plants were used as food and/or as flavoring additives [3]. The importance of these plants is evident from multiple studies conducted worldwide, such as in Brazil [4], China [5], Ethiopia [6], Guatemala [7], Iceland [8], India [9], Japan [10], and Tunisia [11]. In Europe, Schulp et al. [12] wrote a review on the identification of WEPs throughout Europe, finding them in 17 countries. In fact, there are many studies focusing on the Mediterranean region because it has a great diversity of WEPs, especially in Greece, Italy, Portugal and Spain. In all of these studies, nutritional potential or bioactive profiles were reported [2,13–17]. Despite all the studies carried out over the last 10 years, the full potential of WEPs has still not been fully reached and this is a hot topic that deserves

deeper attention by the scientific community considering especially the economy and their role in the sustainability of rural areas.

These plants may play an important role in environmental sustainability as they grow wild and can be used as a functional ingredient to develop new food products. This sustainable character is persuading more and more consumers, chefs and nutritionists to introduce WEPs in their dishes and the diets they prescribe. The relevance of this review is supported by the growing global demand for a change in eating habits, where key new trends are essential and include: (i) reduction in gasses' emissions, (ii) growth of sustainable crops, and (iii) greater environmental awareness. An example of this change and this new trend is the guide that FAO published in 2017 on wild food plants, or current campaigns for the consumption of edible insects as a new and sustainable source of protein [18]. The future must be more sustainable and WEPs can make a significant contribution to this change.

2. Scientific Literature Review

To identify interesting scientific publications dealing with the composition and relevance of WEPs, this review was based on the 2020 update of the PRISMA approach [19]. The literature was searched in different databases: (i) Scopus, (ii) FSTA and (iii) ScienceDirect; the keywords used were the following: "wild edible plants", "WEP", "edible plants", "ruderal plants", and "wild edible plants food". Most of the articles that were selected (1999 to 2022) were included in the Journal Citation Reports (JCR). The selection process is shown in Figure 1. The review is structured in different sections: (i) proximate characterization, (ii) sugars and organic acids, (iii) mineral content, (iv) fatty acids, (v) phenolic content and flavonoids, and (vi) economic value of WEPs.




Figure 1. Diagram of selection of scientific publications.

3. Wild Edible Plants, WEPs

This review collects information on 115 WEPs (Table S1, Supplementary Material) located around the world and belonging to 47 different families. The families with the highest representation were Asteraceae > Brassicaceae > Fabaceae > Lamiaceae.

4. Proximate Characterization

Proximate analysis is used in foods to estimate the values of energy, moisture, protein, lipids, water, ash, and carbohydrate in the samples under study. Proteins, lipids and carbohydrates contribute to the total energy content in an organism, while ash and water also contribute to the organism mass [20]. Therefore, these determinations are essential for correct nutritional labelling, whose main objective is to provide data on macro- and micro-nutrients [21]. Nutritional labelling is regulated in the United States by the Food and Drugs Administration (FDA) and in the European Union by the European Food Safety Authority (EFSA). The main values reflected on food labels are: energy, protein, total fat, saturated fat, total carbohydrate, total sugars and sodium; all are expressed in g, mg or µg per 100 g [22].

The different parameters of the proximate characterization were compiled and summarized for 47 WEPs. The families with the highest representation in the proximate characterization were Asteraceae (9) > Lamiaceae (5) > Polygonaceae (4) (Table 1).

The water content or **moisture** is one of the most important parameters in a plant. For a plant to produce 1 kg of organic matter, it needs to absorb 500 kg of water, which is subsequently eliminated by different processes (e.g., transpiration or evaporation) [23]. In herbaceous plants, the moisture content usually reaches ~90% of the fresh weight, and only on rare occasions (such as intense water stress conditions) it is below 70%. The functions of the water content in the plant are essential, because it maintains cell turgor, facilitates the transport of solutes through the plant, participates in the reduction of CO₂ through photosynthesis, and even in the cooling of the leaves during hot hours [24]. The average moisture content in the WEPs averaged ~80%. It should be noted that, out of the 115 WEP reviewed, the highest content was found in *Silybum marianum* (Asteraceae family) when it reached 93.4 g per 100 g [25,26]; on the contrary, the plant with the lowest moisture content was *Thymus pulegioides* (Lamiaceae family) with 47.6 g per 100 g [27]. In WEPs of the Lamiaceae family, it was observed that moisture values remained between 47.6 and 73.0 g per 100 g; these values were below the average (Table 1).

Studies dealing with 45 out of the 115 studied WEPs (from 24 families), provided results on ash contents. The two WEPs that showed the highest ash contents were: *Blumea lacera* (Asteraceae family) and *Hygrophila schulli* (Amaranthaceae family) with values of 24.05% and 23.36%, respectively [28]. On the contrary, the family that showed the lowest ash values was the Polygonaceae, which included four plants (*Rumex acetosella*, *Rumex induratus*, *Rumex papillaris*, and *Rumex pulcher*) with values of 1.2, 1.0, 1.0 and 1.9 g per 100 g, respectively [29,30].

Table 1. Proximate composition reported in wild edible plants, WEPs.

| Plant Species | Part of Plant | Unit | Moisture | Ash | Proteins | Fat | Carbohydrates | Fibre | Energy [§] | Reference |
|--|---------------|-------------------------|---------------------|---------------------|---------------------|-----------------------|----------------------|---------------------|---------------------|-----------|
| <i>Asystasia gangetica</i> (L.) T. Anderson | - | % | 70.21 ± 0.98 | 17.35 ± 0.26 | 7.84 ± 0.12 | 2.04 ± 0.03 | 10.63 ± 0.23 | 8.14 ± 0.55 | 92.27 ± 0.27 | [31] |
| <i>Achyranthes aspera</i> L. | - | % | 53.34 ± 0.58 | 23.26 ± 0.65 | 12.60 ± 0.11 | 1.196 ± 0.01 | 14.35 ± 0.14 | 16.89 ± 0.34 | 118.62 ± 0.06 | [31] |
| <i>Allium ampeloprasum</i> L. | Bulbs | g/100 g | 78.3 (76.0–81.5) | 0.8 (0.5–1.0) | 1.7 (1.2–2.0) | 0.18 (0.12–0.23) | 16.6 (12.0–20.9) | 4.2 (3.6–4.7) | - | [29] |
| <i>Allium ampeloprasum</i> L. | Bulbs | g/100 g | 78.3 (76.3–80.3) | 0.79 (0.59–0.99) | 1.67 (1.31–2.03) | 0.34 (0.13–0.61) | 16.6 (12.8–19.7) | 4.23 (3.72–4.74) | 85 (65–103) | [17] |
| <i>Amaranthus viridis</i> L. | - | % | 55.80 ± 0.23 | 13.31 ± 0.40 | 13.99 ± 0.12 | 1.40 ± 0.02 | 19.84 ± 0.07 | 6.54 ± 0.28 | 148.02 ± 0.28 | [31] |
| <i>Anchusa azurea</i> Mill. | Leaves | g/100 g | 91.2 (88.9–92.7) | 1.9 (1.8–2.1) | 1.9 (1.1–2.8) | 0.15 (0.07–0.23) | 1.3 (0.9–1.8) | 3.9 (3.5–4.4) | - | [29] |
| <i>Apium nodiflorum</i> (L.) Lag | Stems | g/100 g | 92.0 (90.0–94.0) | 1.7 (1.0–3.3) | 1.6 (1.1–2.1) | 0.10 (0.07–0.14) | 1.2 (0.7–2.1) | 2.7 (1.9–3.4) | - | [29] |
| <i>Asparagus acutifolius</i> L. | Stems | g/100 g | 84.6 ± 3.8 | 12.3 ± 0.0 | 22.4 ± 0.1 | 3.99 ± 0.33 | 61.3 ± 0.3 | - | 371 ± 1 † | [32] |
| <i>Asparagus acutifolius</i> L. | Shoots | g/100 g | 85.4 (81.2–88.5) | 2.23 (0.93–3.70) | 2.40 (1.69–3.25) | 0.61 (0.32–0.99) | 3.56 (1.03–4.67) | 4.83 (4.71–6.63) | 40 (23–56) | [17] |
| <i>Beta maritima</i> L. | Leaves | g/100 g | 87.3 (75.4–91.4) | 2.68 (2.00–5.60) | 3.10 (1.80–3.91) | 0.34 (0.18–0.70) | 1.71 (0.75–4.30) | 4.38 (3.29–9.50) | 31 (16–59) | [17] |
| <i>Beta vulgaris</i> subsp. <i>maritima</i> | Leaves | g/100 g | 84.5 (75.4–89.1) | 3.4 (2.0–5.6) | 2.6 (1.8–3.6) | 0.24 (0.16–0.40) | 3.6 (2.9–4.3) | 5.9 (3.9–9.5) | - | [29] |
| <i>Berberis aristata</i> DC. | Leaves | g/100 g dw ^ρ | 87.44 ± 2.22 | 15.46 ± 0.35 | 19.11 ± 0.78 | 2.14 ± 0.32 | 45.19 ± 0.56 | 18.10 ± 2.03 | - | [28] |
| <i>Blumea lacera</i> (Burm. f.) DC. | Leaves | g/100 g dw | 77.78 ± 2.68 | 24.05 ± 0.69 | 22.52 ± 0.97 | 0.93 ± 0.09 | 31.82 ± 1.26 | 20.68 ± 2.55 | - | [28] |
| <i>Borago officinalis</i> L. | Leaves | g/100 g | 86.9 (86.5–87.3) | 2.4 (2.2–2.5) | 1.2 (1.0–1.4) | 0.16 (0.13–0.19) | 9.5 (9.2–9.7) | - | 44 † | [30] |
| <i>Borago officinalis</i> L. | Leaves | g/100 g | 87.2 (86.4–88.8) | 2.35 1.93–2.91 | 2.35 (1.93–2.91) | 0.16 (0.13–0.19) | 9.45 (7.23–10.7) | - | 44 (34–50) | [17] |
| <i>Bryonia dioica</i> Jacq. | Stems | g/100 g dw | 82.9 ± 2.3 | 8.79 ± 0.01 | 16.6 ± 0.4 | 15.1 ± 1.9 | 59.5 ± 1.2 | - | 440 † | [32] |
| <i>Bryonia dioica</i> Jacq. | Shoots | g/100 g | 85.9 (70.9–90.8) | 1.48 (1.00–3.30) | 3.97 (1.00–11.9) | 1.12 (0.10–2.90) | 4.21 (0.80–10.37) | 4.60 (3.40–10.7) | 55 (14–141) | [17] |
| <i>Cichorium intybus</i> L. | Leaves | g/100 g fw ^γ | 86.4 (84.8–87.9) | 1.8 (1.7–2.1) | 2.9 (1.5–4.3) | 0.13 (tr–0.25) | 3.5 (1.8–4.7) | 6.1 (5.1–6.7) | 157 (137–180) | [25] |
| <i>Cichorium intybus</i> L. | Leaves | g/100 g | 87.9 (75.0–94.5) | 1.65 (1.25–2.10) | 1.83 (0.20–4.30) | 0.46 (Traces–0.92) | 3.50 (1.80–4.7) | 3.6 (1.20–6.70) | 33 (10–58) | [17] |
| <i>Chondrilla juncea</i> L. | Leaves | g/100 g | 87.8 ± 0.88 | 1.8 ± 0.11 | 1.9 ± 0.07 | 0.5 ± 0.04 | 2.0 ± 0.08 | 5.8 ± 0.32 | 19.6 | [33] |
| <i>Chondrilla juncea</i> L. | Leaves | g/100 g | 83.4 (65.9–89.7) | 2.41 (1.39–4.35) | 2.50 (1.83–6.13) | 0.80 (0.09–1.50) | 3.58 (1.49–9.69) | 7.70 (4.10–13.4) | 44 (22–104) | [17] |
| <i>Cynara cardunculus</i> L. | Flowers | g/100 g fw | 84.94 | 1.13 | 3.27 | 0.15 | 10.51 | 5.4 | 47 | [34] |
| <i>Enhydra fluctuans</i> Lour. | - | % | 67.69 ± 0.78 | 15.15 ± 0.44 | 8.00 ± 0.06 | 1.10 ± 0.01 | 9.64 ± 0.06 | 15.37 ± 0.21 | 80.53 ± 0.16 | [31] |
| <i>Erythrina variegata</i> L. | Leaves | g/100 g dw | 87.44 ± 2.22 | 20.15 ± 0.53 | 21.12 ± 1.58 | 1.55 ± 0.15 | 39.63 ± 1.11 | 17.55 ± 1.98 | - | [28] |

Table 1. Cont.

| Plant Species | Part of Plant | Unit | Moisture | Ash | Proteins | Fat | Carbohydrates | Fibre | Energy [§] | Reference |
|--|----------------|------------|--------------|---------------|---------------|-------------|---------------|---------------|------------------------|-----------|
| <i>Foeniculum vulgare</i> Mill. | Leaves | g/100 g | 86.7 | - | 3.8 | - | 4.9 | 3.5 | 48 | [35] |
| <i>Foeniculum vulgare</i> Mill. | Leaves | g/100 g | 82.4 | 2.34 | 2.76 | 0.42 | 9.67 | 3.87 | 63 (14–130) | [17] |
| <i>Glechoma hederacea</i> L. | Leaves | g/100 g | 73.1 ± 8.05 | 3.47 ± 0.1 | 1.34 ± 0.00 | 1.18 ± 0.23 | 21.0 ± 0.17 | - | 99.96 ± 0.80 | [27] |
| <i>Humulus lupulus</i> L. | Leaves | g/100 g | 85.5 | 1.4 (0.9–2.0) | 4.3 (3.1–5.1) | 0.20 | 1.6 (1.4–1.8) | 5.2 (4.3–6.4) | - | [29] |
| <i>Humulus lupulus</i> L. | Shoots | g/100 g | 85.8 | 1.35 | 4.25 | 0.37 | 1.85 | 4.85 | 39 (29–55) | [17] |
| <i>Hygrophilla schulli</i> (Hamilt.) M.R. Almeida & S.M. Almeida | Leaves | g/100 g dw | 91.23 ± 1.01 | 23.36 ± 0.66 | 17.19 ± 1.49 | 1.92 ± 0.18 | 43.46 ± 0.42 | 14.07 ± 1.21 | - | [28] |
| <i>Ipomoea aquatica</i> Forssk. | - | % | 69.11 ± 0.72 | 16.37 ± 0.67 | 13.82 ± 0.08 | 2.19 ± 0.08 | 10.51 ± 0.08 | 7.44 ± 0.27 | 117.27 ± 0.24 | [31] |
| <i>Malva sylvestris</i> L. | Flowers | g/100 g | 72.49 ± 1.89 | 10.54 ± 0.30 | 8.50 ± 0.51 | 2.84 ± 0.37 | 78.12 ± 0.44 | - | 372.02 ± 2.13 ‡ | [36] |
| <i>Malva sylvestris</i> L. | Leaves | g/100 g | 81.0 | 3.21 | 3.00 | 0.56 | 2.23 | 4.76 | 35 (23–50) | [17] |
| <i>Mentha pulegium</i> L. | Inflorescences | g/100 g | 59.47 ± 9.22 | 5.92 ± 0.09 | 7.12 ± 0.49 | 2.22 ± 0.22 | 84.74 ± 0.59 | - | 387.44 ± 0.53 ‡ | [37] |
| <i>Montia fontana</i> subsp. <i>amportitana</i> Sennen | Leaves | g/100 g | 91.47 ± 1.18 | 1.13 ± 0.13 | 1.76 ± 0.14 | 1.94 ± 0.13 | 1.81 ± 0.55 | 4.44 ± 0.34 | 31.48 ± 1.18 | [38] |
| <i>Nasturtium officinale</i> R. Br. | Leaves | g/Kg | 931 ± 10 | 9.4 ± 0.9 | 22.4 ± 0.7 | 1.43 ± 0.08 | 35.6 ± 0.9 | - | 1023 ± 15 ^a | [39] |
| <i>Oldenlandia corymbosa</i> Aiton. | - | % | 60.28 ± 0.40 | 8.34 ± 0.39 | 10.52 ± 0.10 | 2.16 ± 0.06 | 9.08 ± 0.37 | 7.26 ± 0.30 | 97.94 ± 0.04 | [31] |
| <i>Origanum vulgare</i> L. | Leaves | g/100 g | 51.82 ± 5.11 | 2.87 ± 0.07 | 2.28 ± 0.03 | 2.81 ± 0.33 | 40.22 ± 0.28 | - | 195.31 ± 0.96 | [27] |
| <i>Papaver rhoeas</i> L. | Leaves | g/100 g | 91.0 | - | 2.9 | - | 3.1 | 2.5 | 36 | [35] |
| <i>Papaver rhoeas</i> L. | Leaves | g/100 g | 88.3 | 2.50 | 3.50 | 0.64 | 3.35 | 4.40 | 42 (24–78) | [17] |
| <i>Portulaca oleracea</i> L. | Leaves | % | 81.5 | 28.9 | 27.8 | 0.141 | - | 10.0 | - | [40] |
| <i>Portulaca oleracea</i> L. | Leaves | g/100 g | 92.6 | 1.88 | 3.00 | 0.35 | 1.98 | 1.20 | 25 (19–32) | [17] |
| <i>Pterospartum tridentatum</i> (L.) Willk. | Flowers | g/100 g | 60.8 ± 0.16 | 2.36 ± 0.00 | 15.92 ± 0.60 | 2.69 ± 0.51 | 79.03 ± 0.74 | - | 404.01 ± 4.05 | [41] |
| <i>Rumex acetosella</i> L. | Leaves | g/100 g | 89.9 ± 1.01 | 10.93 ± 1.06 | 7.85 ± 1.86 | 2.35 ± 0.28 | 78.87 ± 1.50 | - | 368.03 ± 3.98 ‡ | [30] |
| <i>Rumex induratus</i> Boiss. & Reut | Leaves | g/100 g | 90.29 ± 0.53 | 11.07 ± 0.30 | 13.54 ± 0.28 | 3.97 ± 0.14 | 71.42 ± 0.28 | - | 375.55 ± 0.36 ‡ | [30] |

Table 1. Cont.

| Plant Species | Part of Plant | Unit | Moisture | Ash | Proteins | Fat | Carbohydrates | Fibre | Energy [§] | Reference |
|--|----------------|-------------------------|----------------------|---------------------|---------------------|---------------------|---------------------|---------------------|-------------------------------|-----------|
| <i>Rumex papillaris</i> Boiss. & Reut | Leaves | g/100 g | 89.1 (87.8–90.7) | 1.0 (0.4–1.3) | 2.4 (1.6–3.5) | 0.22 (0.26–0.28) | 2.0 (1.6–2.7) | 4.4 (4.0–5.0) | - | [29] |
| <i>Rumex pulcher</i> L. | Leaves | g/100 g | 86.6 (87.4–89.2) | 1.9 (1.1–3.1) | 3.2 (1.9–5.5) | 0.20 (0.10–0.32) | 3.3 (1.5–4.5) | 4.7 (4–5.2) | - | [29] |
| <i>Raphanus raphanistrum</i> L. | Leaves | g/100 g fw | 89.9 ± 0.6 | 1.58 ± 0.08 | 4.04 ± 0.01 | 0.23 ± 0.03 | 4.22 ± 0.08 | - | 35.1 ± 0.1 | [42] |
| <i>Scolymus hispanicus</i> L. | Leaves | g/100 g fw | 84.1 (81.8–92.7) | 3.19 (1.7–5.2) | 1.8 (0.3–5.3) | 0.09 (0.08–0.11) | 3.4 (1.1–9.2) | 7.0 (3.1–12.) | 167 (53–280) [¥] | [25] |
| <i>Sesbania sesban</i> (L.) Merr. | Leaves | g/100 g dw | 90.13 ± 1.55 | 18.68 ± 0.22 | 15.65 ± 1.10 | 0.97 ± 0.05 | 49.51 ± 0.72 | 15.19 ± 1.79 | - | [28] |
| <i>Silene vulgaris</i> (Moench) Garcke | Leaves | g/100 g | 87.1 (80.4–88.5) | 0.3 | 3.3 (3.0–3.6) | 0.70 | 3.4 (2.9–3.9) | 2.8 (2.6–3.1) | - | [43] |
| <i>Silene vulgaris</i> (Moench) Garcke | Leaves | g/100 g | 85.9 (86.6–88.5) | 1.53 (0.20–4.33) | 2.47 (1.31–3.60) | 0.67 (0.31–1.31) | 2.32 (1.03–3.90) | 4.36 (2.60–6.63) | 34 (17–56) | [17] |
| <i>Silybum marianum</i> (L.) Gaertn. | Leaves | g/100 g fw | 93.4 (92.9–93.8) | 1.5 (1.0–1.9) | 0.6 (0.5–0.8) | 0.01 (tr–0.03) | 1.1 (0.5–1.7) | 2.6 (2.3–2.9) | 51.8 (42.2–61.4) [¥] | [25] |
| <i>Sonchus asper</i> L. | Leaves | g/Kg | 864.3 ± 11.2 | 30.4 ± 3.0 | 32.5 ± 3.2 | 6.8 ± 0.7 | 19.8 ± 1.1 | 35.6 ± 2.2 | 1110 ± 120 | [44] |
| <i>Sonchus oleraceus</i> L. | Leaves | g/Kg | 872.4 ± 14.0 | 29.9 ± 1.8 | 31.7 ± 1.5 | 7.5 ± 0.9 | 18.2 ± 1.4 | 32.5 ± 2.4 | 1110 ± 120 | [44] |
| <i>Sonchus oleraceus</i> L. | Leaves | g/100 g | 87.6 (83.0–91.9) | 2.17 (1.58–3.00) | 2.22 (1.11–3.48) | 0.60 (0.20–1.28) | 2.29 (0.94–4.20) | 3.37 (2.60–5.57) | 33 (16–56) | [17] |
| <i>Sonchus oleraceus</i> L. | Leaves | g/100 g fw | 88.25 (83.2–91.0) | 2.04 (1.6–2.7) | 2.22 (1.3–3.5) | 0.29 (0.20–0.41) | 2.51 (0.9–4.2) | 4.3 (3.5–5.6) | 127 (91–163) [¥] | [25] |
| <i>Sonchus oleraceus</i> L. | Leaves | g/100 g wb [°] | 89.3 ± 3.04 | 1.5 ± 0.01 | 3.0 ± 0.13 | 0.4 ± 0.01 | - | 5.5 ± 0.35 | - | [45] |
| <i>Sonchus tenerrimus</i> L. | Leaves | g/Kg | 877.3 ± 20.8 | 30.2 ± 1.9 | 31.8 ± 2.0 | 5.2 ± 0.4 | 13.2 ± 1.8 | 31.2 ± 1.6 | 935 ± 105 | [44] |
| <i>Tamus communis</i> L. | Leaves | g/100 g | 86.2 (84.6–89.0) | 1.4 (0.9–2.4) | 3.2 (2.5–3.8) | 0.17 (0.10–0.22) | 2.2 (1.9–2.7) | 4.7 (3.5–6.0) | - | [29] |
| <i>Tamus communis</i> L. | Shoots | g/100 g | 85.2 (82.0–89.0) | 1.25 (0.90–2.40) | 3.13 (2.52–3.85) | 0.49 (0.10–1.28) | 5.20 (1.80–11.7) | 4.35 (3.50–6.00) | 46 (25–85) | [17] |
| <i>Taraxacum obovatum</i> (Willd.) DC. | Leaves | g/100 g fw | 83.3 (79.2–86.7) | 2.13 (1.8–2.5) | 1.57 (1.02–2.09) | 0.22 (0.19–0.27) | 3.34 (1.63–5.39) | 7.01 (5.4–8.7) | 152 (114–205) [¥] | [25] |
| <i>Thymus mastichina</i> L. | Leaves | g/100 g | 54.67 ± 7.03 | 2.67 ± 0.08 | 2.2 ± 0.5 | 3.80 ± 0.10 | 36.64 ± 0.08 | - | 189.65 ± 0.44 | [27] |
| <i>Thymus pulegioides</i> L. | Inflorescences | g/100 g | 47.6 ± 12.60 | 4.94 ± 0.62 | 5.53 ± 1.40 | 0.18 ± 0.02 | 89.35 ± 1.54 | - | 381.14 ± 1.76 [‡] | [37] |
| <i>Viola x Wittrockiana</i> | Flowers | g/100 g dw | 87.76 (wet matter) | 7.92 | 10.14 | 1.67 | 80.27 | - | 376.67 [‡] | [4] |
| <i>Umbilicus rupestris</i> (Salisb.) Dandy | Leaves | g/100 g fw | 93 ± 1 | 0.91 ± 0.01 | 1.83 ± 0.06 | 0.255 ± 0.002 | 3.90 ± 0.03 | - | 25.2 ± 0.1 | [46] |

[§] kcal/100 g; [‡] kcal/100 g dw; [°] kJ/kg; [¥] kJ/100 g; Mean value (minimum-maximum); [°] dw = dry weight; [¥] fw = fresh weight; [°] wb = wet bases.

With respect to protein, three WEPs stood out for their high protein content, each belonging to a different family (Portulacaceae, Asteraceae, and Asparagaceae): *Portulaca oleracea* with 27.8 g per 100 g [40], followed by *Blumea lacera* and *Asparagus acutifolius* with 22.52 and 22.40 g per 100 g, respectively [28,32]. On the contrary, the plant with the lowest protein content was *Silybum marianum* (Asteraceae family), with a protein content as low as 0.6 g per 100 g [25,26].

Fat content was only reported in 47 of the available 115 WEPs. *Bryonia dioica* (Cucurbitaceae family) showed the highest content at 15.1 per 100 g dry weight (Martins et al., 2011). The family with the highest number of plants (Asteraceae) had, in general, low contents, reaching its maximum with *Silybum marianum* and *Enhydra fluctuans* (1.1 g per 100 g), and its minimum with *Scolymus hispanicus* (0.09 g per 100 g) [25,26,31].

The scientific literature only provides the carbohydrates content for 46 WEPs out of the 115 reviewed plants. In this regard, *Thymus pulegioides* and *Mentha pulegium* had the highest content, reaching 89.35 and 84.74 per 100 g, respectively [37]; both plants belong to the Lamiaceae family. Within this family, plants showed a wide variability in carbohydrate content, with *Thymus pulegioides* having the highest value (89.35 g per 100 g), while *Glechoma hederacea* had the lowest with 21 g per 100 g [27].

The content of dietary fiber was only found for 36 WEPs. The WEPs that stood out for their high fiber content were *Blumea lacera* (20.68 g per 100 g dw) and *Berberis aristata* (18.10 g per 100 g dw), belonging to the Asteraceae and Berberaceae families, respectively [28]. In general, the fiber content of the Lamiaceae and Brassicaceae plants was not analyzed and thus was not reported [27,37,39,42].

Energy is calculated from the determination of food macro-nutrients including protein, fiber, carbohydrates, fat and alcohol [47]. Nowadays, part of society is willing to have a balanced diet, but unfortunately, most consumers, due to their lifestyle, replace traditional diets with diets high in sugars and refined fats, which leads to large caloric intakes; these diets result in increased incidence of coronary heart disease, strokes, type II diabetes, and obesity [48,49]. However, in the USA the actual daily intake for men is 2800 kcal per day and for women 2000–2200 kcal per day, which are comparable to the 2030 kcal that a person 2 m tall and 88 kilos should ingest, this value being the highest of those established for men and women by the FAO [50]. This difference in daily kilocalories shows a gap between the ideal intake and that which is averaged in countries such as the USA. Energy data was only available for 28 WEPs out of the 115 plants reviewed. The plant with the highest energy value was *Bryonia dioica* with 440 kcal per 100 g dw [32]. Regarding families, the Polygonaceae showed high contents, for instance, *Rumex acetosella* and *Rumex induratus* with 368 and 376 kcal per 100 g dw, respectively [30].

5. Sugars and Organic Acids

5.1. Sugars

The role of the sugars, which are generated through the photosynthesis process, in plants is fundamental; they are the main source of carbon and energy for the plants, and participate in the plant metabolism control, for example participating in multiple biological processes, from embryogenesis to plant senescence [51,52]. Sugars can be classified into monosaccharides, disaccharides, and polysaccharides. Apart from the biological importance of sugars in plants, it is necessary to highlight their importance in the health of humans. Sugars are the main source of energy for multiple metabolic processes, as well as making a necessary contribution for cells to stay alive [53].

Sugar contents of 22 WEPs (from 16 families) are summarized in Table 2. In general, the predominant sugars were fructose, glucose and sucrose. It should be noted that, of the three main sugars, glucose was that with the highest total content in all 22 plants, followed by fructose and subsequently sucrose.

Table 2. Sugars reported in wild edible plants, WEPs.

| Plant Species | Part of Plant | Unit | Sugars | | | Reference |
|---|----------------|-------------|-----------------|----------------|----------------|-----------|
| | | | Fructose | Glucose | Sucrose | |
| <i>Asparagus acutifolius</i> L. | Stems | g/100 g dw | 2.49 ± 0.13 | 1.98 ± 0.04 | 4.27 ± 0.12 | [32] |
| <i>Borago officinalis</i> L. | Leaves | g/100 g dw | 0.14 ± 0.03 | 0.58 ± 0.06 | 1.52 ± 0.13 | [30] |
| <i>Bryonia dioica</i> Jacq. | Stems | g/100 g dw | 3.45 ± 0.08 | 2.97 ± 0.09 | 0.572 ± 0.014 | [32] |
| <i>Chenopodium ambrosioides</i> L. | Leaves | g/100 g dw | 0.24 ± 0.01 | 0.46 ± 0.01 | 1.43 ± 0.12 | [54] |
| <i>Foeniculum vulgare</i> Mill. | Leaves | g/100 g fw | 0.49 ± 0.05 | 0.76 ± 0.12 | 0.04 ± 0.00 | [55] |
| <i>Glechoma hederacea</i> L. | Leaves | g/100 g fw | 0.15 ± 0.01 | 0.08 ± 0.02 | 0.40 ± 0.06 | [27] |
| <i>Helichrysum stoechas</i> (L.) Moench | Stems | g/100 g dw | 1.02 ± 0.04 | 0.59 ± 0.02 | 1.84 ± 0.09 | [56] |
| <i>Malva sylvestris</i> L. | Flowers | g/100 g dw | 8.72 ± 0.14 | 7.36 ± 0.13 | 2.74 ± 0.05 | [36] |
| <i>Mentha pulegium</i> L. | Inflorescences | g/100 g dw | 2.39 ± 0.11 | 3.37 ± 0.22 | 4.62 ± 0.28 | [37] |
| <i>Montia fontana</i> subsp. <i>amporitana</i> Sennen | Leaves | g/100 g dw | 0.76 ± 0.17 | 1.00 ± 0.02 | 0.44 ± 0.05 | [30] |
| <i>Nasturtium officinale</i> R. Br. | Leaves | mg/kg | 1104 ± 31 | 696 ± 20 | 233 ± 51 | [39] |
| <i>Origanum vulgare</i> L. | Leaves | g/100 g fw | 0.19 ± 0.01 | 0.58 ± 0.01 | 0.30 ± 0.00 | [27] |
| <i>Portulaca oleracea</i> L. | Leaves | mg/100 g fw | 290 (202–352) ‡ | 81.25 (59–118) | 161.5 (75–271) | [57] |
| <i>Pterospartum tridentatum</i> (L.) Willk. | Flowers | g/100 g dw | 3.49 ± 0.11 | 1.19 ± 0.05 | 0.58 ± 0.03 | [41] |
| <i>Rubus ulmifolius</i> Schott | Flowers | g/100 g dw | 1.66 ± 0.21 | 2.23 ± 0.19 | 1.34 ± 0.15 | [56] |
| <i>Rumex acetosella</i> L. | Leaves | g/100 g dw | 0.60 ± 0.00 | 0.73 ± 0.01 | 0.21 ± 0.07 | [30] |
| <i>Rumex induratus</i> Boiss. & Reut | Leaves | g/100 g dw | 1.71 ± 0.09 | 1.26 ± 0.20 | 1.25 ± 0.31 | [30] |
| <i>Raphanus raphanistrum</i> L. | Leaves | g/100 g fw | 0.153 ± 0.004 | 0.348 ± 0.003 | - | [42] |
| <i>Tamus communis</i> L. | Leaves | g/100 g dw | 3.83 ± 0.13 | 1.80 ± 0.14 | 0.695 ± 0.05 | [32] |
| <i>Thymus mastichina</i> L. | Leaves | g/100 g fw | 0.45 ± 0.01 | 0.97 ± 0.11 | 0.02 ± 0.00 | [27] |
| <i>Thymus pulegioides</i> L. | Inflorescences | g/100 g dw | 0.22 ± 0.00 | 0.33 ± 0.03 | 1.06 ± 0.02 | [37] |
| <i>Umbilicus rupestris</i> (Salisb.) Dandy | Leaves | g/100 g fw | - | - | 0.082 ± 0.002 | [46] |

‡ Mean value (minimum-maximum).

The most commonly studied plant family regarding sugars was Lamiaceae with five plants. Regarding fructose, the WEPs that had the highest content were *Malva sylvestris* (Malvaceae family) and *Tamus communis* (Dioscoraceae family), with 8.72 and 3.83 g/100 g dw, respectively [32,36]. *Malva sylvestris* also showed the highest glucose content at 7.35 g/100 g dw [36]. The sucrose concentration in these 22 WEPs was mainly dominated by two plants: *Mentha pulegium*, which had 4.62 g/100 g dw, and *Asparagus acutifolius* with a sucrose concentration of 4.27 g/100 g dw [32,37].

5.2. Organic Acids

At a general level, organic acids are weak acids, which can be classified mainly by four criteria: (i) nature of the carbon chain (aliphatic, aromatic, etc.); (ii) saturation or unsaturation properties; (iii) substituted or unsubstituted characteristics; and (iv) number of functional groups. Acids play an essential role in the physiology of plants, participating in processes such as pH regulation, balancing the redox potential cells, the Krebs cycle, or even in organoleptic properties such as color, taste and aroma of both fruits and vegetables [58–60].

Three organic acids (oxalic, malic and shikimic) were mainly found in WEPs, and especially in four families: Asteraceae, Brassicaceae, Crassulaceae, and Portulacaceae (Table S2).

One of the most representative organic acids in the Asteraceae family is oxalic acid [25,26], which show a high content in six plants of this family [61]. Being a small group of plants in which organic acids were determined, a PCA (Principal Component Analysis) was carried out to understand these results in a visual way (Figure 2). The values can be found in Table S2. PCA showed a high percentage of correlation (92.08%) among organic acids and WEPs, being able to differentiate three groups. In the first cluster, malic acid was

associated with *Raphanus raphanistrum* (Brassicaceae family); the second cluster was based on the association of oxalic acid with *Sonchus oleraceus* (Asteraceae family); while in the third group both oxalic acid and shikimic acid were associated with *Hymenonema graecum* (Asteraceae family).




Figure 2. Principal Component Analysis (PCA) of organic acids present in wild edible plants, WEPs.

In this way, the highest contents of malic acid were reported in *Raphanus raphanistrum* and *Sonchus oleraceus* (580 and 415 mg/100 g, respectively) [42,61]. The WEPs in which the highest amounts of oxalic acid were found, in decreasing order were: *Hymenonema graecum* > *Sonchus oleraceus* > *Raphanus raphanistrum*, with concentrations between 972 and 706 mg/100 g [42,61]. Regarding shikimic acid, two plants stood out for their contents: *Hymenonema graecum* and *Sonchus oleraceus*, with 244 and 166 mg/100 g, respectively; the other WEPs where this acid was found had values of below 100 mg/100 g [61].

6. Mineral Elements

The Mediterranean diet is considered by many experts as one of the best in the world at a nutritional level, and one of its main strengths is the contribution of minerals and vitamins [62]. In body composition, minerals represent fourth place in abundance, reaching values of up to 6.1% of the body weight for a person of 65 kg [50]. This importance of minerals for the human body determines health problems due to mineral deficiency; in this way, key elements which are often linked to deficiency are iron, zinc, and iodine. Iron deficiency can affect as many as 18% of the world's children under 5 years of age, along with pregnant women [63]. In addition, zinc and iodine affect 17% and 28% of the world's population, respectively [63]. According to Zeece [64], the main source of minerals is the soil, because from there it passes to plants and through the food chain to humans. The daily intake of minerals establishes the recommended intake for each of these elements; in Table S3, it is possible to find these values according to the Food & Drug Administration (FDA) [65].

Principal Component Analysis (PCA) (Figure 3) was carried out after adjusting all values in a common unit, mg/100 g fw. PCA showed a total percentage of correlation (70.13%) among minerals contents and WEPs. Mineral composition was reported only for 17 of the 115 plants (from 10 families) covered in this review. In this way, most of the plants which had their mineral profile analyzed ($n = 7$) belonged to the Asteraceae family. The macro-elements identified were calcium (Ca), magnesium (Mg), potassium (K), and sodium (Na); while the micro-elements were copper (Cu), iron (Fe), manganese (Mn), and zinc (Zn). In Table S4 [25,34,35,38,57,66–68], values for the contents of these minerals can be found.




Figure 3. Principal Component Analysis of minerals present in wild edible plants, WEPs.

Regarding Figure 3, three different groups can be easily observed. On the one hand, Ca was isolated from the entire group of minerals as not being associated with any plant. The first group contained the highest number of minerals (Cu, Fe, Mn, Mn, and Zn). The second and third groups were linked to K and Na, respectively.

Calcium was not associated with any specific WEPs, although *Foeniculum vulgare* was that with the highest content of this mineral, followed by *Chondrilla juncea* (341 and 301 mg/100 g, respectively) [25,35].

The first group was mainly associated with two plants: *Chondrilla juncea* and *Malva sylvestris*. Regarding Cu and Mn, the WEPs that had the highest simultaneous contents, were *Chondrilla juncea* (0.43 and 0.97 mg/100 g, respectively) followed by *Malva sylvestris* (0.33 and 0.76 mg/100 g, respectively); although the WEP that had the highest Mn content was *Montia fontana* with 1.08 mg/100 g [38]. Regarding Fe, Mg, and Zn, *Malva sylvestris* had the highest contents (5.82, 715 and 1.98 mg/100 g, respectively) compared to *Chondrilla juncea* (3.97, 40.80 and 1.63 mg/100 g, respectively) [25]. In the second group, K was exclusively associated with *Scolymus hispanicus*, the content of which was much higher than in the rest of the WEPs (1040 mg/100 g) [25,26]. In the third group, Na was exclusively associated with *Beta maritima*, containing the highest concentration of this mineral, 171 mg/100 g [66].

The mineral values clearly demonstrated that many WEPs have a high nutritional value, for example, *Chondrilla juncea*, which can provide almost 50% of the recommended daily intake of Cu after consumption of 200 g of the plant. *Montia fontana*, with its Mn content, could also cover approximately 50% of the recommended daily amount after consumption of 200 g (Table S3) [65].

7. Fatty Acids

Of the 115 WEPs evaluated in this review, only 43 (from 21 families) had FA profiles published; the most commonly studied families regarding FAs were: Asteraceae (8) > Lamiaceae (5) > Polygonaceae (4) > Brassicaceae (3). As can be seen in Table 3, the most representative FAs in WEPs were palmitic, oleic, linoleic, and linolenic acids.

Table 3. Fatty acids reported in wild edible plants, WEPS.

| Plant Species | Part of Plant | Unit | C16:0 (Palmitic Acid) | C18:1 (Oleic Acid) | C18:2 (Linoleic Acid) | C18:3 (Linolenic Acid) | Reference |
|---|----------------|------------------|-----------------------------|-----------------------|-----------------------------|------------------------------|-----------|
| <i>Allium ampeloprasum</i> L. | Bulbs | % | 26.42 ± 0.30 | 7.39 ± 0.42 | 53.45 ± 0.27 | nd | [67] |
| <i>Anchusa azurea</i> Mill. | Leaves | % | 10.45 ± 0.62 | 2.20 ± 0.00 | 12.16 ± 0.11 | 64.74 ± 0.23 | [69] |
| <i>Apium nodiflorum</i> (L.) Lag | Stems | % | 16.29 ± 0.96 | 3.33 ± 0.02 | 24.60 ± 0.77 | 43.46 ± 0.08 | [70] |
| <i>Asparagus acutifolius</i> L. | Stems | % | 17.5 ± 0.2 | 4.94 ± 0.35 | 44.5 ± 1.3 | 23.7 ± 0.9 | [32] |
| <i>Bellis perennis</i> L. | Aerial parts | % | 1.5 | - | 5.1 | 13.6 | [71] |
| <i>Beta maritima</i> L. | Leaves | % | 11.03 ± 0.15 | 3.51 ± 0.01 | 21.28 ± 0.04 | 57.80 ± 0.03 | [70] |
| <i>Borago officinalis</i> L. | Leaves | % | 12.03 ± 0.70 | 2.08 ± 0.20 | 9.50 ± 1.25 | 12.26 ± 1.90 | [30] |
| <i>Bryonia dioica</i> Jacq. | Stems | % | 13.5 ± 0.3 | 1.52 ± 0.09 | 6.39 ± 0.16 | 70.3 ± 0.1 | [32] |
| <i>Cichorium intybus</i> L. | Leaves | % | 10.64 ± 0.63 | 1.61 ± 0.04 | 21.14 ± 0.06 | 60.45 ± 0.41 | [70] |
| <i>Chenopodium ambrosioides</i> L. | Leaves | % | 14.16 ± 0.03 | 6.90 ± 0.12 | 19.23 ± 0.12 | 48.54 ± 0.13 | [54] |
| <i>Chondrilla juncea</i> L. | Leaves | % | 12.96 ± 0.47 | 1.91 ± 0.01 | 19.92 ± 0.17 | 56.27 ± 0.13 | [70] |
| <i>Daucus carota</i> L. | Roots | % | tr | - | 1.0 | - | [71] |
| <i>Diplotaxis eruroides</i> (L.) DC | Aerial parts | % | 18.23 | 3.42 | 39.31 ‡ | - | [11] |
| <i>Diplotaxis virgata</i> (Cav.) DC | Flowers | % | 14.35 | - | 36.01 ‡ | - | [11] |
| <i>Foeniculum vulgare</i> Mill. | Leaves | % | 20.15 ± 0.09 | 4.35 ± 0.37 | 23.25 ± 0.07 | 43.55 ± 0.40 | [55] |
| <i>Glechoma hederacea</i> L. | Leaves | % | 12.23 ± 0.23 | 35.12 ± 0.27 | 8.15 ± 0.08 | 27.87 ± 0.20 | [27] |
| <i>Helichrysum stoechas</i> (L.) Moench | Stems | % | 13.24 ± 0.16 | 6.15 ± 0.79 | 25.67 ± 0.08 | 22.79 ± 1.86 | [56] |
| <i>Humulus lupulus</i> L. | Leaves | % | 19.52 ± 0.61 | 1.88 ± 0.10 | 29.72 ± 0.85 | 38.16 ± 0.02 | [70] |
| <i>Malva sylvestris</i> L. | Flowers | % | 9.79 ± 1.07 | 3.31 ± 0.42 | 11.96 ± 0.42 | 67.79 ± 0.96 | [36] |
| <i>Mentha pulegium</i> L. | Inflorescences | % | 14.82 ± 0.09 | 5.77 ± 0.20 | 16.27 ± 0.33 | 37.00 ± 0.35 | [37] |
| <i>Montia fontana</i> subsp. <i>amporitana</i> Sennen | Leaves | % | 17.22 ± 1.06 | 2.37 ± 0.38 | 18.71 ± 0.45 | 55.57 ± 0.80 | [30] |
| <i>Origanum vulgare</i> L. | Leaves | % | 4.95 ± 0.10 | 5.08 ± 0.01 | 23.22 ± 0.14 | 62.34 ± 0.04 | [27] |
| <i>Papaver rhoeas</i> L. | Leaves | % | 9.66 ± 0.39 | 1.36 ± 0.00 | 16.53 ± 0.01 | 64.98 ± 0.07 | [70] |
| <i>Portulaca oleracea</i> L. | Leaves | % | 24.7 (23.4–26.9) ¥ | 12.4 (9.7–15.1) | 28.8 (25.1–32.9) | 23.6 (17.9–28.4) | [57] |
| <i>Pterospartum tridentatum</i> (L.) Willk. | Flowers | % | 14.84 ± 0.83 | 9.22 ± 1.09 | 19.59 ± 0.67 | 29.50 ± 1.98 | [41] |
| <i>Rumex acetosella</i> L. | Leaves | % | 11.23 ± 0.73 | 3.43 ± 0.32 | 20.18 ± 0.48 | 51.34 ± 1.41 | [30] |
| <i>Rumex induratus</i> Boiss. & Reut | Leaves | % | 9.36 ± 0.71 | 2.20 ± 0.05 | 13.76 ± 0.01 | 58.84 ± 1.03 | [30] |
| <i>Rumex papillaris</i> Boiss. and Reut. | Leaves | % | 11.20 ± 0.32 | 5.80 ± 0.14 | 22.79 ± 0.19 | 51.77 ± 0.14 | [70] |
| <i>Rumex pulcher</i> L. | Leaves | % | 9.30 ± 0.11 | 4.22 ± 0.01 | 17.03 ± 0.16 | 62.97 ± 0.03 | [70] |
| <i>Raphanus raphanistrum</i> L. | Leaves | mg/100 g fw ¶ | 25.2 ± 2.3 | 2.4 ± 0.3 | 24 ± 2 | 171 ± 16 | [42] |
| <i>Sambucus nigra</i> L. | Aerial parts | % | tr | - | 16.1 | 2.6 | [71] |
| <i>Scolymus hispanicus</i> L. | Leaves | % | 20.65 ± 0.85 | 6.41 ± 0.07 | 26.44 ± 0.26 | 30.55 ± 0.23 | [70] |
| <i>Silene vulgaris</i> (Moench) Garcke | Leaves | % | 13.5 (13.1–15.1) | 2.4 (2.1–2.7) | 22.4 (18.9–24.4) | 54.5 (51.2–56.9) | [43] |
| <i>Silybum marianum</i> (L.) Gaertn. | Leaves | % | 28.69 ± 1.60 | 3.86 ± 0.10 | 31.01 ± 0.63 | 21.60 ± 0.25 | [70] |
| <i>Sonchus oleraceus</i> L. | Leaves | % | 10.43 ± 0.70 | 0.92 ± 0.10 | 13.78 ± 0.61 | nd | [70] |
| <i>Tamus communis</i> L. | Leaves | % | 17.0 ± 0.7 | 7.51 ± 0.18 | 42.0 ± 0.3 | 27.5 ± 0.4 | [32] |
| <i>Taraxacum obovatum</i> (Willd.) DC. | Leaves | % | 11.83 ± 0.09 | 3.24 ± 0.01 | 17.64 ± 0.08 | 58.53 ± 0.23 | [70] |
| <i>Thymus mastichina</i> L. | Leaves | % | 10.22 ± 0.20 | 9.82 ± 0.18 | 11.83 ± 0.06 | 45.65 ± 0.55 | [27] |
| <i>Thymus pulegioides</i> L. | Inflorescences | % | 16.70 ± 0.22 | 11.40 ± 0.10 | 12.98 ± 0.52 | 36.69 ± 0.25 | [37] |
| <i>V. x Wittrockiana</i> | Flowers | g/100 g dw £ | 36.41 | 8.27 | 32.30 | nd | [4] |
| <i>Umbilicus rupestris</i> (Salisb.) Dandy | Leaves | % | 10.6 ± 0.8 | 0.641 ± 0.002 | 18.3 ± 0.6 | 62 ± 2 | [46] |

‡ Linoleic acid C18: 2 y Linoleic acid C18:2; nd = not detected; tr = trace; ¥ Mean value (minimum-maximum);

£ dw = dry weight; ¶ fw = fresh weight.

Linolenic acid was by far the FA with the highest percentage in most plants, showing in eight of these 42 plants a percentage above 60%; the list of plants in decreasing order of abundance was as follows: *Bryonia dioica* Jacq. (70.3%) > *Malva sylvestris* (67.8%) > *Papaver rhoeas* (65.0%) > *Anchusa azurea* (64.7%) > *Rumex pulcher* (63.0%) > *Origanum vulgare* (62.3%) > *Umbilicus rupestris* (62.0%) > *Cichorium intybus* (62.0%) [27,32,46,56,70]. It is important to highlight that this compound was present in a wide range of families.

Linoleic acid was the second most abundant FA, highlighting its importance in WEPs, especially in *Allium ampeloprasum* (53.5%) > *Asparagus acutifolius* (44.5%) > *Tamus communis* (42.0%) [32,67,72]; these plants belong to the Amaryllidaceae, Asparagaceae and Dioscoreaceae families, respectively.

Another of the most representative FAs, palmitic acid, was present in all the WEPs analyzed. The WEP that obtained the highest value in palmitic acid was *Viola x wittrockiana* (Violaceae family) with 36.41 g/100 g dw [4]. *Silybum marianum* (Asteraceae), *Allium ampeloprasum* (Amaryllidaceae) and *Portulaca oleracea* (Portulacaceae) had values of 28.7%, 26.4% and 24.7%, respectively [57,67,70,72].

Regarding oleic acid, the highest content was found in *Glechoma hederacea* with 35.1% [27], followed by *Portulaca oleracea* with 12% [57]; whilst, the WEP with the lowest content of oleic acid was *Umbilicus rupestris* (0.6%) [46].

In general, the main FAs identified in WEPs were linolenic and linoleic acids. On the other hand, it should be noted that palmitic acid was identified in all plants, although at lower concentrations.

8. Phenolic Content and Flavonoids

Phenolic compounds are the most abundant secondary metabolites in plants, helping in various functions of great importance, such as pigmentation, growth, and/or resistance to pathogens, also playing a fundamental role in maintaining redox homeostasis of cells [73–75]. Apart from the importance in the physiology of plants themselves, it has been clearly demonstrated that these compounds have antioxidant activity, anti-inflammatory effects, help in the reduction of oxidative stress, and can even help in the prevention of tumors [76–79].

In the references evaluated in this review, instudying up to 115 WEPs (from 47 families), the total phenolic contents (TPC) were provided for 100 plants; for 51 plants, the total flavonoids content (TFC) was also provided (Table 4) [5,30,32,36,39,42,46,54,56,57,67,69,72,80,81]. Regarding TPC and TFC, the most commonly studied families were Asteraceae (17) > Brassicaceae (7) > Fabaceae (6) = Lamiaceae (6). These analyses were performed by HPLC [31,54,81–84] and spectrophotometric method [4,5,28,30,32,36,37,39,41,56,57,67,69,72,80].

Table 4. Total phenolic content (TPC) and total flavonoids content (TFC) in wild edible plants.

| Plant Species | Part of Plants | Unit TPC | TPC | Unit TFC | TFC | Reference |
|--|----------------|-------------------|-----------------------------|------------------|------------------------|-----------|
| <i>Asystasia gangetica</i> (L.) T.Anderson | - | mg GAE/g de † | 91.797 ± 0.295 | mg RE/g de § | 20.132 ± 0.093 | [31] |
| <i>Achyranthes aspera</i> L. | - | mg GAE/g de | 74.831 ± 0.243 | mg RE/g de | 20.793 ± 0.122 | [31] |
| <i>Ageratum conyzoides</i> (L.) L. | Flowers | mg GAE/g | 4.63 ± 0.52 | - | - | [81] |
| <i>Allamanda cathartica</i> L. | Flowers | mg GAE/g | 4.16 ± 0.11 | - | - | [81] |
| <i>Allium ampeloprasum</i> L. | Bulbs | mg GAE/g | 5.70 ± 0.62 | mg CE/g ° | 0.86 ± 0.5 | [67] |
| <i>Amaranthus viridis</i> L. | - | mg GAE/g de | 50.700 ± 0.079 | mg RE/g de | 19.970 ± 0.252 | [31] |
| <i>Anagallis arvensis</i> (L.) | Aerial parts | mg GAE/g | 27.54 ± 0.92 | mg QE/g £ | 26.15 ± 0.85 | [84] |
| <i>Anchusa azurea</i> Mill. | Leaves | mg GAE/g extract. | 148.62 (146.62–150.62) ¥ | mg CE/g extract. | 84.81 (80.78–88.84) | [72] |
| <i>Apium nodiflorum</i> (L.) Lag. | Leaves | mg GAE/g extract. | 80.47 ± 4.41 | mg CE/g extract. | 45.48 ± 1.61 | [69] |
| <i>Asparagus acutifolius</i> L. | Stems | mg GAE/g extract. | 624 ± 28 | mg CE/g extract. | 57.8 ± 2.4 | [32] |

Table 4. Cont.

| Plant Species | Part of Plants | Unit TPC | TPC | Unit TFC | TFC | Reference |
|--|----------------|-------------------------|--------------------|------------------|--------------------|-----------|
| <i>Asparagus acutifolius</i> L. | Leaves | mg CAE/kg ww \uparrow | 43.1 | mg QE/kg ww | 2262.9 | [83] |
| <i>Bahuinia purpurea</i> L. | Flowers | mg GAE/g | 6.14 \pm 0.30 | - | - | [81] |
| <i>Beta maritima</i> L. | Leaves | mg GAE/g extract. | 61.91 \pm 7.51 | mg CE/g extract. | 21.55 \pm 0.87 | [72] |
| <i>Berberis aristata</i> DC. | Leaves | mg GAE/g dw | 135.56 \pm 3.26 | mg QE/g dw | 82.05 \pm 0.78 | [28] |
| <i>Bidens pilosa</i> L. | Flowers | mg GAE/g | 8.12 \pm 0.41 | - | - | [81] |
| <i>Blumea lacera</i> (Burm. f.) DC. | Leaves | mg GAE/g dw | 95.23 \pm 1.35 | mg QE/g dw | 59.87 \pm 0.93 | [28] |
| <i>Bombax malabaricum</i> DC. | Flowers | mg GAE/g | 3.88 \pm 0.17 | - | - | [81] |
| <i>Borago officinalis</i> L. | Leaves | mg CAE/kg ww | 64.0 | mg QE/kg ww | 189.4 | [83] |
| <i>Bougainvillea spectabilis</i> Willd. | Flowers | mg GAE/g | 6.87 \pm 0.23 | - | - | [81] |
| <i>Brassica campestris</i> L. | Flowers | mg GAE/g | 3.32 \pm 0.09 | - | - | [81] |
| <i>Brunfelsia acuminata</i> (Pohl) Benth. | Flowers | mg GAE/g | 4.08 \pm 0.25 | - | - | [81] |
| <i>Bryonia dioica</i> Jacq. | Stems | mg GAE/g extract. | 258 \pm 22 | mg CE/g extract. | 18.1 \pm 1.2 | [32] |
| <i>Calliandra haematocephala</i> Hassk. | Flowers | mg GAE/g | 14.43 \pm 0.71 | - | - | [81] |
| <i>Camellia japonica</i> L. | Flowers | mg GAE/g | 5.14 \pm 0.29 | - | - | [81] |
| <i>Chaenomeles sinensis</i> (Thouin) Koehne | Flowers | mg GAE/g | 13.93 \pm 0.34 | - | - | [81] |
| <i>Chrysanthemum coronarium</i> L. | Flowers | mg GAE/g | 3.76 \pm 0.29 | - | - | [81] |
| <i>Chrysanthemum morifolium</i> Ramat | Flowers | mg GAE/g | 3.75 \pm 0.11 | - | - | [81] |
| <i>Cichorium intybus</i> L. | Leaves | mg CAE/kg ww | 158.6 | mg QE/kg ww | 1066.0 | [83] |
| <i>Chenopodium ambrosioides</i> L. | Leaves | mg GAE/100 g dw | 822.33 \pm 12.25 | mg CE/g extract. | 768.27 \pm 10.70 | [54] |
| <i>Chondrilla juncea</i> L. | Leaves | mg GAE/g extract. | 37.66 \pm 2.44 | mg CE/g extract. | 7.43 \pm 0.28 | [72] |
| <i>Dianthus caryophyllus</i> L. | Flowers | mg GAE/g | 5.50 \pm 0.28 | - | - | [81] |
| <i>Dianthus chinensis</i> L. | Flowers | mg GAE/g | 5.27 \pm 0.25 | - | - | [81] |
| <i>Diplotaxis eruroides</i> DC. | Leaves | mg CAE/kg ww | 48.7 | mg QE/kg ww | 2876.7 | [83] |
| <i>Enhydra fluctuans</i> Lour. | - | mg GAE/g de | 70.338 \pm 0.103 | mg RE/g de | 21.759 \pm 0.039 | [31] |
| <i>Erythrina variegata</i> L. | Leaves | mg GAE/g dw | 170.33 \pm 2.18 | mg QE/g dw | 53.42 \pm 1.23 | [28] |
| <i>Erythrina variegata</i> L. | Flowers | mg GAE/g | 3.90 \pm 0.29 | - | - | [81] |
| <i>Foeniculum vulgare</i> Mill. | Leaves | mg GAE/g meth. extract. | 42.16 \pm 0.98 | mg CE/g extract. | 9.72 \pm 0.70 | [69] |
| <i>Gerbera jamesonii</i> Bolus ex Hook.f. | Flowers | mg GAE/g | 4.89 \pm 0.15 | - | - | [81] |
| <i>Gladiolus x hybridus</i> C.Morren | Flowers | mg GAE/g | 2.30 \pm 0.23 | - | - | [81] |
| <i>Glechoma hederacea</i> L. | Leaves | mg GAE/g | 196.61 \pm 6.09 | mg CE/g | 95.02 \pm 2.73 | [80] |
| <i>Helianthus annuus</i> L. | Flowers | mg GAE/g | 1.86 \pm 0.22 | - | - | [81] |
| <i>Helichrysum stoechas</i> (L.) Moench | Inflorescences | mg GAE/g | 184.42 \pm 0.35 | mg CE/g | 34.75 \pm 0.83 | [56] |
| <i>Hibiscus rosa-sinensis</i> L. | Flowers | mg GAE/g | 6.80 \pm 0.63 | - | - | [81] |
| <i>Humulus lupulus</i> L. | Leaves | mg GAE/g | 55.83 \pm 1.34 | mg CE/g | 9.56 \pm 0.65 | [69] |
| <i>Hygrophilla schulli</i> (Hamilt.) M.R.Almeida & S.M.Almeida | Leaves | mg GAE/g dw | 148.05 \pm 3.21 | mg QE/g dw | 87.12 \pm 0.86 | [28] |

Table 4. Cont.

| Plant Species | Part of Plants | Unit TPC | TPC | Unit TFC | TFC | Reference |
|---|----------------|--------------------|-------------------|------------------|------------------|-----------|
| <i>Impatiens walleriana</i> Hook.f. | Flowers | mg GAE/g | 7.62 ± 0.16 | - | - | [81] |
| <i>Ipomoea aquatica</i> Forssk. | - | mg GAE/g de | 45.449 ± 0.130 | mg RE/g de | 13.941 ± 0.040 | [31] |
| <i>Ipomoea cairica</i> (L.) Sweet | Flowers | mg GAE/g | 1.77 ± 0.13 | - | - | [81] |
| <i>Iris japonica</i> Thunb. | Flowers | mg GAE/g | 0.63 ± 0.03 | - | - | [81] |
| <i>Jasminum nudiflorum</i> Lindl. | Flowers | mg GAE/g | 3.08 ± 0.09 | - | - | [81] |
| <i>Jatropha integerrima</i> Jacq. | Flowers | mg GAE/g | 17.22 ± 0.77 | - | - | [81] |
| <i>Lantana camara</i> L. | Flowers | mg GAE/g | 3.50 ± 0.08 | - | - | [81] |
| <i>Ligustrum sinense</i> Lour. | Flowers | mg GAE/g | 6.22 ± 0.11 | - | - | [81] |
| <i>Lilium brownii</i> F.E.Br. ex Miellez | Flowers | mg GAE/g | 1.27 ± 0.13 | - | - | [81] |
| <i>Limonium sinuatum</i> (L.) Mill. | Flowers | mg GAE/g | 34.17 ± 1.17 | - | - | [81] |
| <i>Loropetalum chinense</i> var. <i>rubrum</i> Yieh | Flowers | mg GAE/g | 11.46 ± 0.26 | - | - | [81] |
| <i>Magnolia soulangeana</i> Soul.-Bod. | Flowers | mg GAE/g | 5.30 ± 0.22 | - | - | [81] |
| <i>Malva sylvestris</i> L. | Flowers | mg GAE/g extract. | 386.45 ± 8.54 | mg CE/g extract. | 210.81 ± 7.99 | [36] |
| <i>Malva viscosa</i> arboreus Cav. | Flowers | mg GAE/g | 3.12 ± 0.41 | - | - | [81] |
| <i>Matthiola incana</i> (L.) R.Br. | Flowers | mg GAE/g | 1.70 ± 0.08 | - | - | [81] |
| <i>Mentha pulegium</i> L. | Inflorescences | mg GAE/g extract. | 331.69 ± 19.63 | mg CE/g extract. | 139.85 ± 1.27 | [37] |
| <i>Montia fontana</i> L. | Leaves | mg GAE/g extract. | 75.53 ± 7.05 | mg CE/g extract. | 16.67 ± 0.62 | [69] |
| <i>Nasturtium officinale</i> R. Br. | Aerial parts | g GAE/kg extract | 87 ± 2 | g CE/kg extract | 36 ± 1 | [39] |
| <i>Oldenlandia corymbosa</i> Aiton | - | mg GAE/g de | 47.184 ± 0.060 | mg RE/g de | 15.848 ± 0.125 | [31] |
| <i>Oncidium varicosum</i> Lindl. | Flowers | mg GAE/g | 4.46 ± 0.40 | - | - | [81] |
| <i>Origanum vulgare</i> subsp. <i>virens</i> | Inflorescences | mg GAE/g | 368.58 ± 18.18 | mg CE/g | 224.15 ± 0.96 | [80] |
| <i>Orostachys fimbriata</i> (Turcz.) A. Berger | Flowers | mg GAE/g | 12.36 ± 0.43 | - | - | [81] |
| <i>Osmanthus fragrans</i> Lour. | Flowers | mg GAE/g | 16.00 ± 0.57 | - | - | [81] |
| <i>Papaver corymbosa</i> DC. | Flowers | mg GAE/g | 2.20 ± 0.07 | - | - | [81] |
| <i>Papaver rhoeas</i> L. | Leaves | mg GAE/g extract. | 25.86 ± 3.52 | mg CE/g extract. | 12.00 ± 0.46 | [72] |
| <i>Pelargonium hortorum</i> L.H. Bailey | Flowers | mg GAE/g | 25.68 ± 1.02 | - | - | [81] |
| <i>Phaseolus vulgaris</i> L. | Flowers | mg GAE/g | 1.86 ± 0.10 | - | - | [81] |
| <i>Platyodon grandiflorus</i> (Jacq.) A.DC. | Flowers | mg GAE/g | 4.57 ± 0.28 | - | - | [81] |
| <i>Portulaca oleracea</i> L. | Leaves | mg GAE/g extract. | 12.89 (7.65–20.1) | mg CE/g extract. | 1.76 (0.12–5.30) | [57] |
| <i>Pterospartum tridentatum</i> (L.) Willk. | Flowers | mg CIAE/g extract. | 523.42 ± 36.09 | mg QE/g extract. | 58.12 ± 5.78 | [41] |
| <i>Rhapniopsis indica</i> (L.) Lindl. | Flowers | mg GAE/g | 7.97 ± 0.29 | - | - | [81] |
| <i>Rhododendron simsii</i> Planch | Flowers | mg GAE/g | 6.75 ± 0.22 | - | - | [81] |

Table 4. Cont.

| Plant Species | Part of Plants | Unit TPC | TPC | Unit TFC | TFC | Reference |
|---|----------------|----------------------|----------------|---------------------|---------------|-----------|
| <i>Rhoeo discolor</i> (L'Hér.) Hance | Flowers | mg GAE/g | 2.56 ± 0.05 | - | - | [81] |
| <i>Rubus ulmifolius</i> Schott | Flowers | mg GAE/g extract. | 257.89 ± 3.28 | mg CE/g extract. | 172.45 ± 3.42 | [56] |
| <i>Rumex acetosella</i> L. | Leaves | mg GAE/g extract. | 141.58 ± 3.67 | mg CE/g extract. | 67.91 ± 3.02 | [30] |
| <i>Rumex induratus</i> Boiss. & Reut. | Leaves | mg GAE/g extract. | 117.08 ± 2.54 | mg CE/g extract. | 89.78 ± 2.81 | [30] |
| <i>Rumex papillaris</i> Boiss. & Reut. | Leaves | mg GAE/g extract. | 104.18 ± 4.17 | mg CE/g extract. | 39.49 ± 3.26 | [72] |
| <i>Rumex pulcher</i> L. | Leaves | mg GAE/g extract. | 73.44 ± 5.32 | mg CE/g extract. | 26.14 ± 0.87 | [72] |
| <i>Salvia splendens</i> Sellow ex Roem. & Schult. | Flowers | mg GAE/g | 2.57 ± 0.07 | - | - | [81] |
| <i>Sesbania sesban</i> (L.) Merr. | Leaves | mg GAE/g dw | 167.66 ± 2.37 | mg QE/g dw | 97.16 ± 1.38 | [28] |
| <i>Silene vulgaris</i> (Moench) Garcke | Leaves | mg GAE/g extract. | 26.72 ± 1.63 | mg CE/g extract. | 21.65 ± 5.53 | [69] |
| <i>Silybum marianum</i> (L.) Gaertn. | Leaves | mg GAE/g extract. | 3.72 ± 0.36 | mg CE/g extract. | 1.13 ± 0.27 | [72] |
| <i>Sinapis incana</i> (L.) Maly | Leaves | mg CAE/kg ww | 92.2 | mg QE/kg ww | 1364.7 | [83] |
| <i>Sinapis nigra</i> (L.) W.D.J.Koch | Leaves | mg CAE/kg ww | 44.3 | mg QE/kg ww | 1545.6 | [83] |
| <i>Sonchus oleraceus</i> L. | Leaves | mg GAE/g extract. | 51.33 ± 1.75 | mg CE/g extract. | 14.83 ± 0.98 | [72] |
| <i>Sophora viciifolia</i> Hance | Flowers | mg GAE/g dry extract | 143.8 ± 8.7 | mg RE/g dry extract | 237.2 ± 10.3 | [5] |
| <i>Strelitzia reginae</i> Banks ex Aiton | Flowers | mg GAE/g | 9.40 ± 0.58 | - | - | [81] |
| <i>Tamus communis</i> L. | Leaves | mg GAE/g extract. | 49.51 ± 4.07 | mg CE/g extract. | 9.33 ± 1.44 | [69] |
| <i>Taraxacum obovatum</i> (Willd.) DC. | Leaves | mg GAE/g extract. | 58.26 ± 0.90 | mg CE/g extract. | 30.03 ± 0.66 | [72] |
| <i>Thymus mastichina</i> L. | Inflorescences | mg GAE/g | 165.29 ± 1.11 | mg CE/g | 83.85 ± 1.42 | [80] |
| <i>Thymus pulegioides</i> L. | Inflorescences | mg GAE/g extract. | 210.49 ± 21.16 | mg CE/g extract. | 128.24 ± 6.00 | [37] |
| <i>Viola x Wittrockiana</i> | Flowers | mg GAE/g | 6.08 | - | - | [4] |
| <i>Wedelia trilobata</i> (L.) Hitchc. | Flowers | mg GAE/g | 3.85 ± 0.03 | - | - | [81] |
| <i>Youngia japonica</i> (L.) DC. | Flowers | mg GAE/g | 1.11 ± 0.03 | - | - | [81] |
| <i>Zantedeschia aethiopica</i> (L.) Spreng | Flowers | mg GAE/g | 3.07 ± 0.07 | - | - | [81] |

[‡] Mean value (minimum-maximum); [‡] GAE = gallic acid equivalents; [§] RE = rutin equivalents; [°] CE = catequin equivalents; [£] QE = quercetin equivalent; [¶] CAE = caffeic acid equivalent; CIAE = chlorogenic acid equivalents.

The WEP with the highest TPC was *Tamus communis* reaching 404 mg GAE/g, this plant being the only from the Diocoreaceae family [82]. Regarding the Asteraceae family (17 WEPs), the TPC ranged between 1.1 mg GAE/g (*Youngia japonica*) to 184 mg GAE/g (*Helichrysum stoechas*) [81,82]. It is necessary to highlight that the Polygonaceae family, although only data for four plants was available, showed values above the average, ranging between 73.44 mg GAE/g (*Rumex pulcher*) and 142 mg GAE/g (*Rumex acetosella*) [30,72].

Data regarding TFC (mg EC/g) were found for 31 WEPs, with *Origanum vulgare* (Lamiaceae family) and *Malva sylvestris* (Malvaceae family) having the highest values at 224 and 211 mg CE/g, respectively [36,80]. Regarding the 7 WEPs whose flavonoids content was expressed in mg RE/g, the highest flavonoids content was shown by *Sophora viciifolia*

(Fabaceae family) with values of 237 mg rutin/g dry extract [5]. On the other hand, of the 6 WEPs whose flavonoids content was expressed in mg of quercetin, three stood out for their high content: *Sesbania sesban* (Fabaceae family) > *Hygrophila schulli* (Acanthaceae family) > *Berberis aristata* (Berberidaceae family) (97.16, 87.12 and 82.05 mg QE/g, respectively) [28]. *Diplotaxis eruroides* (Brassicaceae family) and *Asparagus acutifolius* (Asparagaceae family) showed the highest content of flavonoids expressed in mg QE/kg WW, at 2877 and 2263, respectively [83].

These data show the great importance of the WEPs due to their bioactive composition, and also their great diversity, with so many families represented.

9. Economic Value of Wild Edible Plants (WEPs)

Nowadays, a fundamental aspect in society is the economy and its linkage with sustainability is essential. Therefore, the economic value of WEPs must be highlighted and studied, in which edible flowers can play a key role; they can be prepared for sale in different formats, such as fresh, dried, or even candied. This variability together with the color of these flowers makes them highly attractive and a good business proposition within the flowers market. Globalization and online marketing have created an emerging market for this type of product worldwide. The sale of these flowers is usually in punnets of between 6 and 15 flowers, depending on the type of flower and the season. The price of these punnets also depends on the type of flower, but is usually between 8 and 17 euros, some of them reaching approximately 40 euros [85–88]. In recent years, haute cuisine has valued this type of flowers, either as a decoration on dishes, or in search of new flavors, aromas or appearance. One of the best restaurants in the world is “Mugaritz” with two Michelin stars and directed by Andoni Luis Aduriz; this restaurant has been innovative and pioneering when it comes to basing its dishes on wild plants, also using edible flowers.

Apart from this direct way of selling edible flowers, there are also lollipops with edible flowers inside, or crystallized. The price range is between 21 and 60 euros [85].

Other ways in which these plants contribute their economic value is through disclosure through books. In 2004, the book “Clorofilia” by Andoni Luis Aduriz was released, where information was collected on 50 wild plants, their flowering calendar, taxonomic information and recipes [89]. Another informative book on wild plants and herbs was published by the Basque Culinary Center, which contains detailed information on 180 varieties of wild plants/herbs from a botanical and culinary point of view [90].

So far, we have studied the entire market for WEPs, either directly or through other sources; however, for many families these plants are their livelihood. In the study by Mokra et al. [91], the need to promote strategic plans at the national level in Ethiopia for the sustainable use and domestication of these plants is highlighted. This would aim at socio-economic improvement and to help in achieving one of the SDG targets (2. End hunger and malnutrition). In northeast India, WEPs are critical to the survival of ethnic communities. A survey was conducted among 30 local vendors and 550 households. The results were overwhelming, registering, in consumption or sale, five wild edible mushrooms and 158 wild plants (78.8% of them being edible). In most households, wild plants influenced family income, accounting for between 5 and 75% of family income. All these results clearly demonstrated the importance of these wild plants for the subsistence and survival of many rural communities [92].

Not only are these plants important in ethnic communities, but the study carried out by Matsuura [93] showed their importance for the Japanese population. This study indicated that, in rural areas near Fukushima, it is not possible to harvest WEPs or edible mushrooms, which precludes a complete diet full of all the essential traditional products. The results concluded that the collection rate should remain very low for a few years within a radius of between 12 and 30 km from Fukushima (Kawauchi village, where the study was carried out) due to safety reasons. These data show the importance in these communities for their livelihood.

With all these data, the economic importance of WEPs is evident, being essential for many families around the world to survive.

10. Conclusions

The topic of this review is a step into the future, because developing countries are facing a serious problem, the fast growth of their population and the consequent increase food needs/availability. Furthermore, in many rural communities it is a necessity, because WEPs are not only a food but a way of earning a living. WEPs are a natural source of minerals, vitamins, fiber, and antioxidants, and at the same time they are inexpensive, as they are not cultivated. Therefore, they are a real alternative in trying to reduce this gap between food production and demand, especially to produce natural and sustainable food additives, such as flavorings and aromas. In addition, the food industry could take advantage of the properties of WEPs to develop nutritional, organic, and sustainable foods. Nevertheless, more research is needed to go beyond just their composition information (which is the most commonly found in the scientific databases for this type of plant); information on their technological and functional properties is needed to include them in industrial processes. Toxicological studies are also needed to determine their activities (e.g., anti-mutagenic, etc.). Thus, a deep investigation into WEPs is needed to start developing commercial products based on these inexpensive and sustainable plants/ingredients. Sustainability, nutrition, and the agri-food industry converge around these plants.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/foods12051012/s1>, Table S1: List of Wild Edible Plants, WEPs; Table S2: Organic acids content in Wild Edible Plants, WEPs; Table S3: Daily value of minerals recommended by Food & Drug Administration (FDA); Table S4: Mineral elements in Wild Edible Plants, WEP's. [25,34,35,38,42,46,57,61,66–68]

Author Contributions: Conceptualization: J.C.-V., F.B., F.H. and Á.A.C.-B.; writing—original draft preparation: J.C.-V.; writing—review and editing: J.C.-V., F.B., F.H. and Á.A.C.-B.; supervision: F.H. and Á.A.C.-B. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Heywood, V. Use and Potential of Wild Plants in Farm Households. FAO Farm Systems Management Series 15. 1999. Available online: https://www.fao.org/3/W8801E/w8801e00.htm#toc_00 (accessed on 3 October 2022).
- Bacchetta, L.; Visioli, F.; Cappelli, G.; Caruso, E.; Martin, G.; Nemeth, E.; Bacchetta, G.; Bedini, G.; Wezel, A.; van Asseldonk, T.; et al. A manifesto for the valorization of wild edible plants. *J. Ethnopharmacol.* **2016**, *191*, 180–187. [CrossRef] [PubMed]
- Peña-Chocarro, L. Agricultura y alimentación vegetal en el poblado de la Edad del Bronce de Peñalosa (Baños de la Encina, Jaén). *Complutum* **2000**, *11*, 209–219.
- Da Silva, L.A.; Fischer, S.Z.; Zambiasi, R.C. Proximal composition, bioactive compounds content and color preference of *Viola x Wittrockiana* flowers. *Int. J. Gastron. Food Sci.* **2020**, *22*, 100236. [CrossRef]
- Tai, Z.; Cai, L.; Dai, L.; Dong, L.; Wang, M.; Yang, Y.; Cao, Q.; Ding, Z. Antioxidant activity and chemical constituents of edible flower of *Sophora viciifolia*. *Food Chem.* **2011**, *126*, 1648–1654. [CrossRef]
- Berihun, T.; Molla, E. Study on the Diversity and Use of Wild Edible Plants in Bullen District Northwest Ethiopia. *J. Bot.* **2017**, *2017*, 8383468. [CrossRef]
- Turreira-García, N.; Theilade, I.; Meilby, H.; Sørensen, M. Wild edible plant knowledge, distribution and transmission: A case study of the Achí Mayans of Guatemala. *J. Ethnobiol. Ethnomed.* **2015**, *11*, 52. [CrossRef]
- Svanberg, I.; Egišson, S. Edible wild plant use in the Faroe Islands and Iceland. *Acta Soc. Bot. Pol.* **2012**, *81*, 233–238. [CrossRef]
- Dewanjee, S.; Dua, T.K.; Sahu, R. Potential anti-inflammatory effect of *Leea macrophylla* Roxb. leaves: A wild edible plant. *Food Chem. Toxicol.* **2013**, *59*, 514–520. [CrossRef]
- Chen, B.; Qiu, Z. Consumers' Attitudes towards Edible Wild Plants: A Case Study of Noto Peninsula, Ishikawa Prefecture, Japan. *Int. J. For. Res.* **2012**, *2012*, 872413. [CrossRef]
- Salah, N.B.; Casabianca, H.; Jannet, H.B.; Chenavas, S.; Sanglar, C.; Fildier, A.; Bouzouita, N. Phytochemical and Biological Investigation of Two *Diplotaxis* Species Growing in Tunisia: *D. virgata* & *D. erucoides*. *Molecules* **2015**, *20*, 18128–18143. [CrossRef]

12. Schulp, C.J.E.; Thuiller, W.; Verburg, P.H. Wild food in Europe: A synthesis of knowledge and data of terrestrial wild food as an ecosystem service. *Ecol. Econ.* **2014**, *105*, 292–305. [[CrossRef](#)]
13. Clemente-Villalba, J.; Ariza, D.; García-Garvía, J.M.; Sánchez-Bravo, P.; Noguera-Artiaga, L.; Issa-Issa, H.; Hernández, F.; Carbonell-Barrachina, Á.A. Characterization and potential use of *Diplotaxis erucoides* as food ingredient for a sustainable modern cuisine and comparison with commercial mustards and wasabis. *Eur. Food Res. Technol.* **2020**, *246*, 1429–1438. [[CrossRef](#)]
14. Corrêa, R.C.G.; Di Gioia, F.; Ferreira, I.C.F.R.; Petropoulos, S.A. Wild greens used in the Mediterranean diet. In *The Mediterranean Diet*, 2nd ed.; Preedy, V.R., Watson, R.R., Eds.; Academic Press: Cambridge, MA, USA, 2020; pp. 209–228.
15. Fratianni, A.; D’Agostino, A.; Niro, S.; Bufano, A.; Paura, B.; Panfili, G. Loss or Gain of Lipophilic Bioactive Compounds in Vegetables after Domestic Cooking? Effect of Steaming and Boiling. *Foods* **2021**, *10*, 960. [[CrossRef](#)] [[PubMed](#)]
16. Guarrera, P.M.; Savo, V. Perceived health properties of wild and cultivated food plants in local and popular traditions of Italy: A review. *J. Ethnopharmacol.* **2013**, *146*, 659–680. [[CrossRef](#)]
17. Tardío, J.; Sánchez-Mata, M.d.C.; Morales, R.; Molina, M.; García-Herrera, P.; Morales, P.; Díez-Marqués, C.; Fernández-Ruiz, V.; Cámara, M.; Pardo-de-Santayana, M.; et al. Ethnobotanical and Food Composition Monographs of Selected Mediterranean Wild Edible Plants. In *Mediterranean Wild Edible Plants: Ethnobotany and Food Composition Tables*; Sánchez-Mata, M.d.C., Tardío, J., Eds.; Springer: New York, NY, USA, 2016; pp. 273–470.
18. Commission on Genetic Resources for Food and Agriculture (FAO). *Voluntary Guidelines for the Conservation and Sustainable Use of Crop Wild Relatives and Wild Food Plants*; Food and Agriculture Organization of the United Nations (FAO): Rome, Italy, 2017.
19. Page, M.J.; McKenzie, J.E.; Bossuyt, P.M.; Boutron, I.; Hoffmann, T.C.; Mulrow, C.D.; Shamseer, L.; Tetzlaff, J.M.; Akl, E.A.; Brennan, S.E.; et al. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *Syst. Rev.* **2021**, *10*, 89. [[CrossRef](#)] [[PubMed](#)]
20. Thangaraj, P. Proximate Composition Analysis. In *Pharmacological Assays of Plant-Based Natural Products*; Springer International Publishing: Cham, Switzerland, 2016; pp. 21–31.
21. Melini, V.; Melini, F. Compositional and nutritional analysis. In *Innovative Food Analysis*; Galanakis, C.M., Ed.; Academic Press: Cambridge, MA, USA, 2021; pp. 1–39.
22. Cunningham, J.; Sobolewski, R. Food composition databases for nutrition labelling: Experience from Australia. *J. Food Compos. Anal.* **2011**, *24*, 682–685. [[CrossRef](#)]
23. Chrispeels, M.J.; Morillon, R.; Maurel, C.; Gerbeau, P.; Kjellbom, P.; Johansson, I. Aquaporins of plants: Structure, function, regulation, and role in plant water relations. In *Current Topics in Membranes*; Academic Press: Cambridge, MA, USA, 2001; Volume 51, pp. 277–334.
24. Giménez, C.; Gallardo, M.; Thompson, R.B. Plant–Water Relations. In *Reference Module in Earth Systems and Environmental Sciences*; Elsevier: Amsterdam, The Netherlands, 2013.
25. García-Herrera, P.; Sánchez-Mata, M.C.; Cámara, M.; Fernández-Ruiz, V.; Díez-Marqués, C.; Molina, M.; Tardío, J. Nutrient composition of six wild edible Mediterranean Asteraceae plants of dietary interest. *J. Food Compos. Anal.* **2014**, *34*, 163–170. [[CrossRef](#)]
26. Sanchez-Bel, P.; Romojaro, A.; Egea, I.; Pretel, M.T. Wild edible plants as potential antioxidant or nutritional supplements for beverages minimally processed. *LWT Food Sci. Technol.* **2015**, *62*, 830–837. [[CrossRef](#)]
27. Barros, L.; Carvalho, A.; Ferreira, I. From famine plants to tasty and fragrant spices: Three Lamiaceae of general dietary relevance in traditional cuisine of Trás-os-Montes (Portugal). *LWT Food Sci. Technol.* **2011**, *44*, 543–548. [[CrossRef](#)]
28. Alam, M.K.; Rana, Z.H.; Islam, S.N.; Akhtaruzzaman, M. Comparative assessment of nutritional composition, polyphenol profile, antidiabetic and antioxidative properties of selected edible wild plant species of Bangladesh. *Food Chem.* **2020**, *320*, 126646. [[CrossRef](#)]
29. García-Herrera, P. *Plantas Silvestres de Consumo Tradicional en España: Caracterización de su Valor Nutricional y Estimación de su Actividad Antifúngica*; Universidad Complutense de Madrid: Madrid, Spain, 2014.
30. Pereira, C.; Barros, L.; Carvalho, A.M.; Ferreira, I.C.F.R. Nutritional composition and bioactive properties of commonly consumed wild greens: Potential sources for new trends in modern diets. *Food Res. Int.* **2011**, *44*, 2634–2640. [[CrossRef](#)]
31. Datta, S.; Sinha, B.K.; Bhattacharjee, S.; Seal, T. Nutritional composition, mineral content, antioxidant activity and quantitative estimation of water soluble vitamins and phenolics by RP-HPLC in some lesser used wild edible plants. *Heliyon* **2019**, *5*, e01431. [[CrossRef](#)]
32. Martins, D.; Barros, L.; Carvalho, A.M.; Ferreira, I.C.F.R. Nutritional and in vitro antioxidant properties of edible wild greens in Iberian Peninsula traditional diet. *Food Chem.* **2011**, *125*, 488–494. [[CrossRef](#)]
33. Ranfa, A.; Maurizi, A.; Romano, B.; Bodesmo, M. The importance of traditional uses and nutraceutical aspects of some edible wild plants in human nutrition: The case of Umbria (central Italy). *Plant Biosyst.* **2014**, *148*, 297–306. [[CrossRef](#)]
34. Gostin, A.-I.; Waisundara, V.Y. Edible flowers as functional food: A review on artichoke (*Cynara cardunculus* L.). *Trends Food Sci. Technol.* **2019**, *86*, 381–391. [[CrossRef](#)]
35. Trichopoulou, A.; Vasilopoulou, E.; Hollman, P.; Chamalides, C.; Foufa, E.; Kaloudis, T.; Kromhout, D.; Miskaki, P.; Petrochilou, I.; Poulima, E.; et al. Nutritional composition and flavonoid content of edible wild greens and green pies: A potential rich source of antioxidant nutrients in the Mediterranean diet. *Food Chem.* **2000**, *70*, 319–323. [[CrossRef](#)]
36. Barros, L.; Carvalho, A.M.; Ferreira, I.C.F.R. Leaves, flowers, immature fruits and leafy flowered stems of *Malva sylvestris*: A comparative study of the nutraceutical potential and composition. *Food Chem. Toxicol.* **2010**, *48*, 1466–1472. [[CrossRef](#)]

37. Fernandes, Â.S.F.; Barros, L.; Carvalho, A.M.; R. Ferreira, I.C.F. Lipophilic and hydrophilic antioxidants, lipid peroxidation inhibition and radical scavenging activity of two Lamiaceae food plants. *Eur. J. Lipid Sci. Technol.* **2010**, *112*, 1115–1121. [[CrossRef](#)]
38. Tardío, J.; Molina, M.; Aceituno-Mata, L.; Pardo-de-Santayana, M.; Morales, R.; Fernández-Ruiz, V.; Morales, P.; García, P.; Cámara, M.; Sánchez-Mata, M.C. *Montia fontana* L. (Portulacaceae), an interesting wild vegetable traditionally consumed in the Iberian Peninsula. *Genet. Resour. Crop Evolut.* **2011**, *58*, 1105. [[CrossRef](#)]
39. Pinela, J.; Barreira, J.C.M.; Barros, L.; Antonio, A.L.; Carvalho, A.M.; Oliveira, M.B.P.P.; Ferreira, I.C.F.R. Postharvest quality changes in fresh-cut watercress stored under conventional and inert gas-enriched modified atmosphere packaging. *Postharvest Biol. Technol.* **2016**, *112*, 55–63. [[CrossRef](#)]
40. Alu'datt, M.H.; Rababah, T.; Alhamad, M.N.; Al-Tawaha, A.; Al-Tawaha, A.R.; Gammoh, S.; Ereifej, K.I.; Al-Karaki, G.; Hamasha, H.R.; Tranchant, C.C.; et al. Herbal yield, nutritive composition, phenolic contents and antioxidant activity of purslane (*Portulaca oleracea* L.) grown in different soilless media in a closed system. *Ind. Crops Prod.* **2019**, *141*, 111746. [[CrossRef](#)]
41. Pinela, J.; Barros, L.; Carvalho, A.M.; Ferreira, I.C.F.R. Influence of the drying method in the antioxidant potential and chemical composition of four shrubby flowering plants from the tribe Genisteae (Fabaceae). *Food Chem. Toxicol.* **2011**, *49*, 2983–2989. [[CrossRef](#)] [[PubMed](#)]
42. Iyda, J.H.; Fernandes, Â.; Ferreira, F.D.; Alves, M.J.; Pires, T.; Barros, L.; Amaral, J.; Ferreira, I. Chemical composition and bioactive properties of the wild edible plant *Raphanus raphanistrum* L. *Food Res. Int.* **2019**, *121*, 714–722. [[CrossRef](#)] [[PubMed](#)]
43. Alarcón, R.; Ortiz, L.T.; García, P. Nutrient and fatty acid composition of wild edible bladder campion populations [*Silene vulgaris* (Moench.) Garcke]. *Int. J. Food Sci. Technol.* **2006**, *41*, 1239–1242. [[CrossRef](#)]
44. Guil-Guerrero, J.L.; Giménez-Giménez, A.; Rodríguez-García, I.; Torija-Isasa, M.E. Nutritional composition of *Sonchus* species (*S. asper* L., *S. oleraceus* L. and *S. tenerrimus* L.). *J. Sci. Food Agric.* **1998**, *76*, 628–632. [[CrossRef](#)]
45. Panfili, G.; Niro, S.; Bufano, A.; D'Agostino, A.; Fratianni, A.; Paura, B.; Falasca, L.; Cinquanta, L. Bioactive Compounds in Wild Asteraceae Edible Plants Consumed in the Mediterranean Diet. *Plant Foods Hum. Nutr.* **2020**, *75*, 540–546. [[CrossRef](#)]
46. Harumi Iyda, J.; Fernandes, Â.; Calhelha, R.C.; Alves, M.J.; Ferreira, F.D.; Barros, L.; Amaral, J.S.; Ferreira, I.C.F.R. Nutritional composition and bioactivity of *Umbilicus rupestris* (Salisb.) Dandy: An underexploited edible wild plant. *Food Chem.* **2019**, *295*, 341–349. [[CrossRef](#)]
47. Mennella, I. Energy Balance and Body Weight Control. In *Encyclopedia of Food Security and Sustainability*; Ferranti, P., Berry, E.M., Anderson, J.R., Eds.; Elsevier: Oxford, UK, 2019; pp. 374–377.
48. Tilman, D.; Clark, M. Global diets link environmental sustainability and human health. *Nature* **2014**, *515*, 518–522. [[CrossRef](#)]
49. Willett, W.; Rockström, J.; Loken, B.; Springmann, M.; Lang, T.; Vermeulen, S.; Garnett, T.; Tilman, D.; DeClerck, F.; Wood, A.; et al. Food in the Anthropocene: The EAT-Lancet Commission on healthy diets from sustainable food systems. *Lancet* **2019**, *393*, 447–492. [[CrossRef](#)]
50. Latham, M.C. Composición corporal, funciones de los alimentos, metabolismo y energía. In *Nutrición Humana en el Mundo en Desarrollo*; Food and Agriculture Organization of the United Nations (FAO): Rome, Italy, 2002; Volume Alimentación y nutrición.
51. Li, L.; Sheen, J. Dynamic and diverse sugar signaling. *Curr. Opin. Plant Biol.* **2016**, *33*, 116–125. [[CrossRef](#)]
52. Shah, D.; Sajjad, N.; Ali, R.; Nazir, N.; Hassan, S.; Shah, S. Sugar Regulates Plant Growth and Development Under In Vitro Conditions. In *Plant Signaling Molecules*; Khan, M.I.R., Reddy, P.S., Ferrante, A., Khan, N.A., Eds.; Woodhead Publishing: Sawston, UK, 2019; pp. 257–268.
53. Cowan, J.W.; Sakr, A.H.; Shadarevian, S.B.; Sabry, Z.I. Composition of edible wild plants of lebanon. *J. Sci. Food Agric.* **1963**, *14*, 484–488. [[CrossRef](#)]
54. Barros, L.; Pereira, E.; Calhelha, R.C.; Dueñas, M.; Carvalho, A.M.; Santos-Buelga, C.; Ferreira, I.C.F.R. Bioactivity and chemical characterization in hydrophilic and lipophilic compounds of *Chenopodium ambrosioides* L. *J. Funct. Foods* **2013**, *5*, 1732–1740. [[CrossRef](#)]
55. Barros, L.; Carvalho, A.M.; Ferreira, I.C.F.R. The nutritional composition of fennel (*Foeniculum vulgare*): Shoots, leaves, stems and inflorescences. *LWT Food Sci. Technol.* **2010**, *43*, 814–818. [[CrossRef](#)]
56. Barros, L.; Oliveira, S.; Carvalho, A.M.; Ferreira, I.C.F.R. In vitro antioxidant properties and characterization in nutrients and phytochemicals of six medicinal plants from the Portuguese folk medicine. *Ind. Crops Prod.* **2010**, *32*, 572–579. [[CrossRef](#)]
57. Petropoulos, S.A.; Karkanis, A.; Fernandes, Â.; Barros, L.; Ferreira, I.C.F.R.; Ntatsi, G.; Petrotos, K.; Lykas, C.; Khah, E. Chemical Composition and Yield of Six Genotypes of Common Purslane (*Portulaca oleracea* L.): An Alternative Source of Omega-3 Fatty Acids. *Plant Foods Hum. Nutr.* **2015**, *70*, 420–426. [[CrossRef](#)] [[PubMed](#)]
58. Chahardoli, A.; Jalilian, F.; Memariani, Z.; Farzaei, M.H.; Shokoohinia, Y. Analysis of organic acids. In *Recent Advances in Natural Products Analysis*; Sanches Silva, A., Nabavi, S.F., Saeedi, M., Nabavi, S.M., Eds.; Elsevier: Amsterdam, The Netherlands, 2020; pp. 767–823.
59. Igamberdiev, A.U.; Bykova, N.V. Role of organic acids in the integration of cellular redox metabolism and mediation of redox signalling in photosynthetic tissues of higher plants. *Free Rad. Biol. Med.* **2018**, *122*, 74–85. [[CrossRef](#)]
60. Vallarino, J.G.; Osorio, S. Organic Acids. In *Postharvest Physiology and Biochemistry of Fruits and Vegetables*; Yahia, E.M., Ed.; Woodhead Publishing: Sawston, UK, 2019; pp. 207–224.
61. Petropoulos, S.A.; Fernandes, Â.; Tzortzakakis, N.; Sokovic, M.; Ciric, A.; Barros, L.; Ferreira, I.C.F.R. Bioactive compounds content and antimicrobial activities of wild edible Asteraceae species of the Mediterranean flora under commercial cultivation conditions. *Food Res. Int.* **2019**, *119*, 859–868. [[CrossRef](#)]

62. Mesías, M.; Seiquer, I.; Delgado-Andrade, C. The Mediterranean diet and mineral composition. In *The Mediterranean Diet*, 2nd ed.; Preedy, V.R., Watson, R.R., Eds.; Academic Press: Amsterdam, The Netherlands, 2020; pp. 151–163.
63. Shankar, A.H. Mineral Deficiencies. In *Hunter's Tropical Medicine and Emerging Infectious Diseases*, 10th ed.; Ryan, E.T., Hill, D.R., Solomon, T., Aronson, N.E., Endy, T.P., Eds.; Elsevier: London, UK, 2020; pp. 1048–1054.
64. Zeece, M. Vitamins and minerals. In *Introduction to the Chemistry of Food*; Zeece, M., Ed.; Academic Press: Amsterdam, The Netherlands, 2020; pp. 163–212.
65. U.S. Food & Drugs Administration (FDA). *Interactive Nutrition Facts Label*; FDA: Silver Spring, MD, USA, 2020.
66. Guerrero, J.L.G.; Madrid, P.C.; Isasa, M.E.T. Mineral elements determination in wild edible plants. *Ecol. Food Nutr.* **1999**, *38*, 209–222. [[CrossRef](#)]
67. García-Herrera, P.; Morales, P.; Fernández-Ruiz, V.; Sánchez-Mata, M.C.; Cámara, M.; Carvalho, A.M.; Ferreira, I.C.F.R.; Pardo-de-Santayana, M.; Molina, M.; Tardío, J. Nutrients, phytochemicals and antioxidant activity in wild populations of *Allium ampeloprasum* L., a valuable underutilized vegetable. *Food Res. Int.* **2014**, *62*, 272–279. [[CrossRef](#)]
68. Disciglio, G.; Tarantino, A.; Frabboni, L.; Gagliardi, A.; Giuliani, M.M.; Tarantino, E.; Gatta, G. Qualitative characterisation of cultivated and wild edible plants: Mineral elements, phenols content and antioxidant capacity. *Ital. J. Agron.* **2017**, *12*, 383–394. [[CrossRef](#)]
69. Morales, P.; Carvalho, A.M.; Sánchez-Mata, M.C.; Cámara, M.; Molina, M.; Ferreira, I.C.F.R. Tocopherol composition and antioxidant activity of Spanish wild vegetables. *Genet. Resour. Crop Evolut.* **2012**, *59*, 851–863. [[CrossRef](#)]
70. Morales, P.; Ferreira, I.C.; Carvalho, A.M.; Sánchez-Mata, M.C.; Cámara, M.; Tardío, J. Fatty acids profiles of some Spanish wild vegetables. *Food Sci. Int.* **2012**, *18*, 281–290. [[CrossRef](#)]
71. Marrelli, M.; Russo, N.; Chiochio, I.; Statti, G.; Poli, F.; Conforti, F. Potential use in the treatment of inflammatory disorders and obesity of selected wild edible plants from Calabria region (Southern Italy). *S. Afr. J. Bot.* **2020**, *128*, 304–311. [[CrossRef](#)]
72. Morales, P.; Ferreira, I.C.F.R.; Carvalho, A.M.; Sánchez-Mata, M.C.; Cámara, M.; Fernández-Ruiz, V.; Pardo-de-Santayana, M.; Tardío, J. Mediterranean non-cultivated vegetables as dietary sources of compounds with antioxidant and biological activity. *LWT Food Sci. Technol.* **2014**, *55*, 389–396. [[CrossRef](#)]
73. Granato, D.; Santos, J.S.; Maciel, L.G.; Nunes, D.S. Chemical perspective and criticism on selected analytical methods used to estimate the total content of phenolic compounds in food matrices. *TrAC Trends Anal. Chem.* **2016**, *80*, 266–279. [[CrossRef](#)]
74. Kołton, A.; Długosz-Grochowska, O.; Wojciechowska, R.; Czaja, M. Biosynthesis Regulation of Foliates and Phenols in Plants. *Sci. Horticult.* **2022**, *291*, 110561. [[CrossRef](#)]
75. Noreen, H.; Semmar, N.; Farman, M.; McCullagh, J.S.O. Measurement of total phenolic content and antioxidant activity of aerial parts of medicinal plant *Coronopus didymus*. *Asian Pac. J. Trop. Med.* **2017**, *10*, 792–801. [[CrossRef](#)] [[PubMed](#)]
76. Cerdá-Bernad, D.; Valero-Cases, E.; Pastor, J.-J.; Frutos, M.J. Saffron bioactives crocin, crocetin and safranal: Effect on oxidative stress and mechanisms of action. *Crit. Rev. Food Sci. Nutr.* **2020**, *62*, 1–18. [[CrossRef](#)]
77. Liu, G.; Zhu, W.; Zhang, J.; Song, D.; Zhuang, L.; Ma, Q.; Yang, X.; Liu, X.; Zhang, J.; Zhang, H.; et al. Antioxidant capacity of phenolic compounds separated from tea seed oil in vitro and in vivo. *Food Chem.* **2022**, *371*, 131122. [[CrossRef](#)]
78. Radünz, M.; Hackbart, H.C.D.S.; Bona, N.P.; Pedra, N.S.; Hoffmann, J.F.; Stefanello, F.M.; Da Rosa Zavareze, E. Glucosinolates and phenolic compounds rich broccoli extract: Encapsulation by electrospraying and antitumor activity against glial tumor cells. *Colloids Surf. B Biointerfaces* **2020**, *192*, 111020. [[CrossRef](#)]
79. Villela-Castrejón, J.; Antunes-Ricardo, M.; Gutiérrez-Urbe, J.A. Bioavailability and anti-inflammatory activity of phenolic acids found in spray-dried nejayote after its in vitro digestion. *J. Funct. Foods* **2017**, *39*, 37–43. [[CrossRef](#)]
80. Barros, L.; Heleno, S.A.; Carvalho, A.M.; Ferreira, I.C.F.R. Lamiaceae often used in Portuguese folk medicine as a source of powerful antioxidants: Vitamins and phenolics. *LWT Food Sci. Technol.* **2010**, *43*, 544–550. [[CrossRef](#)]
81. Li, A.-N.; Li, S.; Li, H.-B.; Xu, D.-P.; Xu, X.-R.; Chen, F. Total phenolic contents and antioxidant capacities of 51 edible and wild flowers. *J. Funct. Foods* **2014**, *6*, 319–330. [[CrossRef](#)]
82. Pinela, J.; Carvalho, A.M.; Ferreira, I.C.F.R. Wild edible plants: Nutritional and toxicological characteristics, retrieval strategies and importance for today's society. *Food Chem. Toxicol.* **2017**, *110*, 165–188. [[CrossRef](#)]
83. Salvatore, S.; Pellegrini, N.; Brenna, O.V.; Del Rio, D.; Frasca, G.; Brighenti, F.; Tumino, R. Antioxidant Characterization of Some Sicilian Edible Wild Greens. *J. Agric. Food Chem.* **2005**, *53*, 9465–9471. [[CrossRef](#)]
84. Saleem, H.; Zengin, G.; Ahmad, I.; Htar, T.T.; Naidu, R.; Mahomoodally, M.F.; Ahemad, N. Therapeutic propensities, phytochemical composition, and toxicological evaluation of *Anagallis arvensis* (L.): A wild edible medicinal food plant. *Food Res. Int.* **2020**, *137*, 109651. [[CrossRef](#)] [[PubMed](#)]
85. Innoflower. Innoflower—Flores Comestibles. Available online: <https://www.innoflower.com/> (accessed on 9 June 2022).
86. Shopify Inc. Petit Ingredient. Available online: <https://www.petitingredient.com.au/> (accessed on 9 June 2022).
87. Flower Girl NYC. Available online: <http://www.flowergirlnyc.com/> (accessed on 9 June 2022).
88. Fine Food Specialist Limited. Fine Food Specialist. Available online: <https://www.finefoodspecialist.co.uk/> (accessed on 9 June 2022).
89. Andoni Luis Aduriz. *Clorofilia*; IXO Editorial: San Sebastian, Spain, 2004.
90. Basque Culinary Center. *Silvestre*; Planeta: Barcelona, Spain, 2022.

91. Mokria, M.; Gebretsadik, Y.; Birhane, E.; McMullin, S.; Ngethe, E.; Hadgu, K.M.; Hagazi, N.; Tewolde-Berhan, S. Nutritional and ecoclimatic importance of indigenous and naturalized wild edible plant species in Ethiopia. *Food Chem. Mol. Sci.* **2022**, *4*, 100084. [[CrossRef](#)] [[PubMed](#)]
92. Chaudhury, G.; Basumatari, M.; Darji, C.B.; Ahmed, A.F.; Borah, D.; Sah, R.K.; Devi, A.; Hazarika, N.; Dutta, G. Economic significance of wild bioresources to rural communities in the Eastern Himalayan state of Assam, Northeast India. *Trees For. People* **2021**, *5*, 100102. [[CrossRef](#)]
93. Matsuura, T. Assessment of potentially reusable edible wild plant and mushroom gathering sites in eastern Fukushima based on external radiation dose. *J. Environ. Radioact.* **2021**, *227*, 106465. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.