

Postsynaptic SNARE Proteins: Role in Synaptic Transmission and Plasticity

María Pilar Madrigal, Adrián Portalés, María Pérez SanJuan and Sandra Jurado *

Instituto de Neurociencias CSIC-UMH, 03550 San Juan de Alicante, Spain

Abstract—Soluble N-ethylmaleimide sensitive factor attachment protein receptor (SNARE) proteins mediate membrane fusion events in eukaryotic cells. Traditionally recognized as major players in regulating presynaptic neurotransmitter release, accumulative evidence over recent years has identified several SNARE proteins implicated in important postsynaptic processes such as neurotransmitter receptor trafficking and synaptic plasticity. Here we analyze the emerging data revealing this novel functional dimension for SNAREs with a focus on the molecular specialization of vesicular recycling and fusion in dendrites compared to those at axon terminals and its impact in synaptic transmission and plasticity.

This article is part of a Special Issue entitled: SNARE proteins: a long journey of science in brain physiology and pathology: from molecular. © 2018 IBRO. Published by Elsevier Ltd. All rights reserved.

Keywords: postsynaptic SNARE proteins, synaptic plasticity, neurotransmitter receptor trafficking, postsynaptic exocytosis.

INTRODUCTION

Membrane fusion is mediated by the formation of the SNARE complex assembly via the interaction of SNARE proteins. SNAREs are categorized in three families depending on their subcellular localizations. Synaptosomal-associated proteins (SNAP) and syntaxins (Stx) belong to the target SNARE (t-SNAREs) family and are located at target membranes. Synaptobrevin (Syb) (a.k.a VAMP, from vesicle-associated membrane protein) are vesicle SNAREs (v-SNARE) enriched in vesicle membranes. Syntaxin and synaptobrevins have a single copy of the conserved SNARE motif whereas SNAPs contribute with two copies. SNARE motif interaction forms tight four-helix helical bundles resistant to SDS (Sutton et al., 1998). The formation of the SNARE complex brings into close apposition the target and vesicle

membranes during an exothermic process that overcomes the energy barrier required for membrane fusion (Südhof and Rothman, 2009).

Arguably, the most extensively studied example of SNARE-dependent fusion has been the presynaptic release of neurotransmitter (Südhof (2008b)). The canonical SNARE complex in the presynaptic region is constituted by the association of synaptosomal-associated protein of 25 kDa (SNAP-25) in combination with Stx-1 and Syb-2/VAMP-2 (Jahn and Scheller, 2006; Rizo and Rosenmund, 2008; Südhof and Rothman, 2009). Calcium sensor proteins such as synaptotagmins are crucial for culminating calcium-dependent exocytosis (McMahon et al., 1995; Fernández-Chacón et al., 2001; Xu et al., 2007; Cai et al., 2008; Schonn et al., 2008; Südhof and Rothman, 2009) in a millisecond timescale. Interestingly, specific synaptotagmin isoforms have been found to be implicated in postsynaptic exocytosis during activity-dependent plasticity (Wu et al., 2017).

SNARE-dependent fusion is highly efficient thus it is unsurprising neurons employ this machinery for different membrane fusion events including dendritic exocytosis. At the postsynaptic compartment rapid changes in the composition of neurotransmitter receptors via exo- and endocytosis regulate synaptic transmission and plasticity. In the mammalian brain, excitatory transmission is mediated by glutamate receptor activation, mostly AMPA and NMDA receptors (AMPA and NMDAR) which undergo constitutive and activity-dependent trafficking. Membrane dynamics of AMPAR have been most extensively studied as an underlying mechanism of synaptic plasticity, a neuronal correlate of learning and

*Corresponding author.

E-mail address: sjurado@umh.es (S. Jurado).

Abbreviations: Aa, aminoacid; AMPAR, α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor; BoNTB, botulinum toxin serotype B; CA1, cornu ammonis 1; CNS, central nervous system; GluA1-A2, AMPA receptor subunit A1-A2; GluN2A-2B, glutamate receptor subunit 2A-2B; GABA_AR, γ -aminobutyric acid receptor subunit A; GluK5, glutamate ionotropic receptor kainate type subunit 5; IPSC, inhibitory post-synaptic currents; KARs, kainic acid receptors; KO, knock out; LTD, long-term depression; LTP, long-term potentiation; NMDAR, N-methyl-D-aspartic receptor; PSD-95, postsynaptic density-95; PICK-1, protein interacting with C kinase 1; Rab3A, ras-related protein 3A; shRNA, short hairpin RNA; SDS, sodium dodecyl sulfate; SM, Sec1/Munc18-like; SNARE, soluble N-ethylmaleimide sensitive factor attachment protein receptor; SNAP, synaptosomal-associated proteins; Stx, syntaxin; Syb, synaptobrevin; TeTx, tetanus toxin; VAMP, vesicle-associated membrane protein; VGCC, voltage-gated calcium channels.

<https://doi.org/10.1016/j.neuroscience.2018.11.012>

0306-4522/© 2018 IBRO. Published by Elsevier Ltd. All rights reserved.

memory (Malenka and Bear, 2004). Robust activation of calcium-permeable NMDAR often triggers AMPAR insertion into the plasma membrane thus eliciting long-term synaptic potentiation (LTP) (Collingridge et al., 1983; Malinow and Malenka, 2002; Brecht and Nicoll, 2003; Shepherd and Huganir, 2007; Newpher and Ehlers, 2008; Hayashi et al., 2000). AMPARs may reach synapses from perisynaptic sites (Lu et al., 2007; Petrini et al., 2009) from where they can laterally diffuse into the postsynaptic density (PSD) where are stabilized by synaptic scaffolds like PSD-95 (Henley et al., 2011; Kennedy and Ehlers, 2011; Opazo and Choquet, 2011). In the other hand, mild stimulation of NMDARs regularly yields long-term depression (LTD) of synaptic transmission which involves plasma membrane removal of AMPARs (Malenka and Bear, 2004). The elucidation of the mechanisms involved in AMPAR trafficking is crucial given the role of synaptic plasticity in experience-dependent plasticity (Malenka and Bear, 2004; Neves et al., 2008), and as a neuronal property that appears impaired in numerous neuropsychiatric and neurological disorders (Clapp et al., 2012; Ehlers, 2012).

Despite dendritic fusion events are usually triggered by elevation of calcium in the cytosol, there are important differences between pre- and postsynaptic exocytosis. For once, presynaptic terminals store multiple synaptic vesicles which are docked and primed at the plasma membrane which allows ultra-fast fusion and release upon calcium entry. In contrast, docked or primed postsynaptic vesicles have not been shown. For example, AMPAR-containing endosomes are not accumulated close to the plasma membrane. Instead of being stored at defined regions, AMPAR-containing endosomes move along the dendrite through a myosin-dependent mechanism (Correia et al., 2008; Wang et al., 2008). The lack of an active zone-like region in dendrites may contribute to the fact that AMPAR insertion events in response to NMDAR activation occurs in a time scale of several seconds or even minutes (Yudowski et al., 2007; Yang et al., 2008; Petrini et al., 2009; Patterson et al., 2010). Distinctive properties of postsynaptic release may be likely explained by the involvement of specific SNARE proteins. The composition of the postsynaptic SNARE complexes mediating synaptic transmission and plasticity has just begun to be revealed. Here, we review recent findings on postsynaptic SNARE proteins involved in neurotransmitter receptor trafficking. These new data suggest molecularly distinct SNARE complexes regulate different postsynaptic trafficking pathways providing highly selective control of synaptic transmission and plasticity.

POSTSYNAPTIC SNARE PROTEINS IN SYNAPTIC TRANSMISSION AND PLASTICITY

Role of postsynaptic synaptobrevins/VAMPs

Synaptobrevins are small transmembrane proteins located at vesicle membranes (Ernst and Brunger, 2003; Brunger et al., 2009). The main brain isoforms are Syb-1 and Syb-2, with Syb-2 being much more prominent in the forebrain than Syb-1 which is more abundant in the spinal cord and neuromuscular junctions (Elferink et al., 1989; Trimble et al., 1990; Raptis et al., 2005).

Despite being closely related to Syb-2, Syb-1 is considered to be insensitive to tetanus toxin (TeTx) and botulinum toxin serotype B (BoNTB) which specifically cleaves Syb-2 (Schiavo et al., 2000).

During early development Syb-2 can be found in dendrites due to transcytosis (Sampo et al., 2003; Wisco et al., 2003; Yap et al., 2008; Ascaño et al., 2009) suggesting that besides being an integral molecule of synaptic vesicles Syb-2 reaches postsynaptic membrane compartments. Furthermore, analysis of subcellular expression using western blot and immunogold have revealed Syb-2 in postsynaptic compartments in the CA1 hippocampus of adult rats where it appears associated with AMPAR-containing vesicles, particularly those containing the GluA1 subunit (Hussain and Davanger, 2015). In addition to its presence at AMPAR-containing endosomes, several studies have shown the involvement of Syb-2 in regulating AMPAR trafficking. In a seminal paper, Lledó and colleagues (1998) successfully disrupted LTP by postsynaptically infusing BoNTB. Further, spine growth associated with synaptic potentiation is also blocked by BoNTB (Murakoshi and Yasuda, 2012) suggesting that a Syb-2-containing SNARE complex is required for both AMPAR exocytosis during synaptic potentiation and spine growth.

More recently, experiments in cultures prepared from synaptobrevin-2 KO mice indicated that Syb-2 contributes to maintaining both synaptic and extrasynaptic GluA1-containing AMPARs (Jurado et al., 2013) (Fig. 1). According with this idea, a recent report from Hussain and Davanger (2015) has shown that Syb-2 cleavage with TeTx impairs GluA1-containing AMPARs in the postsynaptic membrane. In addition, a potential role of Syb-2 in constitutive trafficking has been hinted in a recent study showing that Syb-2 also participates in the insertion of GluA2-containing receptors (Araki et al. (2010)). These results suggest Syb-2 may be an integral element of AMPAR-containing endosomes participating in the regulation of both constitutive and activity-dependent trafficking. Furthermore, Syb-2 has also been involved in AMPAR insertion during homeostatic potentiation induced by retinoic acid (Arendt et al., 2015) and even in the exocytosis of GABA_ARs in hippocampal neurons (Gu et al., 2016) (Fig. 1). These findings suggest that similarly to synaptic vesicles (Takamori et al., 2006), Syb-2 is an abundant element of postsynaptic vesicles and thus participates in different trafficking processes in dendrites.

However, some important postsynaptic functions may be taken over by Syb-1. Whereas BoNTB treatment effectively blocks AMPAR insertion during LTP, it does not affect NMDAR function (Lledó et al., 1998; Lüscher et al., 1999). Likewise TeTx does not prevent the enhancement of NMDAR-mediated currents during glycine-induced potentiation (Lu et al., 2001). These findings suggest NMDAR insertion is mediated by a Syb isoform insensitive to BoNTB such as Syb-1. This seems to be the case for constitutive trafficking of NMDARs which has been recently shown to be significantly impaired in response to a specific Syb-1 knockdown (Gu and Huganir, 2016) (Fig. 1). These results suggest different Syb isoforms regulate specific trafficking pathways in dendrites.




Fig. 1. Functions of postsynaptic synaptobrevins. The model summarizes some of our current knowledge of the role of postsynaptic Sybs in regulating neurotransmitter receptor trafficking. Syb-2 has been suggested to be an integral part of GABA_AR-containing vesicles and regulate basal insertion in dendritic membranes. Moreover, Syb-2 may be located at AMPAR-containing endosomes, thus participating in basal and activity-dependent exocytic events either induced by NMDAR activation or during synaptic scaling induced by retinoic acid. In addition, Syb-1, another common isoform of Syb in the CNS, has been suggested to specifically regulate constitutive NMDAR exocytosis.

Role of postsynaptic syntaxins

Syntaxins are small transmembrane proteins that comprised a family of 15 members (Advani et al., 1998) from which only four isoforms (Stxs 1–4) localize to the plasma membrane where they mediate membrane fusion events (Bennett et al., 1993; Gaisano et al., 1996). The two isoforms of Stx-1 (Stx-1a and Stx-1b) are the most heavily studied due to their fundamental role in controlling vesicle docking and exocytosis of presynaptic vesicles (Jahn and Südhof, 1999). According to its function, Stx-1 is mainly expressed in presynaptic compartments (Stx-1a constitutes ~1% of all brain protein; Lang and Jahn, 2008) whereas Stxs 2, 3, and 4 have a wider tissue distribution (Bennett et al., 1993). Syntaxins cluster into membrane microdomains which may serve as hot spots for exocytosis. The formation of syntaxin-enriched

microdomains in the dendrite plasma membrane may serve as a signal for favorable regions of exocytosis that otherwise would go unnoticed in the absence of a structured molecular scaffold like in the case of the presynaptic active zone (Lang et al., 2001; Ohara-Imaizumi et al., 2004; Low et al., 2006; Sieber et al., 2007; Kennedy et al., 2010). Interestingly, complexin a protein critical for calcium-dependent exocytosis (Takahashi et al., 1999; Huang et al., 2000) has been implicated in dendritic fusion during LTP (Ahmad et al., 2012). Moreover, structure-analysis experiments showed that complexin requires the interaction with SNARE proteins to regulate LTP (Ahmad et al., 2012) suggesting the presence of a postsynaptic Stx capable of binding to complexin such as Stx-1 and Stx-3 (Pabst et al., 2000). Here we will focus on the postsynaptic function of Stxs 1, 3 and 4 as currently there are no reports investigating the role of Stx-2 in synaptic transmission and plasticity.

Postsynaptic syntaxin-1

According to their function in regulating neurotransmitter release, Stx-1 is predominantly localized at presynaptic plasma membranes (Koh et al., 1993). However a recent analysis using electron microscopy and quantitative post-embedding immunogold has revealed Stx-1 in dendritic spines (Hussain et al., 2016). Double immunofluorescence with PSD-95 and ultrastructural fractionation of synaptosomes confirmed Stx-1 localization at the PSD (Hussain et al., 2016). Intriguingly, postsynaptic Stx-1 strongly colocalized with the GluN2B subunit of NMDARs suggesting a potential role in NMDAR trafficking. This role may be subunit specific as the insertion of the GluN2A subunit has been reported to depend on Stx-4 (Gu and Haganir, 2016). By affecting GluN2B trafficking, Stx-1 may contribute to shape NMDAR-mediated currents and synaptic plasticity. Conversely, specific postsynaptic knockdown targeted to both Stx-1a and Stx-1b did not affect LTP in hippocampal slices (Jurado et al., 2013) implying that reducing endogenous Stx-1 in the postsynapse does not alter NMDAR function to block synaptic potentiation. Nevertheless, changes in NMDAR subunit composition such swapping GluN2B for GluN2A are known to modify NMDAR function during

normal development (Liu et al., 2004) and pathological conditions (Sprengel et al., 1998; Zhou and Baudry, 2006). Therefore a role of postsynaptic Stx-1 in specifically regulating GluN2B trafficking may be fundamental for maintaining normal NMDAR-mediated transmission. In addition to regulate glutamate receptor trafficking, a specific knockdown targeting both Stx-1a and Stx-1b has been shown to significantly reduce the plasma insertion frequency of γ 2-containing GABA_ARs in rat hippocampal neurons (Gu et al., 2016) suggesting a central role for postsynaptic Stx-1 in regulating constitutive trafficking of excitatory and inhibitory receptors contributing to basal excitatory/inhibitory balance (Fig. 2).

Postsynaptic syntaxin-3

Stx-3 has been recently proposed to control AMPAR exocytosis during LTP through its interaction with complexin (Jurado et al., 2013). *In vivo* stereotaxic injections of a specific shRNA against endogenous Stx-3 dramatically impaired LTP in acute hippocampal slices without affecting neither basal transmission nor presynaptic properties (Fig. 2). Moreover, the same Stx-3 shRNA effectively blocked glycine-dependent LTP in cultured hippocampal neurons and both forms of plasticity were fully rescued by restoring Stx-3 levels (Jurado et al., 2013). Furthermore, similarly to presynaptic Stx-1, Stx-3 requires the binding to SM proteins which promote a conformational change that frees the SNARE motif to interact with the rest of the SNARE proteins (Jurado et al., 2013). According to its postsynaptic function in plasticity, Stx-3 has been shown to localize widespread throughout neurons where it can be found in both dendrites and axons (Jurado et al., 2013). Interestingly, Stx-3 may play a quite distinct role in NMDAR-dependent plasticity by specifically mobilizing AMPARs since the same Stx-3 shRNA does not affect other forms of plasticity like homeostatic plasticity (Arendt et al., 2015) or other type of glutamate receptors like NMDARs (Jurado et al., 2013).

Postsynaptic syntaxin-4

Stx-4 also localizes to the plasma membrane but in contrast to Stx-1 and Stx-3 lacks the 12 aa stretch required for complexin interaction (Pabst et al., 2000). Stx-4 has been suggested to play a role in activity-dependent AMPAR exocytosis (Kennedy et al., 2010; Arendt et al., 2015) and NMDAR constitutive trafficking (Gu and Haganir, 2016) (Fig. 2). In addition, a recent study using a Stx-4 KO model revealed a significant reduction of basal glutamatergic transmission due to a reduction of both AMPARs and NMDARs (Bin et al., 2018). A function of Stx-4 in AMPAR insertion during LTP-like conditions in cultured neurons was suggested by Kennedy et al., (2010). These authors observed a block of recycling endosome exocytosis in the presence of a specific Stx-4 shRNA. These results are conflicting with the findings suggesting the role of complexin in LTP (Ahmad et al., 2012). This apparent discrepancy may in large part be explained by the differences in the methods used to assay the fusion of AMPAR-containing endosomes. In Kennedy et al., (2010), recycling endo-

somes were detected with super ecliptic pHluorin (SEP)-fused transferrin receptors as a proxy to assay AMPAR-containing endosomes which may not accurately reflect the behavior of endogenous AMPARs like those assayed by electrophysiology. In addition, it may be plausible that Stx-4 may affect AMPAR exocytosis during LTP indirectly via affecting basal levels of NMDARs (Gu and Haganir, 2016). Another way Stx-4 may affect synaptic transmission may be through a recently appreciated role in modulating transsynaptic adhesion molecules such as neuroligins (Harris et al. (2016)) which are important for synapse formation and function (Südhof (2008a)). In addition, a recent report using immunogold labeling showed endogenous Stx-4 primarily located on the plasma membrane of astrocytes in mixed neuronal cultures (Tao-Cheng et al., 2015) suggesting Stx-4 may play a role in glial exocytosis and thus impacting synaptic transmission indirectly.

Nonetheless, these results raise the intriguing possibility that different Stx isoforms may coexist in postsynaptic compartments and sort different cargos via independent membrane microdomains (Puthenveedu et al., 2010; Temkin et al., 2011). Indeed, Stx-4 has been implicated in other forms of synaptic plasticity such as retinoic acid-induced potentiation (Arendt et al., 2015). These results suggest a potential model for alternative pathways of AMPAR exocytosis in where different postsynaptic Stxs define specific exocytic sites on the postsynaptic plasma membrane. Thus during LTP, AMPAR-containing vesicles may interact with complexin and membrane microdomains enriched with Stx-3, whereas during retinoic acid-dependent plasticity AMPAR-containing vesicles non-associated to complexin may dock and be released at Stx-4 microdomains.

ROLE OF POSTSYNAPTIC SNAPS

Postsynaptic SNAP-25

SNAP proteins are Q-SNARE proteins which contribute two SNARE motifs to complete the coiled-coil assembly required for the formation of the functional SNARE complex. There are four isoforms of SNAP proteins named according to their molecular weight: SNAP-23, SNAP-25, SNAP-29 and SNAP-47, being SNAP-25 the most heavily studied due to its principal role in neurotransmitter release. SNAP-25 deletion or cleavage by specific botulinum toxins abolishes synaptic transmission without significantly affecting vesicle docking (Sørensen et al., 2003; Pantano and Montecucco, 2014). SNAP-25 also interacts with VGCCs (Catterall and Few, 2008) negatively regulating calcium sensitivity to neuronal activation (Verderio et al., 2004; Condliffe et al., 2010). In addition to its role in exocytosis, SNAP-25 contributes to clathrin-dependent endocytosis (Zhang et al., 2013) and interacts with the endocytic protein intersectin (Okamoto et al., 1999) suggesting a role in maintaining the endo and exocytosis balance in neurons.

A postsynaptic role for SNAP-25 is primarily supported by functional studies since the expression of SNAP-25 at the postsynaptic compartment remains to be definitely probed (Holderith et al., 2012; Kerti et al.,




Fig. 2. Functions of postsynaptic syntaxins. The model represents the role of different postsynaptic syntaxins in regulating neurotransmitter receptor dynamics. Despite being mostly concentrated at presynaptic terminals, Stx-1 has been suggested to play a role in regulating constitutive exocytosis of GABA_ARs, AMPARs and GluN2B-containing NMDA receptors. Interestingly, GluN2A-containing NMDARs insertion may depend on Stx-4, a Stx isoform that does not bind to complexin which mediates AMPAR exocytosis during homeostatic plasticity. In contrast Stx-3, a complexin-binding isoform is the primary regulator of AMPAR exocytosis during NMDAR-dependent LTP. These results suggest dendritic membranes may contain microdomains enriched with different Stx isoforms which define specific regions for postsynaptic receptor exocytosis.

2012; Suh et al., 2010, but also see Jordan et al., 2004; Selak et al., 2009). Manipulations targeted to impair SNAP-25 function impact glutamatergic synaptic transmission by disrupting actin cytoskeleton and spine formation (Tomasoni et al., 2013; Fossati et al., 2015). Acute *in vivo* knockdown of SNAP-25 has been shown to reduce the amount of mature dendritic spines in the CA1 region of the hippocampus (Tomasoni et al., 2013). In agreement with these data, SNAP-25 heterozygous mice show reduced levels of dendritic spines and altered PSD-95 trafficking (Fossati et al., 2015). In turn, SNAP-25 overexpression restores mature dendritic spines identified by the synaptic scaffold PSD-95. SNAP-25 regulates spine formation through an interaction with the plasma membrane and the adaptor protein p140Cap, which also inter-

acts with PSD-95. These data suggest postsynaptic SNAP-25 regulates PSD-95 clustering which in turn may affect synaptic architecture and function (Fossati et al., 2015). These results may be especially relevant for neuropsychiatric disorders like schizophrenia in which significant reductions of synaptic proteins such as SNAP-25 has been described (Yin et al., 2012).

Moreover, SNAP-25 plays a role in kainic acid receptors (KARs) and NMDARs dynamics (Selak et al., 2009; Lau et al., 2010; Jurado et al., 2013). SNAP-25 promotes the endocytosis of GluK5 subunit of KARs through a PICK1-dependent mechanisms (Selak et al., 2009), whereas the PKC-mediated phosphorylation of SNAP-25 on S187, favors NMDAR insertion to the plasma membrane via SNARE-dependent exocytosis (Lau et al., 2010; Jurado et al., 2013). Accordingly, *in vivo* knock-down of SNAP-25 impairs LTP in hippocampal synapses as a result of decreasing synaptic NMDARs required for triggering synaptic plasticity (Jurado et al., 2013). Interestingly, SNAP-25 may regulate NMDAR constitutive trafficking without affecting basal levels of AMPARs (Jurado et al., 2013; Suh et al., 2010; Gu and Huganir, 2016 but see also Gu et al., 2016) suggesting specific endosomal pathways may coexist for sorting different types of glutamatergic receptors (Fig. 3) (Fong et al., 2002; Washbourne et al., 2002; Jeyifous et al., 2009).

Postsynaptic SNAP-23

Although first described as a SNAP-25 homolog in non-neuronal tissue (Ravichandran et al., 1996), later evidence indicated SNAP-23 is also expressed in the CNS (Chen et al., 1999; Verderio et al., 2004). SNAP-23 can functionally replace SNAP-25 in *in vitro* studies, however SNAP-23-containing SNARE complexes are less stable than those constituted by SNAP-25 (Vaidyanathan et al., 2001). Nonetheless SNAP-23 is capable of sustaining important secretion processes such as insulin exocytosis (Sadoul et al., 1997). SNAP-23 has been suggested to play a role in both GABA_A and NMDA receptor trafficking (Suh et al., 2010; Gu et al., 2016). Using a heterozygous SNAP-23 knock-out mouse in combination with knockdown strategies, Suh et al., 2010 showed that reducing endogenous




Fig. 3. Functions of postsynaptic SNAREs. Schematic of the identified roles of postsynaptic SNAREs depicts membrane-bound SNAP-25 and SNAP-23 primarily involved in regulating constitutive trafficking of GABA_AR, AMPA and NMDA receptors. This function suggests the existence of specific membrane microdomains enriched with these SNAREs where exocytosis may be facilitated during baseline conditions. In contrast, activity-dependent insertion (NMDAR-dependent LTP and retinoic acid-induced potentiation) depends on the activity of SNAP-47, a broadly expressed cytosolic isoform. These results suggest that calcium influx increases dendritic exocytosis by recruiting SNARE isoforms like SNAP-47 which by not being attached to the membrane may define temporary regions for fusion along the dendritic membrane. By recruiting SNAP-47, the increase in postsynaptic exocytosis demand during synaptic plasticity can be supported in the absence of specialized regions like the presynaptic active zone.

levels of SNAP-23 significantly decreases NMDAR surface expression and NMDAR-mediated synaptic transmission in hippocampal neuronal cultures and organotypic slices. Despite affecting NMDAR constitutive trafficking in cultured conditions, reducing endogenous SNAP-23 with a specific shRNA *in vivo* failed to impair NMDAR-dependent LTP (Jurado et al., 2013) suggesting that either NMDAR trafficking relies more heavily on SNAP-23 during early stages of hippocampal development or that the remaining synaptic NMDARs are sufficient to support LTP induction in response to robust induction protocols commonly used in hippocampal slices.

In addition to glutamate receptor trafficking, SNAP-23 has also been proposed to participate in GABA_AR dynamics (Gu et al., 2016) (Fig. 3). Analysis of the insertion events corresponding to pHluorin-tagged AMPAR and GABA_AR subunits (GluA2 and γ 2, respectively) in the presence of a specific shRNA targeted to SNAP-23 has revealed that whereas SNAP-23 deletion does not impact constitutive AMPAR trafficking, surface expression of γ 2-containing GABA_AR and miniature IPSC amplitude appear significantly reduced in hippocampal neurons. These results again suggest specific roles of different postsynaptic SNARE proteins in regulating alternative sorting pathways for neurotransmitter receptors in neurons.

Postsynaptic SNAP-47

SNAP-47 is the most recent addition to the SNARE protein family and also the heaviest due to a long N-terminal stretch and an extended loop between its two SNARE motifs (Holt et al., 2006; Takamori et al., 2006). Similarly to SNAP-25, SNAP-47 is highly expressed in the CNS where it can be found in both axonal and dendritic compartments (Jurado et al., 2013; Shimojo et al., 2015; Münster-Wandowski et al., 2017). Nonetheless, SNAP-47's role in exocytosis is unclear due to the formation of labile interactions with other SNARE proteins and the lack of palmitoylated cysteine clusters, important for membrane localization (Holt et al., 2006; Kuster et al., 2015). Nonetheless, the widespread expression of SNAP-47 throughout different subcellular compartments suggests a role in trafficking and membrane fusion (Holt et al., 2006). Indeed, SNAP-47 has been shown to play a function in activity-dependent AMPAR exocytosis (Jurado et al., 2013; Arendt et al., 2015) (Fig. 3). *In vivo* SNAP-47 knockdown in the CA1 region abolishes LTP without affecting presynaptic properties or constitutive AMPAR/NMDAR trafficking highlighting a specific role of this protein in activity-dependent insertion of AMPARs (Jurado et al., 2013). In addition to this form of plasticity, SNAP-47 participates in AMPAR exocytosis during retinoic acid-induced homeostatic plasticity in hippocampal organotypic slices (Arendt et al., 2015). The weak

interactions with other SNARE proteins (Holt et al., 2006) and the lack of membrane anchor motifs may contribute to make SNAP-47 the SNAP of choice for regulating exocytosis events during LTP which seem to occur randomly throughout the postsynaptic membrane (Yudowski et al., 2007; Yang et al., 2008; Petrini et al., 2009; Patterson et al., 2010).

Postsynaptic SNAP-29

SNAP-29 is an unconventional SNAP protein which similarly to SNAP-47, lacks palmitoylation sites for membrane anchoring (Steehmaier et al., 1998). In addition, SNAP-29 contains an N-terminal motif that facilitates the interaction with endocytic adaptors and small GTPases like Rab3A (Rotem-Yehudar et al., 2001; Schardt et al., 2009). In contrast with SNAP-25 and SNAP-23, SNAP-29 has been reported to have a widespread expression throughout neurons and to interact with a wide variety of syntaxin isoforms (Steehmaier et al., 1998; Hohenstein and Roche, 2001). SNAP-29 levels in the CNS are low in comparison with its paralogs, however it has been reported that SNAP-29 mRNA expression may be regulated by electrical activity and a single session of electroconvulsive seizures on the rat frontal cortex induced an upregulation of the SNAP-29 messenger in the hippocampus (Elfvig et al., 2008). The role of SNAP-29 in fusion events is controversial and its function in brain synaptic transmission and plasticity has not yet been determined. Some evidence suggests SNAP-29 can participate in plasma membrane fusion events whereas other studies restrict its function to regulating Golgi membrane dynamics (Wong et al., 1999). However, expression of SNAP-29 in presynaptic superior cervical ganglion neurons in culture significantly inhibited synaptic transmission in an activity-dependent manner (Su et al., 2001). Although we still ignore the function of SNAP-29 in the CNS there is some evidence suggesting alterations of the *snap29* gene may be linked to neurological disorders such as schizophrenia and neuropathy associated with autophagy dysfunction (Saito et al., 2001; Morelli et al., 2014).

CONCLUDING REMARKS

Formation of SNARE complexes is a universal mechanism for membrane fusion and vesicle exocytosis. This efficient machinery regulates multiple neuronal functions being the most studied the presynaptic neurotransmitter release. However, evidence accumulated in recent years is exposing the role of SNAREs in regulating postsynaptic processes such as neurotransmitter receptor trafficking, thus expanding their role of master regulators of synaptic transmission and plasticity. Nonetheless, postsynaptic exocytosis exhibit different properties than presynaptic neurotransmitter release thus is likely supported by molecularly distinct SNARE complexes. Whereas the canonical presynaptic SNARE complex is constituted by SNAP-25, Stx-1 and Syb-2, different postsynaptic forms of exocytosis seem to depend on specific SNARE proteins. For example, AMPAR insertion during LTP

may recruit a complex formed by SNAP-47, Syb-2 and Stx-3 whereas receptors mobilized during retinoic acid-induced plasticity will be attracted to the membrane through a similar complex in which Stx-3 is replaced by Stx-4. In addition, constitutive NMDARs may require the formation of a SNARE complex constituted by SNAP-25 (which may substituted by SNAP-23), Syb-1 and Stx-4. Additionally, exocytosis of GABA_AR-containing endosomes may preferentially recruit a SNARE complex formed by SNAP-23, Syb-2 and Stx-1. These findings suggest specific exocytotic pathways coexist in dendrites in order to regulate neurotransmitter receptor insertion into the plasma membrane. Furthermore, the involvement of different Stxs opens the possibility that syntaxin-enriched microdomains mark specific regions of exocytosis for different neurotransmitter receptors.

Also intriguing is the fact that some secretion events at both the presynaptic and postsynaptic compartments seem to require the participation of the same SNARE molecules like Syb-2, however only the correct combination of SNARE molecules mediates specific exocytic events. How do neurons regulate the trafficking of SNARE molecules to maintain adequate levels to support presynaptic neurotransmitter release and postsynaptic receptor levels in dendrites? Understanding these questions will provide insight not only into the molecular basis of synaptic transmission, but also into general principles of how the function of cellular subdomains are specified.

Moreover, calcium-dependent exocytosis such as presynaptic release and postsynaptic LTP require the participation of calcium binding proteins like synaptotagmin (McMahon et al., 1995; Fernández-Chacón et al., 2001; Xu et al., 2007; Cai et al., 2008; Schonn et al., 2008; Südhof and Rothman, 2009). Indeed, a role of complexin in LTP has been exposed (Ahmad et al., 2012) suggesting the participation of a postsynaptic synaptotagmin. Interestingly, synaptotagmin-1, the major trigger of fast neurotransmitter release, is not required for LTP (Ahmad et al., 2012). However simultaneous ablation of both synaptotagmin-1 and -7, a calcium sensor for short-term facilitation (Jackman et al., 2016), blocks LTP without affecting basal synaptic transmission (Wu et al., 2017). These findings suggest postsynaptic synaptotagmin-1 and -7 can play overlapping roles in AMPAR exocytosis during LTP. Future efforts to elucidate the detailed molecular mechanisms involved in postsynaptic SNARE-dependent exocytosis involved in both synaptic transmission and plasticity will be critical for understanding the neural basis of normal and pathological brain function.

ACKNOWLEDGMENTS

Our laboratory is supported by funds from the National Institute of Aging (NIH, USA; RO1AG049937), the NARSAD Foundation (Young Investigator Award, 22688) and the Spanish State Research Agency, through the “Severo Ochoa” Programme for Centres of Excellence in R&D (SEV-2013-0317).

REFERENCES

- Advani RJ, Bae HR, Bock JB, Chao DS, Doung YC, Prekeris R, Yoo JS, Scheller RH (1998) Seven novel mammalian SNARE proteins localize to distinct membrane compartments. *J Biol Chem* 273:10317–10324.
- Ahmad M, Polepalli JS, Goswami D, Yang X, Kaeser-Woo YJ, Südhof TC, Malenka RC (2012) Postsynaptic complexin controls AMPA receptor exocytosis during LTP. *Neuron* 73:260–267.
- Araki Y, Lin DT, Haganir RL (2010) Plasma membrane insertion of the AMPA receptor GluA2 subunit is regulated by NSF binding and Q/R editing of the ion pore. *Proc Natl Acad Sci U S A* 107:11080–11085.
- Arendt KL, Zhang Y, Jurado S, Malenka RC, Südhof TC, Chen L (2015) Retinoic acid and LTP recruit postsynaptic AMPA receptors using distinct SNARE-dependent mechanisms. *Neuron* 86:442–456.
- Ascaño M, Richmond A, Borden P, Kuruvilla R (2009) Axonal targeting of Trk receptors via transcytosis regulates sensitivity to neurotrophin responses. *J Neurosci* 29:11674–11685.
- Bennett MK, García-Arrarás JE, Elferink LA, Peterson K, Fleming AM, Hazuka CD, Scheller RH (1993) The syntaxin family of vesicular transport receptors. *Cell* 74:863–873.
- Bin NR, Ma K, Harada H, Tien CW, Bergin F, Sugita K, et al. (2018) Crucial role of postsynaptic syntaxin 4 in mediating basal neurotransmission and synaptic plasticity in hippocampal CA1 neurons. *Cell Rep* 23:2955–2966.
- Bredt DS, Nicoll RA (2003) AMPA receptor trafficking at excitatory synapses. *Neuron* 40:361–379.
- Brunger AT, Weninger K, Bowen M, Chu S (2009) Single-molecule studies of the neuronal SNARE fusion machinery. *Annu Rev Biochem* 78:903–928.
- Cai H, Reim K, Varoqueaux F, Tapechum S, Hill K, Sørensen JB, Brose N, Chow RH (2008) Complexin II plays a positive role in Ca²⁺-triggered exocytosis by facilitating vesicle priming. *Proc Natl Acad Sci U S A* 105:19538–19543.
- Catterall WA, Few AP (2008) Calcium channel regulation and presynaptic plasticity. *Neuron* 59:882–901.
- Chen D, Minger SL, Honer WG, Whiteheart SW (1999) Organization of the secretory machinery in the rodent brain: distribution of the t-SNAREs, SNAP-25 and SNAP-23. *Brain Res* 831:11–24.
- Clapp WC, Hamm JP, Kirk IJ, Teyler TJ (2012) Translating long-term potentiation from animals to humans: a novel method for noninvasive assessment of cortical plasticity. *Biol Psychiatry* 71:496–502.
- Collingridge GL, Kehl SJ, McLennan H (1983) Excitatory amino acids in synaptic transmission in the Schaffer collateral-commissural pathway of the rat hippocampus. *J Physiol* 334:33–46.
- Condliffe SB, Corradini I, Pozzi D, Verderio C, Matteoli M (2010) Endogenous SNAP-25 regulates native voltage-gated calcium channels in glutamatergic neurons. *J Biol Chem* 285:24968–24976.
- Correia SS, Bassani S, Brown TC, Lisé MF, Backos DS, El-Husseini A, Passafaro M, Esteban JA (2008) Motor protein-dependent transport of AMPA receptors into spines during long-term potentiation. *Nat Neurosci* 11:457–466.
- Ehlers MD (2012) Hijacking Hebb: noninvasive methods to probe plasticity in psychiatric disease. *Biol Psychiatry* 71:484–486.
- Elferink LA, Trimble WS, Scheller RH (1989) Two vesicle-associated membrane protein genes are differentially expressed in the rat central nervous system. *J Biol Chem* 264:11061–11064.
- Elfving B, Bonefeld BE, Rosenberg R, Wegener G (2008) Differential expression of synaptic vesicle proteins after repeated electroconvulsive seizures in rat frontal cortex and hippocampus. *Synapse* 62:662–670.
- Ernst JA, Brunger AT (2003) High resolution structure, stability, and synaptotagmin binding of a truncated neuronal SNARE complex. *J Biol Chem* 278:8630–8636.
- Fernández-Chacón R, Königstorfer A, Gerber SH, García J, Matos MF, Stevens CF, Brose N, Rizo J, Rosenmund C, Südhof TC (2001) Synaptotagmin I functions as a calcium regulator of release probability. *Nature* 410:41–49.
- Fong DK, Rao A, Crump FT, Craig AM (2002) Rapid synaptic remodeling by protein kinase C: reciprocal translocation of NMDA receptors and calcium/calmodulin-dependent kinase II. *J Neurosci* 22:2153–2164.
- Fossati G, Morini R, Corradini I, Antonucci F, Trepte P, Edry E, et al. (2015) Reduced SNAP-25 increases PSD-95 mobility and impairs spine morphogenesis. *Cell Death Differ* 22:1425–1436.
- Gaisano HY, Ghai M, Malkus PN, Sheu L, Bouquillon A, Bennett MK, et al. (1996) Distinct cellular locations of the syntaxin family of proteins in rat pancreatic acinar cells. *Mol Biol Cell* 7:2019–2027.
- Gu Y, Chiu SL, Liu B, Wu PH, Delannoy M, Lin DT, Wirtz D, Haganir RL (2016) Differential vesicular sorting of AMPA and GABA_A receptors. *Proc Natl Acad Sci U S A* 113:E922–E931.
- Gu Y, Haganir RL (2016) Identification of the SNARE complex mediating the exocytosis of NMDA receptors. *Proc Natl Acad Sci U S A* 113:12280–12285.
- Harris KP, Zhang YV, Piccoli ZD, Perrimon N, Littleton JT (2016) The postsynaptic t-SNARE Syntaxin 4 controls traffic of Neuroigin 1 and Synaptotagmin 4 to regulate retrograde signaling. *eLife* 5. <https://doi.org/10.7554/eLife.13881>. pii: e13881.
- Hayashi Y, Shi SH, Esteban JA, Piccini A, Poncer JC, Malinow R (2000) Driving AMPA receptors into synapses by LTP and CaMKII: Requirement for GluR1 and PDZ domain interaction. *Science* 287:2262–2267.
- Henley JM, Barker EA, Glebov OO (2011) Routes, destinations and delays: recent advances in AMPA receptor trafficking. *Trends Neurosci* 34:258–268.
- Hohenstein AC, Roche PA (2001) SNAP-29 is a promiscuous syntaxin-binding SNARE. *Biochem Biophys Res Commun* 285:167–171.
- Holderith N, Lorincz A, Katona G, Rózsa B, Kulik A, Watanabe M, et al. (2012) Release probability of hippocampal glutamatergic terminals scales with the size of the active zone. *Nat. Neurosci* 15:988–997.
- Holt M, Varoqueaux F, Wiederhold K, Takamori S, Urlaub H, Fasshauer D, Jahn R (2006) Identification of SNAP-47, a novel Qbc-SNARE with ubiquitous expression. *J Biol Chem* 281:17076–17083.
- Huang G-Z, Ujihara H, Takahashi S, Kaba H, Yagi T, Inoue S (2000) Involvement of complexin II in synaptic plasticity in the CA1 region of the hippocampus: the use of complexin II-lacking mice. *Jpn J Pharmacol* 84:179–187.
- Hussain S, Davanger S (2015) Postsynaptic VAMP/synaptobrevin facilitates differential vesicle trafficking of GluA1 and GluA2 AMPA receptor subunits. *PLoS One* 10. doi: 10.1371/e0140868.
- Hussain S, Ringsevjen H, Egbenya DL, Skjervold TL, Davanger S (2016) SNARE protein syntaxin-1 colocalizes closely with NMDA receptor subunit NR2B in postsynaptic spines in the hippocampus. *Front Mol Neurosci* 9:10. <https://doi.org/10.3389/fnmol.2016.00010>.
- Jackman SL, Turecek J, Belinsky JE, Regehr WG (2016) The calcium sensor synaptotagmin 7 is required for synaptic facilitation. *Nature* 529:88–91.
- Jahn R, Scheller RH (2006) SNAREs—engines for membrane fusion. *Nat Rev Mol Cell Biol* 9:631–643.
- Jahn R, Südhof TC (1999) Membrane fusion and exocytosis. *Annu Rev Biochem* 68:863–911.
- Jeyifous O, Waites CL, Specht CG, Fujisawa S, Schubert M, Lin EI, Marshall J, Aoki C, et al. (2009) SAP97 and CASK mediate sorting of NMDA receptors through a previously unknown secretory pathway. *Nat Neurosci* 12:1011–1019.
- Jordan BA, Fernholz BD, Boussac M, Xu C, Grigorean G, Ziff EB, Neubert TA (2004) Identification and verification of novel rodent postsynaptic density proteins. *Mol Cell Proteomics* 3:857–871.
- Jurado S, Goswami D, Zhang Y, Molina AJ, Südhof TC, Malenka RC (2013) LTP requires a unique postsynaptic SNARE fusion machinery. *Neuron* 77:542–558.

- Kennedy MJ, Ehlers MD (2011) Mechanisms and function of dendritic exocytosis. *Neuron* 69:856–875.
- Kennedy MJ, Davison IG, Robinson CG, Ehlers MD (2010) Syntaxin-4 defines a domain for activity-dependent exocytosis in dendritic spines. *Cell* 14:524–535.
- Kerti K, Lorincz A, Nusser Z (2012) Unique somato-dendritic distribution pattern of Kv4.2 channels on hippocampal CA1 pyramidal cells. *Eur J Neurosci* 35:66–75.
- Koh S, Yamamoto A, Inoue A, Inoue Y, Akagawa K, Kawamura Y, et al. (1993) Immunoelectron microscopic localization of the HPC-1 antigen in rat cerebellum. *J Neurocytol* 22:995–1005. <https://doi.org/10.1007/bf01218356>.
- Kuster A, Nola S, Dingli F, Vacca B, Gauchy C, Beaujouan JC, Nunez M, Moncion T, Loew D, Formstecher E, Galli T, Proux-Gillardeaux V (2015) The Q-soluble N-ethylmaleimide-sensitive factor attachment protein receptor (Q-SNARE) SNAP-47 regulates trafficking of selected vesicle-associated membrane proteins (VAMPs). *J Biol Chem* 290:28056–28069.
- Lau CG, Takayasu Y, Rodenas-Ruano A, Paternain AV, Lerma J, Bennett MV, Zukin RS (2010) SNAP-25 is a target of protein kinase C phosphorylation critical to NMDA receptor trafficking. *J Neurosci* 30:242–254.
- Lang T, Bruns D, Wenzel D, Riedel D, Holroyd P, Thiele C, Jahn R (2001) SNAREs are concentrated in cholesterol-dependent clusters that define docking and fusion sites for exocytosis. *EMBO J* 20:2202–2213.
- Lang T, Jahn R (2008) Core proteins of the secretory machinery. *Handb Exp Pharmacol* 184:107–127.
- Liu XB, Murray KD, Jones EG (2004) Switching of NMDA receptor 2A and 2B subunits at thalamic and cortical synapses during early postnatal development. *J Neurosci* 24:8885–8895.
- Lledó PM, Zhang X, Südhof TC, Malenka RC, Nicoll RA (1998) Postsynaptic membrane fusion and long-term potentiation. *Science* 279:399–403.
- Low SH, Vasanthi A, Nanduri J, He M, Sharma N, Koo M, Drazba J, Weimbs T (2006) Syntaxins 3 and 4 are concentrated in separate clusters on the plasma membrane before the establishment of cell polarity. *Mol Biol Cell* 17:977–989.
- Lu J, Helton TD, Blanpied TA, Rácz B, Newpher TM, Weinberg RJ, Ehlers MD (2007) Postsynaptic positioning of endocytic zones and AMPA receptor cycling by physical coupling of dynamin-3 to Homer. *Neuron* 55:874–889.
- Lu W, Man H, Ju W, Trimble WS, MacDonald JF, Wang YT (2001) Activation of synaptic NMDA receptors induces membrane insertion of new AMPA receptors and LTP in cultured hippocampal neurons. *Neuron* 29:243–254.
- Lüscher C, Xia H, Beattie EC, Carroll RC, von Zastrow M, Malenka RC, Nicoll RA (1999) Role of AMPA receptor cycling in synaptic transmission and plasticity. *Neuron* 24:649–658.
- Malenka RC, Bear MF (2004) LTP and LTD: an embarrassment of riches. *Neuron* 44:5–21.
- Malinow R, Malenka RC (2002) AMPA receptor trafficking and synaptic plasticity. *Annu Rev Neurosci* 25:103–126.
- McMahon HT, Missler M, Li C, Südhof TC (1995) Complexins: cytosolic proteins that regulate SNAP receptor function. *Cell* 83:111–119.
- Morelli E, Ginefra P, Mastrodonato V, Beznoussenko GV, Rusten TE, Bilder D, Stenmark H, Mironov AA, Vaccari T (2014) Multiple functions of the SNARE protein Snap29 in autophagy, endocytic, and exocytic trafficking during epithelial formation in *Drosophila*. *Autophagy* 10:2251–2268.
- Münster-Wandowski A, Heilmann H, Bolduan F, Trimbuch T, Yanagawa Y, Vida I (2017) Distinct localization of SNAP47 protein in GABAergic and glutamatergic neurons in the mouse and the rat hippocampus. *Front Neuroanat* 11:56. <https://doi.org/10.3389/fnana.2017.00056>.
- Murakoshi H, Yasuda R (2012) Postsynaptic signaling during plasticity of dendritic spines. *Trends Neurosci* 35:135–143.
- Neves G, Cooke SF, Bliss TV (2008) Synaptic plasticity, memory and the hippocampus: a neural network approach to causality. *Nat Rev Neurosci* 9:65–75.
- Newpher TM, Ehlers MD (2008) Glutamate receptor dynamics in dendritic microdomains. *Neuron* 58:472–497.
- Ohara-Imaizumi M, Nishiwaki C, Kikuta T, Kumakura K, Nakamichi Y, Nagamatsu S (2004) Site of docking and fusion of insulin secretory granules in live MIN6 beta cells analyzed by TAT-conjugated anti-syntaxin 1 antibody and total internal reflection fluorescence microscopy. *J Biol Chem* 279:8403–8408.
- Okamoto M, Schoch S, Südhof TC (1999) EHS1/intersectin, a protein that contains EH and SH3 domains and binds to dynamin and SNAP-25. A protein connection between exocytosis and endocytosis? *J Biol Chem* 274:18446–18454.
- Opazo P, Choquet D (2011) A three-step model for the synaptic recruitment of AMPA receptors. *Mol Cell Neurosci* 46:1–8.
- Pabst S, Hazzard JW, Antonin W, Südhof TC, Jahn R, Rizo J, Fasshauer D (2000) Selective interaction of complexin with the neuronal SNARE complex. Determination of the binding regions. *J Biol Chem* 275:19808–19818.
- Pantano S, Montecucco C (2014) The blockade of the neurotransmitter release apparatus by botulinum neurotoxins. *Cell Mol Life Sci* 71:793–811.
- Patterson MA, Szatmari EM, Yasuda R (2010) AMPA receptors are exocytosed in stimulated spines and adjacent dendrites in a Ras-ERK-dependent manner during long-term potentiation. *Proc Natl Acad Sci U S A* 107:15951–15956.
- Petrini EM, Lu J, Cognet L, Lounis B, Ehlers MD, Choquet D (2009) Endocytic trafficking and recycling maintain a pool of mobile surface AMPA receptors required for synaptic potentiation. *Neuron* 63:92–105.
- Puthenveedu MA, Lauffer B, Temkin P, Vistein R, Carlton P, Thorn K, Taunton J, Weiner OD, Parton RG, von Zastrow M (2010) Sequence-dependent sorting of recycling proteins by actin-stabilized endosomal microdomains. *Cell* 143:761–773.
- Raptis A, Torrejon-Escribano B, Gomez de Aranda I, Blasi J (2005) Distribution of synaptobrevin/VAMP 1 and 2 in rat brain. *J Chem Neuroanat* 30:201–211.
- Ravichandran V, Chawla A, Roche PA (1996) Identification of a novel syntaxin- and synaptobrevin/VAMP-binding protein, SNAP-23, expressed in non-neuronal tissues. *J Biol Chem* 271:13300–13303.
- Rizo J, Rosenmund C (2008) Synaptic vesicle fusion. *Nat Struct Mol Biol* 15:665–674.
- Rotem-Yehudar R, Galperin E, Horowitz M (2001) Association of insulin-like growth factor 1 receptor with EHD1 and SNAP29. *J Biol Chem* 276:33054–33060.
- Sadoul K, Berger A, Niemann H, Weller U, Roche PA, Klip A, et al. (1997) SNAP-23 is not cleaved by botulinum neurotoxin E and can replace SNAP-25 in the process of insulin secretion. *J Biol Chem* 272:33023–33027.
- Saito T, Guan F, Papolos DF, Rajouria N, Fann CS, Lachman HM (2001) Polymorphism in SNAP29 gene promoter region associated with schizophrenia. *Mol Psychiatry* 6:193–201.
- Sampo B, Kaech S, Kunz S, Banker G (2003) Two distinct mechanisms target membrane proteins to the axonal surface. *Neuron* 37:611–624.
- Schardt A, Brinkmann BG, Mitkovski M, Sereda MW, Werner HB, Nave KA (2009) The SNARE protein SNAP-29 interacts with the GTPase Rab 3A: Implications for membrane trafficking in myelinating glia. *J Neurosci Res* 87:3465–3479.
- Schiavo G, Matteoli M, Montecucco C (2000) Neurotoxins affecting neuroexocytosis. *Physiol Rev* 80:717–766.
- Schonn JS, Maximov A, Lao Y, Südhof TC, Sørensen JB (2008) Synaptotagmin-1 and -7 are functionally overlapping Ca²⁺ sensors for exocytosis in adrenal chromaffin cells. *Proc Natl Acad Sci U S A* 105:3998–4003.
- Selak S, Paternain AV, Aller MI, Picó E, Rivera R, Lerma J (2009) A role for SNAP25 in internalization of kainite receptors and synaptic plasticity. *Neuron* 63:357–371.
- Shepherd JD, Huganir RL (2007) The cell biology of synaptic plasticity: AMPA receptor trafficking. *Annu Rev Cell Dev Biol* 23:613–643.

- Shimojo M, Courchet J, Pieraut S, Torabi-Rander N, Sando 3rd R, Polleux F, Maximov A (2015) SNAREs controlling vesicular release of BDNF and development of callosal axons. *Cell Rep* 7:1054–1066.
- Sieber JJ, Willig KI, Kutzner C, Gerding-Reimers C, Harke B, Donnert G, Rammner B, Eggeling C, Hell SW, Grubmüller H, et al. (2007) Anatomy and dynamics of a supramolecular membrane protein cluster. *Science* 317:1072–1076.
- Sørensen JB, Nagy G, Varoqueaux F, Nehring RB, Brose N, Wilson MC, et al. (2003) Differential control of the releasable vesicle pools by SNAP-25 splice variants and SNAP-23. *Cell* 114:75–86.
- Sprengel R, Suchanek B, Amico C, Brusa R, Burnashev N, Rozov A, et al. (1998) Importance of the intracellular domain of NR2 subunits for NMDA receptor function *in vivo*. *Cell* 92:279–289.
- Steegmaier M, Yang B, Yoo JS, Huang B, Shen M, Yu S, Luo Y, Scheller RH (1998) Three novel proteins of the syntaxin SNAP-25 family. *J Biol Chem* 273:34171–34179.
- Su Q, Mochida S, Tian JH, Mehta R, Sheng ZH (2001) SNAP-29: a general SNARE protein that inhibits SNARE disassembly and is implicated in synaptic transmission. *Proc Natl Acad Sci U S A* 98:14038–14043.
- Südhof TC (2008a) Neuroligins and neurexins link synaptic function to cognitive disease. *Nature* 455:903–911.
- Südhof TC (2008b) Neurotransmitter release. *Handb Exp Pharmacol* 184:1–21.
- Südhof TC, Rothman JE (2009) Membrane fusion: grappling with SNARE and SM proteins. *Science* 323:474–477.
- Suh YH, Terashima A, Petralia RS, Wenthold RJ, Isaac JT, Roche KW, Roche PA (2010) A neuronal role for SNAP-23 in postsynaptic glutamate receptor trafficking. *Nat Neurosci* 13:338–343.
- Sutton RB, Fasshauer D, Jahn R, Brunger AT (1998) Crystal structure of a SNARE complex involved in synaptic exocytosis at 2.4 Å resolution. *Nature* 395:347–353.
- Takahashi S, Ujihara H, Huang G-Z, Yagy KI, Sanbo M, Kaba H, Yagi T (1999) Reduced hippocampal LTP in mice lacking a presynaptic protein: complexin II. *Eur J Neurosci* 11:2359–2366.
- Takamori S, Holt M, Stenius K, Lemke EA, Grønborg M, et al. (2006) Molecular anatomy of a trafficking organelle cell. *Cell* 127:831–846.
- Tao-Cheng JH, Pham A, Yang Y, Winters CA, Gallant PE, Reese TS (2015) Syntaxin 4 is concentrated on plasma membrane of astrocytes. *Neuroscience* 286:264–271.
- Temkin P, Lauffer B, Jäger S, Cimermanic P, Krogan NJ, von Zastrow M (2011) SNX27 mediates retromer tubule entry and endosome-to-plasma membrane trafficking of signalling receptors. *Nat Cell Biol* 13:715–721.
- Tomasoni R, Repetto D, Morini R, Elia C, Gardoni F, Di Luca M, Turco E, Defilippi P, Matteoli M (2013) SNAP-25 regulates spine formation through postsynaptic binding to p140Cap. *Nat Commun* 4:2136.
- Trimble WS, Gray TS, Elferink LA, Wilson MC, Scheller RH (1990) Distinct patterns of expression of two VAMP genes within the rat brain. *J Neurosci* 10:1380–1387.
- Vaidyanathan VV, Puri N, Roche PA (2001) The last exon of SNAP-23 regulates granule exocytosis from mast cells. *J Biol Chem* 276:25101–25106.
- Verderio C, Pozzi D, Pravettoni E, Inverardi F, Schenk U, Coco S, et al. (2004) SNAP-25 modulation of calcium dynamics underlies differences in GABAergic and glutamatergic responsiveness to depolarization. *Neuron* 41:599–610.
- Wang Z, Edwards JG, Riley N, Provance Jr DW, Karcher R, Li XD, Davison IG, Ikebe M, Mercer JA, Kauer JA, Ehlers MD (2008) Myosin Vb mobilizes recycling endosomes and AMPA receptors for postsynaptic plasticity. *Cell* 135:535–548.
- Washbourne P, Thompson PM, Carta M, Costa ET, Mathews JR, Lopez-Bendito G, Molnar Z, et al. (2002) Genetic ablation of the t-SNARE SNAP-25 distinguishes mechanisms of neuroexocytosis. *Nat Neurosci* 5:19–26.
- Wong SH, Xu Y, Zhang T, Griffiths G, Lowe SL, Subramaniam VN, Seow KT, Hong W (1999) GS32, a novel Golgi SNARE of 32 kDa, interacts preferentially with syntaxin 6. *Mol Biol Cell* 10:119–134.
- Wu D, Bacaj T, Morishita W, Goswami D, Arendt KL, Xu W, Chen L, Malenka RC, Südhof TC (2017) Postsynaptic synaptotagmins mediate AMPA receptor exocytosis during LTP. *Nature* 544:316–321.
- Xu J, Mashimo T, Südhof TC (2007) Synaptotagmin-1, -2, and -9: Ca²⁺ sensors for fast release that specify distinct presynaptic properties in subsets of neurons. *Neuron* 54:567–581.
- Yang Y, Wang XB, Frerking M, Zhou Q (2008) Delivery of AMPA receptors to perisynaptic sites precedes the full expression of long-term potentiation. *Proc Natl Acad Sci U S A* 105:11388–11393.
- Yap CC, Wisco D, Kujala P, Lasiecka ZM, Cannon JT, Chang MC, Hirling H, Klumperman J, Winckler B (2008) The somatodendritic endosomal regulator NEEP21 facilitates axonal targeting of L1/NgCAM. *J Cell Biol* 180:827–842.
- Yin DM, Chen YJ, Sathyamurthy A, Xiong WC, Mei L (2012) Synaptic dysfunction in schizophrenia. *Adv Exp Med Biol* 970:493–516.
- Yudowski GA, Puthenveedu MA, Leonoudakis D, Panicker S, Thorn KS, Beattie EC, von Zastrow M (2007) Real-time imaging of discrete exocytic events mediating surface delivery of AMPA receptors. *J Neurosci* 27:11112–11121.
- Wisco D, Anderson ED, Chang MC, Norden C, Boiko T, Fölsch H, Winckler B (2003) Uncovering multiple axonal targeting pathways in hippocampal neurons. *J Cell Biol* 162:1317–1328.
- Zhang Z, Wang D, Sun T, Xu J, Chiang HC, Shin W, et al. (2013) The SNARE proteins SNAP25 and synaptobrevin are involved in endocytosis at hippocampal synapses. *J Neurosci* 33:9169–9175.
- Zhou M, Baudry M (2006) Developmental changes in NMDA neurotoxicity reflect developmental changes in subunit composition of NMDA receptors. *J Neurosci* 26:2956–2963.

(Received 17 April 2018, Accepted 10 November 2018)
(Available online 17 November 2018)