

# Antioxidant Activity and Total Phenolics Content of Montpellier Cistus (*Cistus monspeliensis*) Extracts

Mahassine Amensour<sup>1,2\*</sup>, José Angel Pérez-Alvarez<sup>3</sup>, Nadia Skali-Senhaji<sup>2</sup>, Nadia El Barnoussi<sup>1</sup>, Jamal Abrini<sup>2</sup>, Juana Fernández-López<sup>3</sup>

<sup>1</sup>Higher Institute of Nursing Professions and Health Techniques (ISPITS), Laayoune, Morocco

<sup>2</sup>Laboratory of Biology and Health, Team Biotechnology and Applied Microbiology, Department of Biology, Faculty of Science, Abdelmalek Essaadi University, Tetouan, Morocco

<sup>3</sup>IPOA Research Group (REVIV-Generalitat Valenciana and UMH-1), Agrofood Department, Miguel Hernández University Orihuela, Alicante, Spain

Email: \*mamensour@gmail.com

**How to cite this paper:** Amensour, M., Pérez-Alvarez, J.A., Skali-Senhaji, N., El Barnoussi, N., Abrini, J. and Fernández-López, J. (2024) Antioxidant Activity and Total Phenolics Content of Montpellier Cistus (*Cistus monspeliensis*) Extracts. *American Journal of Plant Sciences*, 15, 1049-1068.  
<https://doi.org/10.4236/ajps.2024.1511067>

**Received:** July 24, 2024

**Accepted:** November 18, 2024

**Published:** November 21, 2024

Copyright © 2024 by author(s) and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).  
<http://creativecommons.org/licenses/by/4.0/>



Open Access

## Abstract

To determine the antioxidant activity of the water (WE), methanol (ME), ethanol (EE) and ethyl acetate extracts (EAE) from the leaves of *Cistus monspeliensis*, several methods such as 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical-scavenging assay, 2,2'-azinobis (3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) radical cation decolorization assay, reducing power assay, metal chelating assay, Thiobarbituric acid test (TBARS) and  $\beta$ -carotene linoleic acid system assay were investigated. Because of the important roles of total phenolics and total flavonoids as antioxidants, the amounts of total phenolics and total flavonoids in the extracts were also determined. The WE and ME showed a higher scavenging activity on the DPPH and ABTS radical. Both extracts also exhibited a strong chelating effect on  $Fe^{2+}$  and performed the best in the reducing power assay. On the other hand, the WE and ME were found to have the higher total phenolic contents (23.13 and 23.25 mg GAE/g of extract respectively) and total flavonoid contents (55.08 and 47.77 mg rutin/g of extract respectively). Thus, both extracts are promising alternatives to synthetic substances as food ingredients with antioxidant activity. The ME and the WE showed the similar levels in phenolic content.

## Keywords

*Cistus monspeliensis*, Antioxidant Properties, Polyphenols, Natural Antioxidant

## 1. Introduction

Lipid peroxidation, or reaction of lipid with molecular oxygen, can be described generally as a process in which oxidants, such as free radicals, attack lipids containing carbon-carbon double bonds, particularly polyunsaturated fatty acids (PUFAs) [1]. The lipoperoxidation process has been implicated in several human diseases including atherosclerosis [2] [3], cancer [4], diabetes [5] and neurodegenerative disorders [6] [7]. Oxidation is also a significant contributor to a deteriorative reaction in food during storage and processing. Damage to cells in food products during storage can lead to serious health problems for consumers [8]. Consequently, to combat these effects, synthetic antioxidants are widely used in the food industry to neutralize free radicals that contribute to this deterioration. Common synthetic phenolic antioxidants include butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), tertiary butylhydroquinone (TBHQ), and propyl gallate (PG) [9]. However, these synthetic compounds have been linked to significant health concerns, including carcinogenicity, cytotoxicity, oxidative stress induction, and endocrine disruption [10] [11]. Therefore, the search for natural products with antioxidant activities has become a significant focus in food industry [12] [13]. Their effectiveness in combating stabilizing cellular functions under oxidative stress conditions, reducing inflammation, supporting skin health, and boosting immune function highlights their significance in nutrition [14] [15].

Many medicinal plants are excellent sources of phenolic compound which have been reported to exhibit potent antioxidant activity [16]-[19]. They have been demonstrated to have comparable activity to major biological antioxidants such as tocopherol and ascorbic acid [20]. Nevertheless, the potential antioxidant effect of phenolic compounds is directly dependent on the plant species used, the harvesting time, the growing conditions, and the type of the solvent used for extraction [21].

In the Mediterranean region, one of the most representative medicinal plants is *Cistus*. In Morocco, it is noted that *Cistus* seeds are consumed as appetizers and they also possess aphrodisiac properties [22] [23]. In Spain, tea infusions are prepared for microbial infections [24]. Generally, *Cistus* compounds are particularly appreciated for their balsamic odour, as well as for their fixative properties [25]. The principal active constituents are polyphenolic compounds. Based on their chemical structures, these compounds are divided into various subclasses such as phenolic acid, flavonoids, tannins, anthocyanin [26]-[29]. The essential oil composition is chemically complex, revealing the presence of monoterpenes, sesquiterpenes hydrocarbon, oxygenated diterpenes, diterpenes hydrocarbon and oxygenated sesquiterpenes [30] [31].

One of the most common *Cistus* species in the Mediterranean basin is *Cistus monspeliensis* L. This medicinal plant is an aromatic shrub and is widespread in Morocco, except in desert areas [22]. The leaves are evergreen, elongated and highly aromatic, while the flowers boast five white petals and the fruits are small capsules. Extracts obtained from *C. monspeliensis* have been reported to possess

significant antimicrobial [29] [32] [33], antileishmanial [34], anti-inflammatory and analgesic [35], antiproliferative and cytotoxic activity [36]. Also, the significant inhibitory effects of *C. monspeliensis* extracts against the enzymes  $\alpha$ -amylase and  $\alpha$ -glucosidase demonstrate their potential abilities to reduce the postprandial increase of blood glucose levels in diabetic patients and their capacities to prevent type 2 diabetes [37]. However, these promising results have motivated further studies on this species, studying other biological activities that have not been previously considered.

The aim of the present study was to find total phenolic and flavonoid contents of water, methanol, ethanol and ethyl acetate extracts of Moroccan *C. monspeliensis*. Additionally, another goal of this study was to assess and compare the antioxidant activities of *C. monspeliensis* extract using different methods.

## 2. Materials and Methods

### 2.1. Chemical

1,1-diphenyl-2-picryl-hydrazil (DPPH) ferrozine, Folin-Ciocalteu reagent, gallic acid, rutin, aluminium chloride, 2,2'-azinobis (3-ethylbenzothiazoline-6-sulphonic acid) diammonium salt (ABTS), potassium persulphate, ascorbic acid, butylated hydroxytoluene (BHT), iron (III) chloride, iron (II) chloride, trichloroacetic acid (TCA), Trolox, linoleic acid, tween 40 and  $\beta$ -carotene were from Sigma Chemical Company (Darmstadt, Germany). The solvent used for extraction, 2-thiobarbituric acid (TBA), sodium nitrite (II) and sodium carbonate were from Merck (Darmstadt, Germany). Potassium hexacyanoferrate was from Fluka BioChemika (GmbH Bosch, Germany).

### 2.2. Plant Materials

*C. monspeliensis* (*Cistaceae*) plant was collected from Chefchaouen region (NW of Morocco) during the vegetation period and transported to the Laboratory of Biology and Health in the Abdelmalek Essaadi University. The taxonomic identification of plant material was confirmed by Dr. A. Ennabili (N° INP599 from The National Institute of Medicinal and Aromatic Plants, Sidi Mohamed Ben Abdellah University, Fés, Morocco). Collected plant materials were immediately dried in an oven (Selecta, Barcelona, Spain) at 35°C for 3 days. The leaves of plants were separated from the stem, and ground in a grinder (Moulinex, France).

### 2.3. Preparation of Extracts

Dried powders of leaves from *C. monspeliensis* were extracted with different solvents (methanol, ethanol, ethyl acetate and water). For methanolic, ethanolic and ethyl acetate extractions, a 25 g of dried sample was extracted using 100 mL of methanol, ethanol and ethyl acetate respectively, at room temperature for 7 days. For aqueous extraction, a 25 g aliquot of dried sample was extracted using 100 mL of boiling water for 15 min. Two extraction replicates of each solvent were

prepared. The extracts were filtered from filter paper and the solvents were eliminated using a rotary evaporator (Buchi Heating Bath B-490, Buchi Rotvapor R-200) to obtain a dry extract. The extracts were stored at  $-20^{\circ}\text{C}$  until use.

#### 2.4. Determination of Total Phenolic Content (TPC)

The concentration of total phenols in extracts was measured by UV spectrophotometry, based on a colorimetric oxidation/reduction reaction. The oxidizing agent used was Folin-Ciocalteu reagent [38]. To 0.5 mL of extract, 2.5 mL of Folin-Ciocalteu reagent (10-fold diluted in water) was added and, afterwards (within a time interval from 0.5 to 8 min), 2 mL of sodium carbonate (7.5% w/v) was added. The sample was incubated for 5 min at  $50^{\circ}\text{C}$  and then cooled. For a control sample, 0.5 mL of distilled water was used. The absorbance of the resulting blue-coloured solutions was measured at 760 nm. Quantitative measurements were performed, based on a standard calibration curve of gallic acid in methanol. The mean ( $\pm\text{SD}$ ) results of triplicate analyses were expressed as mg of gallic acid equivalents (GAE) per gram of dry extract.

#### 2.5. Determination of Total Flavonoid Content (TFC)

Total flavonoid content was determined by using a method described by Sakanaka *et al.* [39]. Briefly, 0.25 mL of the extract was mixed with 1.25 mL of distilled water in a test tube, followed by addition of 75  $\mu\text{L}$  of a 5% (w/v) sodium nitrite solution. After 6 min, 150  $\mu\text{L}$  of a 10% (w/v) aluminium chloride solution was added and the mixture was allowed to stand for a further 5 min before 0.5 mL of 1 M sodium hydroxide was added. The mixture was made up to 2.5 mL with distilled water and mixed. The absorbance was measured immediately at 510 nm. Quantitative measurements were performed, based on a standard calibration curve of rutin. The mean ( $\pm\text{SD}$ ) results of triplicate analyses were expressed as mg of rutin equivalents (RE) per gram of dry extract.

#### 2.6. Determination of Antioxidant Capacity

##### 2.6.1. DPPH Radical Scavenging Capacity Assay

The free radical scavenging activity of the extracts of water, methanol, ethanol and ethyl acetate of *C. monspeliensis* leaves were measured by 1,1-diphenyl-2-picrylhydrazil (DPPH) [40]. Briefly, 0.1 mM solution of DPPH in methanol was prepared and 1 mL of this solution was added to 3 mL of plant extract (50, 100, 200, 250, 500, and 1000  $\mu\text{g}/\text{mL}$ ) and was allowed to stand at room temperature for 30 min, and then absorbance was read at 517 nm against blank samples. Lower absorbance of the reaction mixture indicated higher free radical scavenging activity. Trolox and ascorbic acid were also used as controls. The inhibition percentage of the DPPH radical was calculated from the following equation:  $\% I = [(A_{\text{DPPH}} - A_{\text{Extr}})/A_{\text{DPPH}}] * 100$ . Where  $A_{\text{DPPH}}$  is the absorbance value of the DPPH blank sample, and  $A_{\text{Extr}}$  is the absorbance value of the test solution. Each assay was carried out in triplicate.

### 2.6.2. ABTS Radical Cation Decolorization Assay

The antioxidant capacity assay was carried out using the improved ABTS radical cation decolorization assay as described by Re *et al.* [41]. ABTS<sup>+</sup> radical cation was generated by oxidation of ABTS with potassium persulfate. ABTS was dissolved in deionized water to 7 mM concentration, and mixed with 2.45 mM potassium persulfate. The reaction mixture was left to stand at room temperature in the dark for 12 - 16 h before use. The ABTS<sup>+</sup> solution was diluted with ethanol to an absorbance of  $0.700 \pm 0.020$  at 734 nm at 30°C. Then 1 mL of diluted ABTS solution was mixed with 10  $\mu$ L aliquots of plant extract at different concentrations (50, 100, 250, 500, and 1000  $\mu$ g/mL), and the absorbance at 734 nm was measured at 30°C exactly 6 min after mixing. Trolox standard solutions (concentrations from 0 to 2.5 mM) in 80% ethanol were prepared and assayed using the same conditions. Appropriate solvent blanks were run in each assay. The inhibition percentage of the ABTS radical was calculated from the following equation: % I =  $[(A_{\text{ABTS}} - A_{\text{Extr}})/A_{\text{ABTS}}] \times 100$ . Where  $A_{\text{ABTS}}$  is the absorbance value of the ABTS blank sample, and  $A_{\text{Extr}}$  is the absorbance value of the test solution. Where  $A_{\text{ABTS}}$  is the absorbance value of the ABTS blank sample, and  $A_{\text{Extr}}$  is the absorbance value of the test solution. Results were expressed also in terms of trolox equivalent antioxidant capacity (TEAC, mM trolox equivalents per mg dry extract). All determinations were carried out in triplicate.

### 2.6.3. Reducing Power Assay

The reducing power of the prepared *C. monspeliensis* extracts was determined according to the method of Oyaizu [42]. One mL of plant extracts at different concentrations (50, 100, 200, 250, 500, and 1000  $\mu$ g/mL) were mixed with 2.5 mL of phosphate buffer (0.2 M, pH 6.6) and 2.5 mL of a potassium ferricyanide solution (1% w/v). The mixture was incubated in a water bath at 50°C for 20 min. Next, 2.5 mL of a TCA solution (10%, w/v) was added, and the mixture was then centrifuged at 3000 g for 10 min. A 2.5 mL aliquot of the upper layer was combined with 2.5 mL of distilled water and 0.5 mL of a ferric chloride solution (0.1%, w/v), and the absorbance was measured at 700 nm. Increased absorbance of the reaction mixture correlates with greater reducing power. BHT was also used as control. Each assay was carried out in triplicate.

### 2.6.4. Metal Chelating Assay

Metal chelating activity was determined according to the method of Decker and Welch [43] with some modifications. Briefly, 0.5 mL of the plant extract was mixed with 0.05 mL 2 mM FeCl<sub>2</sub> and 0.1 mL 5 mM ferrozine. Total volume was diluted with 2 mL methanol. Then, the mixture was shaken vigorously and left standing at room temperature for 10 min. After the mixture had reached equilibrium, the absorbance of the solution was measured spectrophotometrically at 562 nm. The absorbance of the resulting solution was compared to that of the calibrated ethylene diamine tetra acetic acid (EDTA). Results were expressed in terms of EDTA equivalents (mM EDTA equivalents per gram of dry extract). All

determinations were carried out in triplicate.

#### 2.6.5. Tiobarbituric Acid Reactive Species Test

The method of Kuppusamy *et al.* [44] was modified, to determine the conditions which would induce a high amount of thiobarbituric acid reactive substances (TBARS). The reaction mixture for inducing lipid peroxidation contained 1 mL fowl egg yolk emulsified with 0.1 M phosphate buffer (pH 7.4), to obtain a final concentration of 25 g/L and 100  $\mu$ L of 1 mM Fe<sup>2+</sup>. *Cistus* extracts or BHT was added to the incubation mixture to test the capacity to block lipid peroxidation under these conditions. The mixture was incubated at 37°C for 1 h, afterwards it was treated with 0.5 mL of freshly prepared TCA (15%) and 1 mL of TBA (1%). The reaction tubes were kept in a boiling water bath for 10 min. Upon cooling, the tubes were centrifuged at 3500 g for 10 min to remove precipitated protein. The formation of TBARS was measured by removing 100  $\mu$ L of the supernatant and measuring its absorbance at 532 nm.

Buffered egg with Fe<sup>2+</sup> was used as control. BHT was used as the standard. The inhibition ratio was calculated from the following equation:

$$\% \text{ inhibition} = ((A_{\text{control}} - A_{\text{sample}})/A_{\text{control}}) * 100$$

where  $A_{\text{control}}$  refers to the absorbance of the control and  $A_{\text{sample}}$  is the absorbance of the sample. Each assay was carried out in triplicate.

#### 2.6.6. $\beta$ -Carotene Linoleic Acid System Assay

A stock solution of  $\beta$ -carotene linoleic acid mixture was prepared as follows: 0.5 mg  $\beta$ -carotene was dissolved in 1 mL of chloroform, and 25  $\mu$ L of linoleic acid and 200 mg of Tween 40 were added [45]. The chloroform was removed using a rotary vacuum evaporator. Then, 100 mL of distilled water was added to the residue with vigorous shaking to form an emulsion. A test sample (350  $\mu$ L of the extracts, prepared at 2 g/L concentration) was mixed with 2.5 mL of this emulsion. The reaction mixture was incubated at room temperature for 48 h for auto-oxidation, and then the absorbance was measured at 490 nm. The same procedure was repeated with BHT as positive control. Relative antioxidant activities (RAA%) of the extracts were compared with those of BHT. The assays were carried out in triplicate.

### 2.7. Statistical Analysis

All data was reported as mean  $\pm$  standard error of triplicate determinations and analyzed using one-way analysis of variance (ANOVA) with significant differences between means determined at  $p < 0.05$ .

## 3. Results and Discussion

### 3.1. Determination of Total Phenolic (TPC) and Total Flavonoid Content (TFC)

The total amount of phenolic content present in the water extract (WE), methanolic extract (ME), ethanolic extract (EE) and ethyl acetate extract (EAE) of *C.*



*monspeliensis* is shown in **Table 1**. The ME and the WE showed the similar levels (23.25 and 23.13 mg GAE/g of extract, respectively) in phenolic content ( $p < 0.05$ ). The total phenolic content in EE was 21.44 mg GAE/g of extract, while the lowest content was measured for EAE (14.38 mg GAE/g of extract). The flavonoid content of the extracts is significantly different, the WE and ME are rich of flavonoids with a value of 55.08 and 47.77 mg rutin/g of extract respectively. However, the EAE yielded the lowest ( $p < 0.05$ ) flavonoid content (23.56 mg rutin/g of extract) (**Table 1**). In our experiment, a considerable variability in phenolic and flavonoid contents was observed between extracts. Specifically, the polyphenol content is higher in WE and ME, and lower in EAE. The yield of phenolic compounds increased with increasing solvent polarity. These findings, congruous with previous studies demonstrating that the concentration of polyphenol compounds depends on the polarity of extraction solvents [46]-[50].

**Table 1.** The content of phenolic and the content of flavonoids in different extracts of leaves from *Cistus monspeliensis*.

Plant extracts	Total phenolic* (mg GAE/g of extract)	Total flavonoids* (mg rutin/g of extract)
Methanol	23.25 ± 0.17a	47.77 ± 5.79b
Ethanol	21.44 ± 0.97b	33.21 ± 1.15c
Ethyl acetate	14.38 ± 0.67c	23.56 ± 3.03d
Water	23.13 ± 0.29a	55.08 ± 3.07a

\*Values are means of triplicate determination, a-d: values within a column with different letters are significantly different ( $p < 0.05$ ).


Interestingly, the results confirmed the potential for recovering high amounts of phenolics content with antioxidant properties of different extracts of *C. monspeliensis*. Several reports highlight the contribution of phenolic compounds to the antioxidant potential of different plant extracts [51]-[53]. Phenolic compounds can play an important role in neutralizing and absorbing free radicals, efficiently inhibiting lipid peroxidation [54] [55]. For these reasons, the findings suggest that these extracts could be applied in various contexts, rich in polyphenolic compounds, to delaying, retarding, or preventing the chemical and enzymatic reactions associated with lipid oxidation.

### 3.2. DPPH Radical Scavenging Activity

The radical scavenging activities of the WE, ME, EE and EAE of *C. monspeliensis* leaves were estimated by comparing the percentage inhibition of DPPH radicals by the extracts and those of trolox and ascorbic acid (**Figure 1**).

The results indicated that the DPPH free radicals were scavenged by all plant extracts in a concentration dependent manner. At low concentrations (100 µg/mL), the antioxidant activity of WE (12.53%) was comparable to that of EE (12.62%) ( $p < 0.05$ ). In contrast, the ME exhibited higher antioxidant activity

(45.66%). The antioxidant activity levels of EAE were very low, accounting for less than 2% of the total.



**Figure 1.** Scavenging effects of leaves of *Cistus monspeliensis* extracts on DPPH radicals. Trolox and ascorbic acid were used as a positive control.

At 200 µg/mL, the ME of *C. monspeliensis* demonstrated excellent DPPH scavenging activity (87.29%), comparable to that of synthetic antioxidant standard Trolox and ascorbic acid (96.39% and 96.47% respectively), followed by the WE and EE (44.41% and 26.96% respectively).

At 250 µg/mL, the ME and WE exhibited high ( $p < 0.05$ ) scavenging activity with 92.20% and 71.90% respectively, significantly higher ( $p < 0.05$ ) than EE (41.77%). An exception was observed in the EAE, which did not show a leveling off with increasing concentration (2.98%). However, the radical scavenging activity of this extract (EAE) was much lower than the other extracts ( $p < 0.05$ ); even at higher concentration (1000 µg/mL), the scavenging activity was 13.25%. This observation is fairly common in plant extracts; many non-polar compounds are not able to act as antioxidants [56].

DPPH method has been used as one of basic screening steps for searching new antioxidant compounds in extracts from natural resources [57] [58]. Conventionally, high free radical scavenging ability is regarded as high antioxidant activity. In our study, the WE and ME exhibited the highest antioxidant activity. These variations in activity among the extracts can be explained by the contribution of compounds phenolic radical-scavenging activity which has been widely studied. Effectively, the WE and ME showed the highest phenolic and flavonoid contents, compared with EE and EAE.


### 3.3. ABTS Radical Scavenging Activity

The ABTS radical cation decolourisation test is another method widely used to assess antioxidant activity. Reduction in colour indicates reduction of ABTS radical [59].

**Figure 2** shows the percentage of inhibition on the absorbance of the ABTS<sup>+</sup>



radical of WE, ME, EE and EAE extracts from leaves of *C. monspeliensis* at various concentrations (50 - 1000  $\mu\text{g/mL}$ ). At low concentrations tested (50 - 100  $\mu\text{g/mL}$ ), the WE, ME and EE of *C. monspeliensis* exhibited very similar values (13.67% - 16.03%, 15.08% - 15.88% and 17.81% - 20.77% respectively). At dose of 500  $\mu\text{g/mL}$ , the WE and ME were highly effective ( $p < 0.05$ ) in free radical scavenging activity (52.47% and 67.69% respectively) compared to EE (31.94%). Increasing the concentration of ME and WE (1000  $\mu\text{g/mL}$ ) resulted in increased ( $p < 0.05$ ) antioxidant activity to 90.13% and to 97.82% respectively, in contrast to the 62.30% of radical inhibition of EE. In this assay, the EAE showed insignificant activity; this extract reduced the ABTS radical by 34.4% at the highest dose of 1000  $\mu\text{g/mL}$  ( $p < 0.05$ ).




**Figure 2.** Scavenging effects of leaves of *Cistus monspeliensis* extracts on ABTS radical.

In this study, the ABTS $\cdot$  scavenging activity was expressed in terms of a TEAC (trolox equivalent antioxidant capacity) value after 6 minutes of reaction time, with a higher value indicating a more potent radical-scavenging effect. The results indicate that WE (1.30 mM Trolox/mg) and the ME (1.42 mM Trolox/mg) are a much more powerful antioxidants than that of the EE (0.88 mM Trolox/mg) ( $p < 0.05$ ). This variation in the activity is related to some compounds that may not react as quickly as Trolox, resulting in low TEAC values for early measuring times [60].

### 3.4. Reducing Power Activity

In the reducing power assay, the presence of reductants in the tested samples would result in reducing  $\text{Fe}^{3+}$ /ferricyanide complex to the ferrous form ( $\text{Fe}^{2+}$ ). The  $\text{Fe}^{2+}$  can therefore be monitored by measuring the formation of Perls Prussian blue at 700 nm [61]. High absorbance indicates high reducing power. It can be

observed that the reducing power of the WE of *C. monspeliensis* was superior ( $p < 0.05$ ) than the other extracts at 50 - 500  $\mu\text{g/mL}$  (Figure 3). In fact, the reducing power of WE increased from 0.22 at 50  $\mu\text{g/mL}$  to 1.31 at 500  $\mu\text{g/mL}$ . However, the reducing power of ME increased from 0.12 at 50  $\mu\text{g/mL}$  to 0.93 at 500  $\mu\text{g/mL}$ , and that of EE and EAE increased from 0.04 at 50  $\mu\text{g/mL}$  to 0.31 at 500  $\mu\text{g/mL}$  and from 0.02 at 50  $\mu\text{g/mL}$  to 0.06 at 500  $\mu\text{g/mL}$  respectively. At a dose of 1000  $\mu\text{g/mL}$ , the reducing power of WE and ME was the most effective ( $p < 0.05$ ) (1.62 and 1.57 respectively) and significantly higher ( $p < 0.05$ ) than that EE and EAE (0.65 and 0.13 respectively). However, the reducing power of BHT was relatively more pronounced ( $p < 0.05$ ) than that of the test samples (from 0.30 at 50  $\mu\text{g/mL}$  to 1.87 at 500  $\mu\text{g/mL}$ ).



**Figure 3.** Reducing power of *Cistus monspeliensis* extracts. BHT was used as a positive control.

The reducing properties are generally associated with the presence of reductones [62] [63]. Therefore, the WE and ME of *C. monspeliensis* might contain reductones, which could react with free radicals to stabilize and terminate radical chain reactions. It has been previously reported that the reducing capacity of a compound may serve as a significant indicator of its potential antioxidant activity [64] [65].

### 3.5. Ferrous Ion-Chelating Ability

Ferrozine can quantitatively form complexes with  $\text{Fe}^{2+}$ . In the presence of ion chelating agents, the complex formation is disrupted, resulting in a decrease in the red color of the complex. Table 2 presents the ferrous ion-chelating ability of the different extracts of *C. monspeliensis* calculated as EDTA equivalents/g extract. Apparently, the WE showed the highest ( $p < 0.05$ ) ferrous iron chelating ability (130.54 mM EDTA/g extract). The second-highest abilities were observed in ME

(111.42 mM ETDA/g extract). However, the EE showed relatively lower ( $p < 0.05$ ) chelating activity (82.21 mM ETDA/g extract) when compared to the action obtained from the WE and ME, but higher ( $p < 0.05$ ) than EAE (8.81 mM ETDA/g extract).

**Table 2.** Radical-scavenging activities of *Cistus monspeliensis* extracts.

Plant extracts	ABTS* (TEAC <sub>6min</sub> mM Trolox/mg)	mM ETDA*/g extract
Methanol	1.42 ± 0.01a	111.42 ± 6.01b
Ethanol	0.88 ± 2.58c	82.21 ± 2.58c
Ethyl acetate	nd	8.81 ± 3.54d
Water	1.30 ± 0.29b	130.54 ± 7.29a


\*Values are means of triplicate determination, a-d: values within a column with different letters are significantly different ( $p < 0.05$ ), nd: Not determined.

The ferrous ion ( $Fe^{2+}$ ) is currently recognized as the more pro-oxidative form of iron [66]. Minimizing ferrous ( $Fe^{2+}$ ) ions may provide protection against oxidative damage by inhibiting the production of reactive oxygen species and lipid peroxidation. Metal ions could be inactivated and potentially inhibited by chelating agents, such as a phenolic compound [67]. This study demonstrates that the WE and ME of *C. monspeliensis* leaves have a marked capacity for iron binding, attributed to the presence of polyphenols with potent iron-chelating capacity.

### 3.6. Tiobarbituric Acid Reactive Species

**Figure 4** shows the results for the TBARS values of WE, ME, EE and EAE from *C. monspeliensis*. At 100 µg/mL, the capacity to protect lipids from oxidation of all extracts showed a percentage of inhibition lower than 30%. However, at 500 - 5000 µg/mL ( $p > 0.05$ ), WE exhibited a higher percentage of inhibition compared to the other extracts. The values ranged from 47.67% to 70.86% for WE, from 30.99% to 54.89% for ME and from 24.34% to 52.27% for EE. The result of this assay suggests that the antioxidants in the different extracts required a high concentration to have a significant activity. However, at all concentrations, BHT significantly ( $p > 0.05$ ) reduced the TBARS values compared to the different extracts, with the values ranging between 83.59% at 100 µg/mL and 93.05% at 5000 µg/mL.

Thiobarbituric acid reactive substance (TBARS) assay is another method to detect lipid oxidation. This assay measures malondialdehyde (MDA), which is a split product of an endoperoxide of unsaturated fatty acids resulting from oxidation of lipid substrates [68]. The major strategies for preventing lipid oxidation involve the use of antioxidants. In the past few years, various plant extracts containing phenolic compounds have been demonstrated to be effective antioxidants in food [69]-[72]. Therefore, the antioxidant properties of WE in this assay, which has already shown a high concentration of phenolics, suggest possible alternatives to minimize or retard oxidative deterioration in food.




**Figure 4.** Ability to inhibit lipid peroxidation of *Cistus monspeliensis* extracts. BHT was used as a positive control.

### 3.7. $\beta$ -Carotene Linoleic Acid System Activity

In this assay, antioxidant capacity is determined by measuring the inhibition of volatile organic compounds and the conjugated diene hydroperoxides arising from linoleic acid oxidation [65]. The antioxidant activity of *C. monspeliensis* extracts is shown in **Figure 5**. All extracts showed lower antioxidant activities than BHT ( $p < 0.05$ ). The antioxidant activity of *C. monspeliensis* extracts followed the order: BHT (96.01%) > EA (24.07%) > EM (20.91%) > EE (19.86%) > EEA (6.03%). Results of this assay showed significant difference ( $p < 0.05$ ) between all *C. monspeliensis* extracts and BHT. This result is not surprising, as certain antioxidant compounds demonstrate limited effectiveness in preventing the bleaching of  $\beta$ -carotene. Similar to the present results, several studies have reported that ascorbic acid, a well-known antioxidant, did not show antioxidant properties in the  $\beta$ -carotene bleaching test [73]-[75]. However, the effectiveness of phenolic antioxidants in inhibiting lipid oxidation in emulsions may vary depending on the polarity, partitioning at oil-water interface, and emulsifier type [76]. In the current body of literature related to the  $\beta$ -carotene/linoleic acid assay, the best inhibition was observed in the non-polar sub-fraction (86.07%) of the methanolic extract of *Eucalyptus sargentii* [77].

A well-known theory in the field concerns the “polar paradox” which illustrates the paradoxical behavior of antioxidants in different media and rationalizes the fact that polar antioxidants are more effective in less polar media while nonpolar antioxidants are more effective in relatively more polar media [78] [79]. Some studies have been conducted to circumvent this problem; one strategy is to significantly reduce the polarity of some molecules by grafting aliphatic chains fat, in order to adjust their position in food or cosmetic matrices [80].



**Figure 5.** Degradation rates of *Cistus monspeliensis* extracts by  $\beta$ -carotene-linoleate bleaching method. Concentration sample was 2 g/L. BHT was used as a positive control.

We found that the WE and ME of *C. monspeliensis* exhibited significant activity in DPPH and ABTS radical-scavenging activity, reducing power, and ferrous ion-chelating ability at low concentrations. However, these extracts were less effective in terms of TBARS and  $\beta$ -carotene linoleic acid system assays compared to other assays. This justifies the use of various assay approaches for the determination of antioxidant activity, as suggested in the literature [81]. Due to the complex nature of the different phytochemical classes, the antioxidant capacities of plant extracts cannot be evaluated using a single method. Thus, for an extract or fraction to be considered as a good antioxidant, it is not necessary to obtain good results in all tests.

#### 4. Conclusion

This study supports the idea that *C. monspeliensis* may be a good source of natural antioxidants for use by the food industry. WE and ME of *C. monspeliensis* show a good antioxidant activity in the *in vitro* assays applied, which can be attributed to the high phenolic and flavonoid contents. Further identification and isolation of such bioactive components could potentially clarify the antioxidant properties of *C. monspeliensis* and be further exploited for pharmaceutical and food use.

#### Acknowledgements

The authors would like to thank the Spanish Agency for International Cooperation (AECI) for financing this research project. The authors are grateful to Dr. A. Ennabili (from The National Institute of the Medicinal and Aromatic Plants, Sidi Mohamed Ben Abdellah University) for the identification of the plant.

#### Conflicts of Interest

The authors declare no conflicts of interest regarding the publication of this paper.

## References

- [1] Ayala, A., Muñoz, M.F. and Argüelles, S. (2014) Lipid Peroxidation: Production, Metabolism, and Signaling Mechanisms of Malondialdehyde and 4-Hydroxy-2-Nonenal. *Oxidative Medicine and Cellular Longevity*, **2014**, Article ID: 360438. <https://doi.org/10.1155/2014/360438>
- [2] Berliner, J.A. and Heinecke, J.W. (1996) The Role of Oxidized Lipoproteins in Atherogenesis. *Free Radical Biology and Medicine*, **20**, 707-727. [https://doi.org/10.1016/0891-5849\(95\)02173-6](https://doi.org/10.1016/0891-5849(95)02173-6)
- [3] Gianazza, E., Brioschi, M., Martinez Fernandez, A., Casalnuovo, F., Altomare, A., Aldini, G., *et al.* (2021) Lipid Peroxidation in Atherosclerotic Cardiovascular Diseases. *Antioxidants & Redox Signaling*, **34**, 49-98. <https://doi.org/10.1089/ars.2019.7955>
- [4] Lee, J., Shin, D. and Roh, J. (2023) Lipid Metabolism Alterations and Ferroptosis in Cancer: Paving the Way for Solving Cancer Resistance. *European Journal of Pharmacology*, **941**, Article ID: 175497. <https://doi.org/10.1016/j.ejphar.2023.175497>
- [5] Khan, M., Nouman, M., Hashim, H., Latif, S., Husain, S., Sattar, S., *et al.* (2023) A Correlation Biomarker between BMI and Lipid Peroxidation in Type 2 Diabetes Mellitus with and without Other Complications. *Biological and Clinical Sciences Research Journal*, **2023**, 253. <https://doi.org/10.54112/bcsrj.v2023i1.253>
- [6] Simonian, N.A. and Coyle, J.T. (1996) Oxidative Stress in Neurodegenerative Diseases. *Annual Review of Pharmacology and Toxicology*, **36**, 83-106. <https://doi.org/10.1146/annurev.pa.36.040196.000503>
- [7] Wang, Y., Xu, E., Musich, P.R. and Lin, F. (2019) Mitochondrial Dysfunction in Neurodegenerative Diseases and the Potential Countermeasure. *CNS Neuroscience & Therapeutics*, **25**, 816-824. <https://doi.org/10.1111/cns.13116>
- [8] Nasery, M., Hassanzadeh, M.K., Najaran, Z.T. and Emami, S.A. (2016) Rose (*Rosa × damascena* Mill.) Essential Oils. In: Preedy, V.R., Ed., *Essential Oils in Food Preservation, Flavor and Safety*, Elsevier, 659-665. <https://doi.org/10.1016/b978-0-12-416641-7.00075-4>
- [9] Gonçalves-Filho, D. and De Souza, D. (2022) Detection of Synthetic Antioxidants: What Factors Affect the Efficiency in the Chromatographic Analysis and in the Electrochemical Analysis? *Molecules*, **27**, Article No. 7137. <https://doi.org/10.3390/molecules27207137>
- [10] Liu, R. and Mabury, S.A. (2018) Synthetic Phenolic Antioxidants and Transformation Products in Human Sera from United States Donors. *Environmental Science & Technology Letters*, **5**, 419-423. <https://doi.org/10.1021/acs.estlett.8b00223>
- [11] Felter, S.P., Zhang, X. and Thompson, C. (2021) Butylated Hydroxyanisole: Carcinogenic Food Additive to Be Avoided or Harmless Antioxidant Important to Protect Food Supply? *Regulatory Toxicology and Pharmacology*, **121**, Article ID: 104887. <https://doi.org/10.1016/j.yrtph.2021.104887>
- [12] Lourenço, S.C., Moldão-Martins, M. and Alves, V.D. (2019) Antioxidants of Natural Plant Origins: From Sources to Food Industry Applications. *Molecules*, **24**, Article No. 4132. <https://doi.org/10.3390/molecules24224132>
- [13] Fernández-López, J., Lucas-González, R., Viuda-Martos, M., Sayas-Barberá, E., Navarro, C., Haros, C.M., *et al.* (2019) Chia (*Salvia hispanica* L.) Products as Ingredients for Reformulating Frankfurters: Effects on Quality Properties and Shelf-Life. *Meat Science*, **156**, 139-145. <https://doi.org/10.1016/j.meatsci.2019.05.028>
- [14] Guzmán, D.C., Brizuela, N.O., Mejía, G.B., Olguín, H.J., Reyes, L.S., Perez, A.V., *et al.* (2019) Effect of Cardioxane and Selenium on Lipoperoxidation and Levels of



- Dopamine in Rat Brain and Heart. *Neuroscience and Medicine*, **10**, 354-368. <https://doi.org/10.4236/nm.2019.104026>
- [15] Rahaman, M.M., Hossain, R., Herrera-Bravo, J., Islam, M.T., Atolani, O., Adeyemi, O.S., *et al.* (2023) Natural Antioxidants from Some Fruits, Seeds, Foods, Natural Products, and Associated Health Benefits: An Update. *Food Science & Nutrition*, **11**, 1657-1670. <https://doi.org/10.1002/fsn3.3217>
- [16] Gouvêa, F.d.J., de Oliveira, V.S., Mariano, B.J., Takenaka, N.A.R., Gamallo, O.D., da Silva Ferreira, M., *et al.* (2023) Natural Antioxidants as Strategy to Minimize the Presence of Lipid Oxidation Products in Canned Fish: Research Progress, Current Trends and Future Perspectives. *Food Research International*, **173**, Article ID: 113314. <https://doi.org/10.1016/j.foodres.2023.113314>
- [17] Fernández-López, J., Ponce-Martínez, A.J., Rodríguez-Párraga, J., Solivella-Poveda, A.M., Fernández-López, J.A., Viuda-Martos, M., *et al.* (2023) Beetroot Juices as Colorant in Plant-Based Minced Meat Analogues: Color, Betalain Composition and Antioxidant Activity as Affected by Juice Type. *Food Bioscience*, **56**, Article ID: 103156. <https://doi.org/10.1016/j.fbio.2023.103156>
- [18] Amensour, M., Sendra, E., Pérez-Alvarez, J.A., Skali-Senhaji, N., Abrini, J. and Fernández-López, J. (2010) Antioxidant Activity and Chemical Content of Methanol and Ethanol Extracts from Leaves of Rockrose (*Cistus ladaniferus*). *Plant Foods for Human Nutrition*, **65**, 170-178. <https://doi.org/10.1007/s11130-010-0168-2>
- [19] Gad, A.S. and Sayd, A.F. (2015) Antioxidant Properties of Rosemary and Its Potential Uses as Natural Antioxidant in Dairy Products—A Review. *Food and Nutrition Sciences*, **6**, 179-193. <https://doi.org/10.4236/fns.2015.61019>
- [20] Surai, P.F. (2013) Polyphenol Compounds in the Chicken/Animal Diet: From the Past to the Future. *Journal of Animal Physiology and Animal Nutrition*, **98**, 19-31. <https://doi.org/10.1111/jpn.12070>
- [21] Martins, N., Barros, L. and Ferreira, I.C.F.R. (2016) *In Vivo* Antioxidant Activity of Phenolic Compounds: Facts and Gaps. *Trends in Food Science & Technology*, **48**, 1-12. <https://doi.org/10.1016/j.tifs.2015.11.008>
- [22] Belakhdar, J. (1997) La pharmacopée marocaine traditionnelle. Médecine arabe ancienne et savoirs populaires. IBIS.
- [23] Youbi, A., Ouahidi, I., Mansouri, L., Daoudi, A. and Boust, D. (2016) Ethnopharmacological Survey of Plants Used for Immunological Diseases in Four Regions of Morocco. *European Journal of Medicinal Plants*, **13**, 1-24. <https://doi.org/10.9734/ejmp/2016/12946>
- [24] Morales-Soto, A., Oruna-Concha, M.J., Elmore, J.S., Barrajon-Catalan, E., Micol, V., Roldán, C., *et al.* (2015) Volatile Profile of Spanish Cistus Plants as Sources of Antimicrobials for Industrial Applications. *Industrial Crops and Products*, **74**, 425-433. <https://doi.org/10.1016/j.indcrop.2015.04.034>
- [25] Guimarães, R., Sousa, M.J. and Ferreira, I.C.F.R. (2010) Contribution of Essential Oils and Phenolics to the Antioxidant Properties of Aromatic Plants. *Industrial Crops and Products*, **32**, 152-156. <https://doi.org/10.1016/j.indcrop.2010.04.011>
- [26] Barrajon-Catalan, E., Fernandez-Arroyo, S., Roldán, C., Guillén, E., Saura, D., Segura-Carretero, A., *et al.* (2011) A Systematic Study of the Polyphenolic Composition of Aqueous Extracts Deriving from Several Cistus Genus Species: Evolutionary Relationship. *Phytochemical Analysis*, **22**, 303-312. <https://doi.org/10.1002/pca.1281>
- [27] Tomás-Menor, L., Morales-Soto, A., Barrajon-Catalan, E., Roldán-Segura, C., Segura-Carretero, A. and Micol, V. (2013) Correlation between the Antibacterial Activity and the Composition of Extracts Derived from Various Spanish Cistus Species.

- Food and Chemical Toxicology*, **55**, 313-322.  
<https://doi.org/10.1016/j.fct.2013.01.006>
- [28] Lukas, B., Jovanovic, D., Schmiderer, C., Kostas, S., Kanellis, A., Gómez Navarro, J., *et al.* (2021) Intraspecific Genetic Diversity of *Cistus creticus* L. and Evolutionary Relationships to *Cistus albidus* L. (Cistaceae): Meeting of the Generations? *Plants*, **10**, Article No. 1619. <https://doi.org/10.3390/plants10081619>
- [29] Mastino, P.M., Marchetti, M., Costa, J., Juliano, C. and Usai, M. (2021) Analytical Profiling of Phenolic Compounds in Extracts of Three Cistus Species from Sardinia and Their Potential Antimicrobial and Antioxidant Activity. *Chemistry & Biodiversity*, **18**, e2100053. <https://doi.org/10.1002/cbdv.202100053>
- [30] Mastino, P.M., Marchetti, M., Costa, J. and Usai, M. (2016) Comparison of Essential Oils from Cistus Species Growing in Sardinia. *Natural Product Research*, **31**, 299-307. <https://doi.org/10.1080/14786419.2016.1236095>
- [31] Abu-Orabi, S.T., Al-Qudah, M.A., Saleh, N.R., Bataineh, T.T., Obeidat, S.M., Al-Sheraideh, M.S., *et al.* (2020) Antioxidant Activity of Crude Extracts and Essential Oils from Flower Buds and Leaves of *Cistus creticus* and *Cistus salviifolius*. *Arabian Journal of Chemistry*, **13**, 6256-6266. <https://doi.org/10.1016/j.arabj.2020.05.043>
- [32] Rebaya, A., Igueld Belghith, S., Baghdikian, B., Leddet, V.M., Mabrouki, F., Olivier, E., Cherif, J. and Ayadi, M. (2015) Total Phenolic, Total Flavonoid, Tannin Content, and Antioxidant Capacity of *Halimium halimifolium* (Cistaceae). *Journal of Applied Pharmaceutical Science*, **5**, 52-57.
- [33] Karim, H., Boubaker, H., Askarne, L., Talibi, I., Msanda, F., Boudyach, E.H., *et al.* (2015) Antifungal Properties of Organic Extracts of Eight Cistus L. Species against Postharvest Citrus Sour Rot. *Letters in Applied Microbiology*, **62**, 16-22. <https://doi.org/10.1111/lam.12507>
- [34] Fokialakis, N., Kalpoutzakis, E., Tekwani, B.L., Khan, S.I., Kobaisy, M., Skaltsounis, A.L., *et al.* (2006) Evaluation of the Antimalarial and Antileishmanial Activity of Plants from the Greek Island of Crete. *Journal of Natural Medicines*, **61**, 38-45. <https://doi.org/10.1007/s11418-006-0013-y>
- [35] Sayah, K., Chemlal, L., Marmouzi, I., El Jemli, M., Cherrah, Y. and Faouzi, M.E.A. (2017) *In Vivo* Anti-Inflammatory and Analgesic Activities of *Cistus salviifolius* (L.) and *Cistus monspeliensis* (L.) Aqueous Extracts. *South African Journal of Botany*, **113**, 160-163. <https://doi.org/10.1016/j.sajb.2017.08.015>
- [36] Vitali, F., Pennisi, G., Attaguile, G., Savoca, F. and Tita, B. (2011) Antiproliferative and Cytotoxic Activity of Extracts from *Cistus incanus* L. and *Cistus monspeliensis* L. on Human Prostate Cell Lines. *Natural Product Research*, **25**, 188-202. <https://doi.org/10.1080/14786410802583148>
- [37] Sayah, K., Marmouzi, I., Naceiri Mrabti, H., Cherrah, Y. and Faouzi, M.E.A. (2017) Antioxidant Activity and Inhibitory Potential of *Cistus salviifolius* (L.) and *Cistus monspeliensis* (L.) Aerial Parts Extracts against Key Enzymes Linked to Hyperglycemia. *BioMed Research International*, **2017**, Article ID: 2789482. <https://doi.org/10.1155/2017/2789482>
- [38] AOCS (1990) Official Methods and Recommended Practices of the American Oil Chemists Society. 4th Edition, American Oil Chemists Society.
- [39] Sakanaka, S., Tachibana, Y. and Okada, Y. (2005) Preparation and Antioxidant Properties of Extracts of Japanese Persimmon Leaf Tea (Kakinoha-Cha). *Food Chemistry*, **89**, 569-575. <https://doi.org/10.1016/j.foodchem.2004.03.013>
- [40] Blois, M.S. (1958) Antioxidant Determinations by the Use of a Stable Free Radical. *Nature*, **181**, 1199-1200. <https://doi.org/10.1038/1811199a0>

- [41] Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M. and Rice-Evans, C. (1999) Antioxidant Activity Applying an Improved ABTS Radical Cation Decolorization Assay. *Free Radical Biology and Medicine*, **26**, 1231-1237.  
[https://doi.org/10.1016/s0891-5849\(98\)00315-3](https://doi.org/10.1016/s0891-5849(98)00315-3)
- [42] Oyaizu, M. (1986) Studies on Products of Browning Reaction. Antioxidative Activities of Products of Browning Reaction Prepared from Glucosamine. *The Japanese Journal of Nutrition and Dietetics*, **44**, 307-315.  
<https://doi.org/10.5264/eiyogakuzashi.44.307>
- [43] Decker, E.A. and Welch, B. (1990) Role of Ferritin as a Lipid Oxidation Catalyst in Muscle Food. *Journal of Agricultural and Food Chemistry*, **38**, 674-677.  
<https://doi.org/10.1021/jf00093a019>
- [44] Kuppusamy, U.R., Indran, M. and Balraj, B.R.S. (2002) Antioxidant Effects of Local Fruits and Vegetable Extracts. *Journal of Tropical Medicinal Plants*, **3**, 47-53.
- [45] Dapkevicius, A., Venskutonis, R., van Beek, T.A. and Linssen, J.P.H. (1998) Antioxidant Activity of Extracts Obtained by Different Isolation Procedures from Some Aromatic Herbs Grown in Lithuania. *Journal of the Science of Food and Agriculture*, **77**, 140-146.  
[https://doi.org/10.1002/\(sici\)1097-0010\(199805\)77:1<140::aid-jsfa18>3.0.co;2-k](https://doi.org/10.1002/(sici)1097-0010(199805)77:1<140::aid-jsfa18>3.0.co;2-k)
- [46] Bhebhe, M., Füller, T.N., Chipurura, B. and Muchuweti, M. (2015) Effect of Solvent Type on Total Phenolic Content and Free Radical Scavenging Activity of Black Tea and Herbal Infusions. *Food Analytical Methods*, **9**, 1060-1067.  
<https://doi.org/10.1007/s12161-015-0270-z>
- [47] Sepahpour, S., Selamat, J., Abdul Manap, M., Khatib, A. and Abdull Razis, A. (2018) Comparative Analysis of Chemical Composition, Antioxidant Activity and Quantitative Characterization of Some Phenolic Compounds in Selected Herbs and Spices in Different Solvent Extraction Systems. *Molecules*, **23**, Article No. 402.  
<https://doi.org/10.3390/molecules23020402>
- [48] Venkatesan, T., Choi, Y. and Kim, Y. (2019) Impact of Different Extraction Solvents on Phenolic Content and Antioxidant Potential of *Pinus densiflora* Bark Extract. *BioMed Research International*, **2019**, Article ID: 3520675.  
<https://doi.org/10.1155/2019/3520675>
- [49] Shakun, A.S., Vorobyova, V.I., Chygyrynets, O.E. and Skiba, M.I. (2020) Influence of Solvent on the Component Composition and Antioxidant Properties of Apricot Cake (*Prunus armeniaca* L.) Extracts. *Journal of Chemistry*, **2020**, Article ID: 2913454.  
<https://doi.org/10.1155/2020/2913454>
- [50] Kaczorová, D., Karalija, E., Dahija, S., Bešta-Gajević, R., Parić, A. and Čavar Zeljković, S. (2021) Influence of Extraction Solvent on the Phenolic Profile and Bioactivity of Two Achillea Species. *Molecules*, **26**, Article No. 1601.  
<https://doi.org/10.3390/molecules26061601>
- [51] Banica, F., Bungau, S., Tit, D.M., Behl, T., Otrisal, P., Nechifor, A.C., *et al.* (2020) Determination of the Total Polyphenols Content and Antioxidant Activity of Echinacea Purpurea Extracts Using Newly Manufactured Glassy Carbon Electrodes Modified with Carbon Nanotubes. *Processes*, **8**, Article No. 833.  
<https://doi.org/10.3390/pr8070833>
- [52] Quyen, N.T.C., Quyen, N.T.N., Nhan, L.T.H. and Toan, T.Q. (2020) Antioxidant Activity, Total Phenolics and Flavonoids Contents of *Pandanus amaryllifolius* (Roxb.). *IOP Conference Series: Materials Science and Engineering*, **991**, Article ID: 012019.  
<https://doi.org/10.1088/1757-899x/991/1/012019>
- [53] Khiya, Z., Oualcadi, Y., Gamar, A., Berrekhis, F., Zair, T. and Hilali, F.E. (2021)

- Correlation of Total Polyphenolic Content with Antioxidant Activity of Hydromethanolic Extract and Their Fractions of the *Salvia officinalis* Leaves from Different Regions of Morocco. *Journal of Chemistry*, **2021**, Article ID: 8585313. <https://doi.org/10.1155/2021/8585313>
- [54] Motolinia-Alcántara, E.A., Franco-Vásquez, A.M., Nieto-Camacho, A., Arreguín-Espinosa, R., Rodríguez-Monroy, M., Cruz-Sosa, F., *et al.* (2023) Phenolic Compounds from Wild Plant and *in Vitro* Cultures of *Ageratina pichichensis* and Evaluation of Their Antioxidant Activity. *Plants*, **12**, Article No. 1107. <https://doi.org/10.3390/plants12051107>
- [55] Koraqi, H., Petkoska, A.T., Khalid, W., Sehrish, A., Ambreen, S. and Lorenzo, J.M. (2023) Optimization of the Extraction Conditions of Antioxidant Phenolic Compounds from Strawberry Fruits (*Fragaria X ananassa* Duch.) Using Response Surface Methodology. *Food Analytical Methods*, **16**, 1030-1042. <https://doi.org/10.1007/s12161-023-02469-6>
- [56] Zaitun Hasibuan, P.A. and Mardiana (2018) Antioxidant Activity of N-Hexane, Ethyl Acetate and Ethanol Extract from Lakoocha Leaves (*Artocarpus lacucha* Buch.-Ham.) Using DPPH Method. *Indonesian Journal of Pharmaceutical and Clinical Research*, **1**, 41-47. <https://doi.org/10.32734/ijpcr.v1i2.433>
- [57] Baliyan, S., Mukherjee, R., Priyadarshini, A., Vibhuti, A., Gupta, A., Pandey, R.P., *et al.* (2022) Determination of Antioxidants by DPPH Radical Scavenging Activity and Quantitative Phytochemical Analysis of *Ficus Religiosa*. *Molecules*, **27**, Article No. 1326. <https://doi.org/10.3390/molecules27041326>
- [58] Gulcin, İ. and Alwasel, S.H. (2023) DPPH Radical Scavenging Assay. *Processes*, **11**, Article No. 2248. <https://doi.org/10.3390/pr11082248>
- [59] Ilyasov, I.R., Beloborodov, V.L., Selivanova, I.A. and Terekhov, R.P. (2020) ABTS/PP Decolorization Assay of Antioxidant Capacity Reaction Pathways. *International Journal of Molecular Sciences*, **21**, Article No. 1131. <https://doi.org/10.3390/ijms21031131>
- [60] Mira, L., Silva, M., Rocha, R. and Manso, C.F. (1999) Measurement of Relative Antioxidant Activity of Compounds: A Methodological Note. *Redox Report*, **4**, 69-74. <https://doi.org/10.1179/135100099101534666>
- [61] Chung, Y., Chang, C., Chao, W., Lin, C. and Chou, S. (2002) Antioxidative Activity and Safety of the 50 Ethanolic Extract from Red Bean Fermented by *Bacillus subtilis* IMR-NK1. *Journal of Agricultural and Food Chemistry*, **50**, 2454-2458. <https://doi.org/10.1021/jf011369q>
- [62] Govindan, P. and Muthukrishnan, S. (2013) Evaluation of Total Phenolic Content and Free Radical Scavenging Activity of *Boerhavia erecta*. *Journal of Acute Medicine*, **3**, 103-109. <https://doi.org/10.1016/j.jacme.2013.06.003>
- [63] Liu, Y., Wen, F., Yang, H., Bao, L., Zhao, Z. and Zhong, Z. (2022) The Preparation and Antioxidant Activities of Three Phenyl-Acylchitooligosaccharides. *Heliyon*, **8**, e10624. <https://doi.org/10.1016/j.heliyon.2022.e10624>
- [64] Chaves, N., Santiago, A. and Alías, J.C. (2020) Quantification of the Antioxidant Activity of Plant Extracts: Analysis of Sensitivity and Hierarchization Based on the Method Used. *Antioxidants*, **9**, Article No. 76. <https://doi.org/10.3390/antiox9010076>
- [65] Kalinowska, M., Płońska, A., Trusiak, M., Gołębowska, E. and Gorlewska-Pietluszenko, A. (2022) Comparing the Extraction Methods, Chemical Composition, Phenolic Contents and Antioxidant Activity of Edible Oils from *Cannabis sativa* and *Silybum marianum* Seeds. *Scientific Reports*, **12**, Article No. 20609. <https://doi.org/10.1038/s41598-022-25030-7>

- [66] Maliar, T., Maliarová, M., Blažková, M., Kunštek, M., Uváčková, L., Viskupičová, J., *et al.* (2023) Simultaneously Determined Antioxidant and Pro-Oxidant Activity of Randomly Selected Plant Secondary Metabolites and Plant Extracts. *Molecules*, **28**, Article No. 6890. <https://doi.org/10.3390/molecules28196890>
- [67] Ajiboye, B.O., Ojo, O.A., Oyinloye, B.E., Akuboh, O., Okesola, M.A., Idowu, O. and Kappo, A.P. (2020) *In Vitro* Antioxidant and Inhibitory Activities of Polyphenolic-Rich Extracts of *Syzygium cumini* (Linn) Skeels Leaf on Two Important Enzymes Relevant to Type II Diabetes Mellitus. *Pakistan Journal of Pharmaceutical Sciences*, **33**, 523-529. <https://doi.org/10.36721/PIPS.2020.33.2.REG.523-529.1>.
- [68] Kumar, S., Krishna Chaitanya, R. and Preedy, V.R. (2018) Assessment of Antioxidant Potential of Dietary Components. In: Preedy, V.R. and Watson, R.R., Eds., *HIV/AIDS*, Elsevier, 239-253. <https://doi.org/10.1016/b978-0-12-809853-0.00020-1>
- [69] Yao, Y., Liu, Y., Li, C., Huang, X., Zhang, X., Deng, P., *et al.* (2023) Effects of Rosemary Extract Supplementation in Feed on Growth Performance, Meat Quality, Serum Biochemistry, Antioxidant Capacity, and Immune Function of Meat Ducks. *Poultry Science*, **102**, Article ID: 102357. <https://doi.org/10.1016/j.psj.2022.102357>
- [70] Kandylari, A., Potsaki, P., Bousdouni, P., Kaloteraki, C., Christofilea, M., Almpounioti, K., *et al.* (2023) Development of Dairy Products Fortified with Plant Extracts: Antioxidant and Phenolic Content Characterization. *Antioxidants*, **12**, Article No. 500. <https://doi.org/10.3390/antiox12020500>
- [71] Manzoor, A., Yousuf, B., Pandith, J.A. and Ahmad, S. (2023) Plant-Derived Active Substances Incorporated as Antioxidant, Antibacterial or Antifungal Components in Coatings/Films for Food Packaging Applications. *Food Bioscience*, **53**, Article ID: 102717. <https://doi.org/10.1016/j.fbio.2023.102717>
- [72] Kola, V. and Carvalho, I.S. (2023) Plant Extracts as Additives in Biodegradable Films and Coatings in Active Food Packaging. *Food Bioscience*, **54**, Article ID: 102860. <https://doi.org/10.1016/j.fbio.2023.102860>
- [73] Kulisic, T., Radonic, A., Katalinic, V. and Milos, M. (2004) Use of Different Methods for Testing Antioxidative Activity of Oregano Essential Oil. *Food Chemistry*, **85**, 633-640. <https://doi.org/10.1016/j.foodchem.2003.07.024>
- [74] Ahmadi, F., Kadivar, M. and Shahedi, M. (2007) Antioxidant Activity of *Kelussia odoratissima* Mozaff. in Model and Food Systems. *Food Chemistry*, **105**, 57-64. <https://doi.org/10.1016/j.foodchem.2007.03.056>
- [75] Lelono, R.A.A., Tachibana, S. and Itoh, K. (2009) *In Vitro* Antioxidative Activities and Polyphenol Content of Eugenia Polyantha Wight Grown in Indonesia. *Pakistan Journal of Biological Sciences*, **12**, 1564-1570. <https://doi.org/10.3923/pjbs.2009.1564.1570>
- [76] Laguerre, M., Bayrasy, C., Panya, A., Weiss, J., McClements, D.J., Lecomte, J., *et al.* (2014) What Makes Good Antioxidants in Lipid-Based Systems? The Next Theories Beyond the Polar Paradox. *Critical Reviews in Food Science and Nutrition*, **55**, 183-201. <https://doi.org/10.1080/10408398.2011.650335>
- [77] Safaei-Ghomi, J., Ghadamib, M. and Batooli, H. (2010) Bioactivity of Methanol Extracts of *Eucalyptus sargentii* Maiden Cultivated in Iran. *Digest Journal of Nanomaterials and Biostructures*, **5**, 859-863.
- [78] Shahidi, F. and Zhong, Y. (2011) Revisiting the Polar Paradox Theory: A Critical Overview. *Journal of Agricultural and Food Chemistry*, **59**, 3499-3504. <https://doi.org/10.1021/jf104750m>
- [79] Kiokias, S. and Oreopoulou, V. (2022) Review on the Antioxidant Activity of Phenolics in O/W Emulsions along with the Impact of a Few Important Factors on Their

Interfacial Behaviour. *Colloids and Interfaces*, **6**, Article No. 79.  
<https://doi.org/10.3390/colloids6040079>

- [80] Farooq, S., Abdullah, Zhang, H. and Weiss, J. (2021) A Comprehensive Review on Polarity, Partitioning, and Interactions of Phenolic Antioxidants at Oil-Water Interface of Food Emulsions. *Comprehensive Reviews in Food Science and Food Safety*, **20**, 4250-4277. <https://doi.org/10.1111/1541-4337.12792>
- [81] Munteanu, I.G. and Apetrei, C. (2021) Analytical Methods Used in Determining Antioxidant Activity: A Review. *International Journal of Molecular Sciences*, **22**, Article No. 3380. <https://doi.org/10.3390/ijms22073380>