

Immunomodulatory functions of type I interferons

José M. González-Navajas^{1,2,3}, Jongdae Lee¹, Michael David⁴ and Eyal Raz¹

Abstract | Interferon- α (IFN α) and IFN β , collectively known as type I IFNs, are the major effector cytokines of the host immune response against viral infections. However, the production of type I IFNs is also induced in response to bacterial ligands of innate immune receptors and/or bacterial infections, indicating a broader physiological role for these cytokines in host defence and homeostasis than was originally assumed. The main focus of this Review is the underappreciated immunomodulatory functions of type I IFNs in health and disease. We discuss their function in the regulation of innate and adaptive immune responses, the response to bacterial ligands, inflammasome activation, intestinal homeostasis and inflammatory and autoimmune diseases.

Viral interference

The antagonistic or inhibitory effect induced by one virus or its components on the propagation of another virus.

The inhibitory effect that one viral infection often exerts on the infectivity of a different virus was first observed in 1804 by Edward Jenner, who reported that herpetic infections could prevent the development of vaccinia lesions¹. In retrospect, this observation might be the first documented description of what was later coined as the viral interference phenomenon. This phenomenon was first described in detail for plant viruses in the early 1930s². After that, similar observations were made with bacteriophages³ and with animal viruses⁴. In 1954, Nagano and Kojima reported the inhibition of viral growth in areas of rabbit skin that had been previously inoculated with ultraviolet-inactivated vaccinia virus, and in 1957, during a study of the interference produced by heat-inactivated influenza virus, Isaacs and Lindenmann identified that a secreted factor was responsible for this phenomenon and they termed it interferon (IFN)⁵.

Type I IFNs belong to a family of cytokines that attracted much attention owing to their protective role against viral infection. IFNs are widely expressed cytokines that possess strong antiviral and immunomodulatory properties. The IFN family can be classified into three main types of cytokines — type I, type II and type III IFNs. In humans and mice, the type I IFN family is composed of 16 members, namely 12 IFN α subtypes, IFN β , IFN ϵ , IFN κ and IFN ω ⁶. By contrast, the type II IFN family includes only one cytokine: IFN γ , which also exhibits antiviral activities. The third type of IFNs is the IFN λ family, which includes IFN λ 1 (also known as IL-29), IFN λ 2 (also known as IL-28A) and IFN λ 3 (also known as IL-28B). On the basis of protein sequence and

structure, type III IFNs are markedly different from type I and type II IFNs and are more similar to members of the interleukin-10 (IL-10) family; however, they provoke antiviral responses and induce the activation of IFN-stimulated genes (ISGs)⁷.

Recent evidence has uncovered new roles for this family of cytokines beyond their well-known function in viral interference. This article highlights the function of type I IFNs in modulating immune responses. We map the molecular signalling pathways activated by type I IFNs, and describe the function of these cytokines in the response to bacterial ligands and their role in inflammasome activation. In addition, we discuss the role of type I IFNs in intestinal homeostasis and in inflammatory and autoimmune diseases such as coeliac disease, psoriasis, multiple sclerosis and cancer.

Type I IFN production and signalling

Induction of type I IFNs by bacterial ligands. Type I IFNs can be produced by almost every cell type, including leukocytes, fibroblasts and endothelial cells. The signalling pathways that lead to the induction of type I IFNs differ depending on the stimulus and the responding cell types, but they ultimately lead to the activation of some common signalling molecules, including TNF receptor-associated factor 3 (TRAF3)⁸ and the transcription factors IFN regulatory factor 3 (IRF3) and IRF7. Dimerized IRF3 and IRF7 translocate to the nucleus and, concomitantly with the transcription factor nuclear factor- κ B (NF- κ B), bind to both the *IFNA* and *IFNB* promoters⁹ to initiate the transcription of these IFN genes. Although the most important function of type I IFNs is

¹Division of Rheumatology, Allergy and Immunology, Department of Medicine, University of California San Diego, 9500 Gilman Drive, La Jolla, California 92093-0663, USA.

²Biomedical Research Center in Hepatic and Digestive Diseases (CIBERehd), Instituto de Salud Carlos III, Madrid, Spain.

³Fundación para la Investigación en el Hospital General de Alicante, Alicante, Spain.

⁴Division of Biological Sciences, University of California San Diego, La Jolla, California, USA.

Correspondence to J.M.G.-N. and E.R. e-mails: jog001@ucsd.edu; eraz@ucsd.edu doi:10.1038/nri3133 Published online 6 January 2012

IFN-stimulated genes

(ISGs). These genes contain promoters that are responsive to interferon (IFN) signalling, and they are responsible for the antiviral and immunomodulatory properties of IFNs. Over 400 such genes have been identified by microarray analyses. Some, such as RNA-activated protein kinase, ribonuclease L, MX1 (myxovirus resistance 1) and ISG15 (IFN-stimulated gene of 15 kDa), have well-documented antiviral activities, but the precise biological function of most of these genes is unknown.

Inflammasome

A cytosolic multiprotein complex that activates caspase 1 and regulates the release of IL-1 β and IL-18 in response to exogenous pathogens and endogenous danger signals. This complex minimally consists of a danger-sensing initiator component and the effector component, which is mature caspase 1.

Coeliac disease

An immune-mediated enteropathy triggered by intolerance to dietary ingestion of glutamine- and proline-rich proteins, collectively known as gluten, which is present in wheat, barley, rye and other grains. This disease results in gastrointestinal symptoms such as diarrhoea, nutrient malabsorption and weight loss.

Multiple sclerosis

A chronic inflammatory disease of the central nervous system that causes the progressive destruction of the myelin sheaths around axons in any area of the brain, optic nerve and spinal cord. This results in slower nerve impulses.

typically considered to be in the induction of an antiviral immune response, these cytokines are also induced in response to many bacterial pathogens or their products, mainly through Toll-like receptor (TLR)-dependent pathways^{10,11}.

TLRs are the key sensors of microbial invasion in mammals¹², and they activate an innate defence programme that is crucial for host survival. Each TLR senses a particular subset of microbial signature molecules. Most TLRs that recognize bacterial products are linked to the induction of type I IFNs¹⁰; these TLRs include TLR3, TLR4, TLR7 and TLR9. Signalling through TLR3 and TLR4 induces type I IFN production in a broad range of cell types in a manner dependent on TIR-domain-containing adaptor protein inducing IFN β (TRIF). By contrast, TLR7, TLR8 and TLR9 induce type I IFN production in dendritic cells (DCs) — mainly plasmacytoid DCs (pDCs) — via a pathway dependent on myeloid differentiation primary-response protein 88 (MYD88) (TABLE 1).

However, it became evident that TLR-deficient animals can still produce type I IFNs in response to RNA and DNA ligands^{13,14}. These TLR-independent pathways include the cytoplasmic sensors retinoic-acid-inducible gene I (RIG-I) and melanoma differentiation-associated gene 5 (MDA5). In addition, stimulator of IFN genes (STING) and DNA-dependent activator of IRFs (DAI; also known as DLM1 and ZBP1) have been reported to induce type I IFNs in response to cytosolic DNA^{15,16}. STING is an endoplasmic reticulum-associated protein that has been shown to respond to DNA from various pathogens — including *Listeria monocytogenes* and the DNA virus herpes simplex virus 1 (HSV-1) — in macrophages, DCs and epithelial cells¹⁵. DAI, which was the first cytosolic DNA receptor to be described, recognizes viral, bacterial and mammalian double-stranded DNA and induces type I IFN production through the activation of TANK-binding kinase 1 (TBK1), which

subsequently phosphorylates IRF3 (REF. 16). In addition, DAI stimulates the production of pro-inflammatory cytokines — such as IL-6 and tumour necrosis factor (TNF) — through the activation of the kinase receptor-interacting protein 1 (RIP1), which leads to the phosphorylation of NF- κ B inhibitor- α (I κ B α) and the subsequent activation of NF- κ B¹⁷. However, DAI-deficient cells can still induce type I IFN production in response to foreign DNA¹⁸, suggesting the existence of additional mechanisms that contribute to type I IFN production in response to cytosolic DNA. Indeed, a recent study showed that DNA-dependent RNA polymerase III can use cytosolic DNA as a template to synthesize RNA containing a 5'-triphosphate group, and that this RNA activates the RIG-I-IPS1 (*IFNB*-promoter stimulator 1; also known as MAVS) signalling pathway¹⁹.

Signalling pathways activated by type I IFNs. Type I IFNs signal through a common heterodimeric receptor, known as the IFN α/β receptor (IFNAR), which is expressed by nearly all cell types. This receptor consists of two subunits — IFNAR1 and IFNAR2 (REF. 20) — that are constitutively associated with Janus kinase 1 (JAK1) and non-receptor tyrosine kinase 2 (TYK2)²¹. Activation of JAK1 and TYK2 results in the tyrosine phosphorylation and activation of several signal transducer and activator of transcription (STAT) family members; in most cells these include STAT1, STAT2, STAT3 and STAT5, but in lymphocytes type I IFNs also activate STAT4 and STAT6 (REFS 22,23).

Activation of STAT1 and STAT2 leads to the recruitment of IRF9 and the formation of a STAT1-STAT2-IRF9 complex, which is known as the IFN-stimulated gene factor 3 (ISGF3) complex. This complex then migrates to the nucleus and binds to IFN-stimulated response elements (ISREs) in the promoters of ISGs to initiate gene transcription. Other STAT complexes that do not recruit IRF9, including STAT1 homodimers, bind to IFN γ -activated

Table 1 | **Inducers of type I IFNs and responding cells**

Inducer	Source	Receptor	Localization	Responding cell
ssRNA, dsRNA	Viruses	RIG-I and MDA5	Cytoplasm	Multiple cell types
Cytosolic DNA	Viruses or bacteria	STING, DAI and RNA polymerase III	Cytoplasm	Multiple cell types
dsRNA	Viruses	TLR3-TRIF	Endosomes	Macrophages, cDCs and epithelial cells
LPS	Gram-negative bacteria	TLR4-TRIF	Plasma membrane	Macrophages and cDCs
Viral glycolipids	Viruses	TLR4-TRIF	Plasma membrane	Macrophages and cDCs
ssRNA	Viruses or damaged host cells	TLR7-MYD88	Endosomes	pDCs, cDCs and macrophages
Imiquimod	Synthetic	TLR7-MYD88	Endosomes	pDCs, cDCs and macrophages
ssRNA	Viruses	TLR8-MYD88	Endosomes	cDCs
CpG DNA	Bacteria or viruses	TLR9-MYD88	Endosomes	pDCs, cDCs and macrophages

cDC, conventional DC; DAI, DNA-dependent activator of IRFs; DC, dendritic cell; dsRNA, double-stranded RNA; IFN, interferon; LPS, lipopolysaccharide; MDA5, melanoma differentiation-associated gene 5; MYD88, myeloid differentiation primary-response protein 88; pDC, plasmacytoid DC; RIG-I, retinoic-acid-inducible gene I; ssRNA, single-stranded RNA; STING, stimulator of IFN genes; TLR, Toll-like receptor; TRIF, TIR-domain-containing adaptor protein inducing IFN β .

site (GAS) enhancer elements in the promoters of ISGs^{24,25}. Both type I and type II IFNs can induce the activation of GAS elements through, for example, the formation of STAT1 homodimers. However, in contrast to type I IFNs, IFN γ cannot induce the formation of ISGF3 complexes and therefore is not able to promote the engagement of ISRE sites to activate those genes that have only ISREs in their promoters²¹ (FIG. 1).

In addition, both type I and type II IFNs can induce the recruitment and phosphorylation of STAT3 (REFS 21,26). Following its phosphorylation, STAT3 forms homodimers that translocate to the nucleus, where they bind to STAT3-binding elements (SBEs). STAT1 and STAT3 have a high level of sequence similarity and can heterodimerize and activate similar genes *in vitro*²⁷. However, *in vivo*, the sets of genes that are activated by these two

factors are very different²⁸ and depend on the cell type and the activating cytokine, and in many physiological contexts STAT1 and STAT3 exert opposing effects. For example, in most cell types, STAT1 activates several pro-apoptotic and anti-proliferative genes^{29,30}, whereas STAT3 inhibits apoptosis and promotes proliferation through the induction of anti-apoptotic genes of the B cell lymphoma (BCL) family and of oncogenes such as *MYC*^{31,32}.

STAT1 and STAT3 also have opposite roles in inflammation. IFN γ -dependent STAT1 activation usually mediates a pro-inflammatory response that favours the recruitment of immune cells to the site of inflammation³³, and this increases the production of pro-inflammatory mediators³⁴ and enhances antigen processing and presentation by MHC class I and II molecules^{35,36}. By contrast, STAT3 is a key mediator of IL-10 signalling, which




Figure 1 | Signalling pathways activated by type I and type II IFNs. Different signal transducer and activator of transcription (STAT) family members can be activated by interferons (IFNs). STAT1 homodimers can be formed in response to both type I IFNs and type II IFN (IFN γ). These homodimers bind to IFN γ -activated site (GAS) enhancer elements in the promoters of IFN-stimulated genes, and this results in the induction of genes encoding pro-inflammatory cytokines and apoptotic factors. Type I and type II IFNs can also activate STAT3 homodimers, and this can result in the production of both pro-inflammatory cytokines and anti-inflammatory cytokines (such as interleukin-10 (IL-10)), although the underlying mechanisms are not known. STAT1–STAT2 heterodimers, which are activated by type I IFNs, bind to IFN regulatory factor 9 (IRF9) in the cytosol to form the IFN-stimulated gene factor 3 (ISGF3) complex, which in turn migrates to the nucleus to bind to IFN-stimulated response elements (ISREs) and activate antiviral and antibacterial genes. In addition, type I IFNs stimulate IL-10 production either through the phosphoinositide 3-kinase (PI3K)–AKT pathway or through STAT3 homodimers. Finally, in a STAT-independent manner, type I IFNs activate both p38, which is an upstream activator of several genes regulated by ISREs and GAS elements, and mammalian target of rapamycin (mTOR), which regulates mRNA translation. CREB, cAMP-responsive-element-binding protein; IFNAR, IFN α/β receptor; IFNGR, IFN γ receptor; JAK, Janus kinase; SBE, STAT3-binding element; *TBX21*, T box 21; TYK2, non-receptor tyrosine kinase 2.

negatively regulates pro-inflammatory responses by activated macrophages and DCs and can directly inhibit STAT1 activation³⁷. Interestingly, other receptors, such as the IL-6 receptor, can activate STAT3, but this does not result in an anti-inflammatory response, possibly owing to the participation of additional regulatory factors, such as members of the suppressor of cytokine signalling (SOCS) family^{38,39}. This dual ability of STAT3 to differentially regulate inflammation in different scenarios might explain some of the controversial roles that type I IFNs have in different human conditions (discussed below). Several studies also suggest that cross-regulation between STAT1 and STAT3 and their relative abundance in a cell are the defining factors that determine the biological effects of their upstream activators, such as IL-6 and IFNs^{40,41}.

Type I IFNs were also shown to promote the production of IL-10 by lipopolysaccharide (LPS)-stimulated macrophages⁴² and by LPS-stimulated human peripheral blood mononuclear cells (PBMCs)⁴³. This effect was initially attributed to the activation of STAT3 and IRF1 in IFN α -stimulated human monocytes⁴⁴. However, another study indicated that IFN β stimulates IL-10 production independently of STAT3, by activating a signalling pathway mediated by JAK1 and phosphoinositide 3-kinase (PI3K). This pathway increases the phosphorylation and nuclear translocation of cAMP-responsive element-binding protein (CREB), which promotes IL-10 production in human DCs⁴⁵ (FIG. 1). It is not clear why IL-10 production is mediated by different signalling pathways in response to IFN α (the STAT3 pathway) and IFN β (the PI3K pathway). However, the differential activation of STAT3 versus STAT1, the ability to induce IL-10 production and/or the participation of additional regulatory elements may explain the diverse effects of type I IFNs in several pathological conditions.

In addition to the JAK–STAT signalling pathways, there is evidence that type I IFNs activate other (non-STAT) signalling pathways that have crucial roles in their different biological properties. For example, it has been shown that type I IFNs activate signalling pathways mediated by mitogen-activated protein kinases (MAPKs), specifically p38 and extracellular signal-regulated kinase 1 (ERK1) or ERK2. p38 activity is required for type I IFN-dependent transcription of several genes that are regulated by ISREs and GAS elements in a STAT-independent manner^{46–48}. The functional relevance of this signalling pathway in the biological effects of type I IFNs was evidenced by several studies indicating that p38 is required for the growth-inhibitory and antiviral effects of type I IFNs^{49,50}. In addition to the induction of the p38 signalling cascade, the MAPK/ERK kinase (MEK)–ERK pathway is activated by type I IFNs⁵¹ and participates in the response to viral infection⁵². Finally, as mentioned above, type I IFNs can also induce the activation of the PI3K signalling pathway in a STAT-independent manner^{53,54}. The PI3K pathway has an important role in mediating gene transcription in response to both type I and type II IFNs and is essential for mediating their antiviral effects against the encephalomyocarditis virus *in vitro*⁵⁵. Activation of the

PI3K signalling cascade controls the activation of mammalian target of rapamycin (mTOR), which regulates mRNA translation. Notably, it was shown that type I IFNs activate pathways for the initiation of mRNA translation that are downstream of mTOR, such as the pathway that involves the activation of p70 S6 kinase and the subsequent phosphorylation of the S6 ribosomal protein^{56,57}. The activation of mTOR by type I IFNs was unrelated to the activation of STAT family members and had no effect on gene transcription^{56,57}, indicating that mTOR selectively regulates IFN-induced mRNA translation. In terms of biological relevance, the activation of mTOR signalling has been shown to mediate the antiviral effects of IFN α against the hepatitis C virus⁵⁸.

Modulating immune responses by type I IFNs

Type I IFNs and suppression of intestinal inflammation.

The intestinal microbiota influences the development of local and systemic immune responses and the proliferation and barrier functions of the intestinal epithelium, in part via the activation of host TLRs⁵⁹. Over the last few years, data from our laboratory and others have shown that the systemic administration of TLR9 ligands reduces the severity of colonic injury and inflammation in models of experimental colitis^{60–62}, in part through the TLR9-induced production of type I IFNs by DCs⁶⁰. In addition, IFN α enhances the barrier function of intestinal epithelial cells *in vitro* by activating STAT3, which maintains the expression of several key tight junction molecules, such as members of the claudin family (J.L., unpublished observations). Similarly, activation of TLR3 by the synthetic viral RNA polyinosinic–polycytidylic acid (polyI:C), which induces type I IFN production in many cell types, was shown to protect mice from experimental colitis⁶². Furthermore, *Ifnar*^{-/-} mice were extremely susceptible to dextran-sulphate sodium (DSS)-induced colitis^{60,63}, and the administration of recombinant IFN β to DSS-treated wild-type mice mimicked the anti-inflammatory effects of TLR9 ligands⁶⁰.

Different types of haematopoietic cells contribute to DSS-induced colonic inflammation. The depletion of CD11c^{hi} DCs after DSS administration showed that type I IFNs induced by TLR9-activated CD11c^{hi} DCs attenuate the severity of colitis in this model⁶⁴. This study also showed that, depending on their mode of activation, DCs can enhance or inhibit acute DSS-induced colitis by secreting various cytokines (such as TNF, IL-6, IL-10 and type I IFNs), by regulating the production of chemokines that affect the composition of the cellular infiltrate, and by affecting the rate of resolution of inflammation in the colon⁶⁴. In addition, a recent study using a genetically modified strain of *Lactobacillus acidophilus* that constitutively expresses IFN β reported that the local expression of IFNAR1 by CD103⁺ DCs in the Peyer's patches is necessary for protection against DSS-induced intestinal inflammation⁶³. In this study, saturating expression of IFN β before the induction of colitis resulted in the transient suppression of IFNAR1 expression on intestinal CD103⁺ DCs; this prevented IFN β from signalling during the inflammatory period, mimicking the phenotype of *Ifnar*^{-/-} mice with DSS-induced colitis⁶³.

Peyer's patches

Collections of lymphoid tissue that are located in the mucosa of the small intestine, with an outer epithelial layer that consists of specialized epithelial cells called M cells.

Regulation of adaptive immune responses by type I IFNs. In addition to the T cell-independent protective effects of TLR9-mediated type I IFN induction, the role of TLR9 agonist-induced protection in T cell-dependent intestinal inflammation has also been evaluated. TLR9 signalling in DCs in the lamina propria of the small intestine was shown to regulate the functions of regulatory T (T_{Reg}) cells and effector T cells⁶⁵. In this study, T_{Reg} cell induction was inhibited by DNA derived from commensal bacteria through TLR9, and this was shown to favour the development of protective mucosal immune responses; however the role of type I IFNs in these effects was not clarified⁶⁵. In a T cell-transfer model of colitis, treatment with a TLR9 ligand in the absence of type I IFN signalling failed to induce functionally suppressive $CD4^+CD62L^+$ T cells, whereas pretreatment of the donor mice with recombinant IFN β and subsequent T cell transfer resulted in reduced intestinal inflammation and decreased secretion of pro-inflammatory cytokines⁶⁶. Indeed, a role for type I IFNs in TLR9-induced protection in T cell-dependent colitis was previously shown in germ-free mice; in this study, type I IFNs were shown to induce the expression of regulatory markers on $CD4^+CD62L^+$ T cells⁶⁷. These data correlate with previous reports that type I IFNs induce regulatory capacities in T cells^{68,69}. Therefore, type I IFNs seem to exert a protective effect in the intestinal mucosa in both acute and chronic models of colitis.

Type I IFNs have been shown to have an important role in the differentiation of both $CD4^+$ and $CD8^+$ T cells. Initial studies found that IL-12 signalling in $CD4^+$ T cells and the subsequent activation of STAT4, which in turn promotes T-bet expression, are crucial events in T helper 1 (T_H1) cell differentiation⁷⁰. In addition to IL-12, type I IFNs were shown to induce the tyrosine phosphorylation and DNA binding of STAT4 (REF. 71) and to act directly on human T cells, but not mouse T cells, to drive T_H1 cell development⁷². However, it was later reported that type I IFNs activate STAT4 directly and that this activation is required for IFN γ production during viral infections in mice^{73,74}. Thus, phosphorylation of STAT4 has been detected in response to type I IFN treatment of both human and mouse T cells^{74–76} but, in contrast with IL-12, type I IFNs were unable to sustain STAT4 phosphorylation and therefore were not sufficient to drive T_H1 cell commitment *in vitro*^{75,77}. However, type I IFNs regulate T_H1 cell differentiation and effector functions *in vivo* by synergizing with other cytokines, such as IL-18 and IL-21 (REFS 78–80). Moreover, type I IFNs have been associated with the suppression of T_H2 and T_H17 cell-mediated responses. In human cells, type I IFNs reversed T_H2 cell commitment by suppressing the expression of the T_H2 cell-associated transcription factor GATA-binding protein 3 (GATA3)⁷⁸. Similarly, type I IFNs also suppress the differentiation of T_H17 cells in both mice⁸¹ and humans⁸².

Forkhead box P3 (FOXP3)⁺ T_{Reg} cells have a pivotal role in maintaining immunological tolerance and homeostasis. IFN β treatment markedly improved the frequency and suppressive function of T_{Reg} cells in patients with relapsing–remitting multiple sclerosis^{9,83} and in

patients with chronic hepatitis C virus infection⁸⁴, and also increased *Foxp3* mRNA expression in PBMCs from patients with relapsing–remitting multiple sclerosis⁸⁵. FOXP3⁺ T_{Reg} cells show functional and phenotypical plasticity in response to environmental cues and are capable of secreting pro-inflammatory cytokines, for example during severe inflammation^{86,87}. Recently, a higher frequency of IFN γ -secreting T_{Reg} cells has been reported in untreated patients with relapsing–remitting multiple sclerosis compared with the frequency in control individuals⁸⁸. By contrast, in patients treated with IFN β , the frequency of IFN γ^+ FOXP3⁺ T cells was similar to that in healthy controls⁸⁸. In addition, ongoing studies in our laboratory indicate that type I IFN signalling is essential for the maintenance of FOXP3 expression *in vivo* and for the suppressive activity of T_{Reg} cells in a model of T cell-mediated colitis in mice (S. E. Lee, J.M.G.-N. and E.R., unpublished observations). These data suggest a novel role for type I IFNs in T_H cell differentiation, as well as in the suppressive function of T_{Reg} cells.

Type I IFNs and inflammasome activation. Four different initiator components that activate different types of inflammasome in response to different stimuli have been identified: NOD-, LRR- and pyrin domain-containing 1 (NLRP1), NLRP3, NOD-, LRR- and CARD-containing 4 (NLRC4; also known as IPAF) and absent in melanoma 2 (AIM2)⁸⁹.

Recently, an inhibitory role for type I IFNs in the activation of the inflammasome has been reported⁹⁰. In this study, type I IFNs inhibited IL-1 β production through two different mechanisms. First, IFN β signalling directly inhibited the NLRP1 and NLRP3 inflammasomes (but not the NLRC4 and AIM2 inflammasomes) in a STAT1-dependent manner. Second, type I IFNs induced the production of IL-10, which in turn activated the transcription factor STAT3 in an autocrine manner to reduce the levels of pro-IL-1 α and pro-IL-1 β ⁹⁰ (FIG. 2a). Furthermore, type I IFNs increased the susceptibility of mice to *Candida albicans* infection, an effect that was attributed to the reduction in IL-1 β production⁹⁰. Therefore, these data suggested an inhibitory role of type I IFNs in inflammasome activation.

By contrast, previous work reported a positive effect of type I IFNs in inflammasome activation during *Francisella tularensis* infection⁹¹. The cytosolic pathogens *F. tularensis* and *L. monocytogenes* were both shown to induce a type I IFN response that was essential for caspase 1 activation and IL-1 β production. Consistent with this, *F. tularensis* DNA released into the cytosol activated the production of type I IFNs via IRF3, and this was necessary for the activation of the DNA-sensing AIM2 inflammasome⁹² (FIG. 2b). The discrepancies regarding the role of type I IFN signalling in the production of IL-1 β might be related to the type of inflammasome; that is, type I IFNs mediate the inhibition of the NLRP3 inflammasome but the activation of the AIM2 inflammasome. Of particular interest, however, is the case of *L. monocytogenes*, which has been shown to activate the NLRP3 inflammasome⁸⁹ and to increase IL-1 β production in an IFN β -dependent manner.




Figure 2 | Type I IFNs regulate inflammasome activation. a | Type I interferons (IFNs) inhibit the production of IL-1 β by the inflammasome through two different mechanisms. First, they activate signal transducer and activator of transcription 1 (STAT1), which directly inhibits the NOD-, LRR- and pyrin domain-containing 1 (NLRP1) and NLRP3 inflammasomes but not the absent in melanoma 2 (AIM2) or NOD-, LRR- and CARD-containing 4 (NLRC4) inflammasomes. Second, type I IFNs induce the production of interleukin-10 (IL-10), which binds to the IL-10 receptor (IL-10R) in an autocrine manner and activates STAT3, which in turn reduces the levels of the precursors pro-IL-1 α and pro-IL-1 β . **b** | Type I IFN signalling is required for efficient activation of the AIM2 inflammasome in response to *Francisella tularensis*. After the bacterium enters the phagosome, the phagosome is rapidly acidified. Acidification causes the release of bacterial DNA into the cytosol, and this DNA activates an unidentified DNA sensor, which in turn activates IFN regulatory factor 3 (IRF3) to initiate the production of type I IFNs. IFN β then binds to the IFN α/β receptor (IFNAR) in an autocrine manner to enhance the activation of the AIM2 inflammasome, possibly by increasing phagosomal acidification and/or bactericidal activity, thereby favouring the release of more bacterial DNA. CREB, cAMP-responsive-element-binding protein; TLR, Toll-like receptor.

The cytokines IL-1 α and IL-1 β both bind to IL-1 receptor type 1 (IL-1R1), leading to the activation of NF- κ B, MAPKs and certain IRFs^{93,94}. In a recent study, we showed that IL-1R1 is necessary for the TLR9-dependent activation of a type I IFN and IL-10 response⁹⁵. The mechanism by which IL-1R1 signalling modulates type I IFN production involves changes in the ubiquitylation profile of TRAF3, an E3 ubiquitin ligase that interacts with both MYD88 and TRIF, and the type of ubiquitylation determines whether type I IFNs or pro-inflammatory cytokines are produced^{8,96}. Lysine 48 (K48)-linked polyubiquitylation of TRAF3 leads to its proteosomal degradation and the activation of MAPKs and pro-inflammatory cytokines, whereas K63-linked polyubiquitylation of TRAF3 results in the activation of IRFs and subsequent type I IFN production^{96,97}. K63-linked, but not K48-linked, polyubiquitylation of TRAF3 is greatly reduced in the absence of IL-1 signalling. This effect is mediated by deubiquitylating enzyme A (DUBA; also known as OTUD5)⁹⁵, which specifically cleaves the K63-linked polyubiquitin chain on TRAF3 (REF. 97). Together, these data suggested a model in which IL-1R1 positively regulates TLR-induced type I IFN production (FIG. 3). This mechanism may explain previous observations of IL-1R1-mediated protection against intestinal damage from *Citrobacter rodentium* infection⁹⁸ and DSS-induced colitis^{98,99}.

Type I IFNs and bacterial infections. Although type I IFNs are induced by bacterial pathogens, the role of type I IFNs in the context of bacterial infections is not completely understood. Bacterial induction of type I IFNs can be mediated through the TLR-dependent recognition of bacterial products, such as LPS, or through the TLR-independent recognition of bacterial ligands that are delivered to the host cytosol. Many investigators have reported a variety of beneficial but also detrimental immune functions for type I IFNs during bacterial infection. For example, type I IFNs have an important role in mediating the pathology of LPS-induced septic shock⁸⁴. By contrast, type I IFNs impair the clearance of *L. monocytogenes*^{100,101}, *Mycobacterium tuberculosis*¹⁰² and *Chlamydia muridarum*^{83,103}, and they are detrimental to host survival after infection with *F. tularensis*¹⁰⁴.

Two mechanisms have been proposed to explain these phenomena. The first mechanism suggests that microorganisms induce type I IFN production as a strategy to induce apoptosis in lymphocytes, resulting in the suppression of adaptive immune responses¹⁰⁵. The second mechanism proposes that type I IFNs suppress the production of IL-17A and IL-17F, which are necessary for neutrophil-mediated bacterial clearance¹⁰⁴. By contrast, type I IFNs are necessary for host resistance against

other bacterial pathogens. Mice deficient in IFNAR showed decreased survival and increased bacterial burdens after infection with *Streptococcus pneumoniae* or *Escherichia coli*¹⁰⁶, and this was attributed to the reduced expression of certain cytokines, such as TNF or IFN γ . Moreover, type I IFNs were shown to play an important part in restricting the growth of *Legionella pneumophila* in macrophages¹⁰⁷. Taken together, these data show that type I IFNs have a wide range of immunomodulatory effects in response to bacterial infections, and this clearly expands the old notion that type I IFNs serve only as antiviral cytokines.

Type I IFNs in autoimmune disease

The connection between type I IFNs and several autoimmune and inflammatory disorders is well known, although there is considerable variation in the precise mechanisms and in the role of these cytokines in each condition. Some autoimmune diseases (such as psoriasis and systemic lupus erythematosus (SLE)) are improved by the inhibition of type I IFNs or their upstream regulators. By contrast, other conditions that are characterized by strong T_H1 and/or T_H17 cell responses — such as arthritis, inflammatory bowel disease (IBD) and multiple sclerosis — benefit from the administration of type I IFNs (FIG. 4). All these inflammatory conditions have an important impact on public health and are the focus of numerous analyses that are readily available in the literature. Therefore, they are only briefly discussed here.

Inflammatory bowel disease. IBD encompasses two major clinical entities: Crohn’s disease and ulcerative colitis. Conventional therapies for IBD include 5-aminosalicylic acid (5-ASA), antibiotics (such as ciprofloxacin and metronidazole), corticosteroids, immunosuppressants and TNF-specific antibodies (such as infliximab and adalimumab)¹⁰⁸. However, poor efficacy in some cases (with 5-ASA or antibiotic treatment) or the development of serious side effects (with corticosteroid treatment) led to the search for other biological therapies. Given their mechanisms of action and their beneficial effects on intestinal homeostasis in animal models, type I IFNs have been tested as a treatment for IBD. Although a few clinical studies of IFN α or IFN β therapy in patients with ulcerative colitis initially showed promising results^{109–111}, most of these studies failed to demonstrate a beneficial therapeutic effect^{112–115}. Furthermore, documented cases of the exacerbation of ulcerative colitis during IFN α therapy for chronic hepatitis C infection^{116,117} and of the development of ulcerative colitis in patients with multiple sclerosis following treatment with IFN β 1 α ¹¹⁸ call into question whether type I IFNs have a therapeutic role in IBD.

Coeliac disease. Coeliac disease has a strong genetic component, as it is highly associated with HLA-DQ2 and HLA-DQ8 alleles¹¹⁹. The pathogenesis of coeliac disease involves the polarization of T cells into T_H1 cells and the production of high levels of IFN γ in response to gluten¹²⁰. T_H1 cell polarization is driven by many




Figure 3 | IL-1R1 signalling regulates type I IFN production. Interleukin-1 receptor 1 (IL-1R1) signalling positively modulates the production of type I interferons (IFNs) through the differential ubiquitylation of TNF receptor-associated factor 3 (TRAF3). Lysine 48 (K48)-linked polyubiquitylation of TRAF3 leads to its proteosomal degradation and the production of pro-inflammatory cytokines. By contrast, K63-linked polyubiquitylation of TRAF3 triggers the activation of IFN regulatory factor 3 (IRF3) and the subsequent production of type I IFNs. The absence of IL-1 signalling results in increased levels of deubiquitylating enzyme A (DUBA), which cleaves the K63-linked but not the K48-linked ubiquitin chains on TRAF3. LPS, lipopolysaccharide; MAPK, mitogen-activated protein kinase; NF- κ B, nuclear factor- κ B; TLR, Toll-like receptor.

cytokines, including IL-21, IL-18 and IFN α ¹²¹, the levels of which are usually elevated in the mucosa of untreated patients with coeliac disease¹²². Interestingly, the addition of IFN α -specific, but not IL-18-specific, blocking antibodies to biopsy specimens from patients with coeliac disease inhibits IFN γ production in *ex vivo* organ cultures¹²³. Activation of gluten-specific CD4⁺ T cells requires that gluten antigens be presented by antigen-presenting cells (APCs) that express HLA-DQ2 or HLA-DQ8 (REFS 124,125). The expression of these alleles is relatively low in normal mucosa, but can be upregulated on APCs by type I and type II IFNs, and this facilitates the activation of other inflammatory cells, thereby favouring an inflammatory response to gluten peptides^{122,126,127}. Notably, enteric viruses can induce the production of type I IFNs, leading to the upregulation of HLA-DQ2 or HLA-DQ8 on APCs, and this would result in the perfect milieu for the activation of gluten-specific T_H1 cells. The development of coeliac disease in patients undergoing treatment with recombinant type I IFNs, such as patients with hepatitis C virus infection^{122,128}, also supports this hypothesis.

Systemic lupus erythematosus (SLE). An autoimmune disease characterized by the presence of circulating immune complexes that contain antinuclear antibodies bound to self nucleic acids and other nuclear antigens.

Inflammatory bowel disease (IBD). A chronic inflammatory condition that affects the intestinal tract. The proposed pathogenesis of IBD involves a complex model that includes abnormalities of innate immune function and their relationship with the commensal microbiota, inappropriate release of pro-inflammatory cytokines and other mediators, alterations of the intestinal epithelial barrier, and a cytokine imbalance that promotes the pro-inflammatory activity of adaptive immune cells.




Figure 4 | Type I IFNs in human diseases. Type I interferons (IFNs) are implicated in different human diseases, although their role in each condition varies. Type I IFNs usually have a beneficial impact in inflammatory syndromes, such as inflammatory bowel disease (IBD) and multiple sclerosis. By contrast, some autoimmune diseases, such as psoriasis and systemic lupus erythematosus (SLE), are improved by type I IFN inhibition. This figure summarizes the positive (green) and negative (red) roles of type I IFNs in different human conditions. IL-10, interleukin-10; T_H, T helper.

Psoriasis. Psoriasis is a T cell-mediated chronic inflammatory skin disease that is characterized by the production of large amounts of type I IFNs by pDCs that are recruited to and accumulate in the dermis. High levels of type I IFNs are responsible for the local activation and expansion of pathogenic T cell populations, which in turn trigger the abnormal proliferation and differentiation of keratinocytes and the development of diffuse epidermal hyperplasia, known as acanthosis. The activation of T_H17 cells has been associated with this phenomenon¹²⁹, and the blockade of type I IFN signaling by IFNAR-specific antibodies effectively inhibited the activation of autoreactive T cells and the development of skin lesions in a xenograft model of psoriasis¹³⁰. In addition, antibodies specific for blood DC antigen 2 (BDCA2; also known as CLEC4C) suppressed the production of type I IFNs by pDCs through the activation of SRC family protein tyrosine kinases, and this also inhibited the activation of autoreactive T cells and the development of skin lesions¹³¹.

Systemic lupus erythematosus. The activation of pDCs by nucleic acid-containing immune complexes leads to the production of large amounts of type I IFNs, and elevated levels of IFN α are detected in the blood of patients with SLE^{132,133}. Type I IFNs in the serum of patients with SLE were responsible for the differentiation of monocytes into DCs and the expression of MHC class II molecules and the co-stimulatory molecules CD80 and CD86 by these DCs. In addition, normal monocytes cultured with serum from patients with SLE, but not those cultured with autologous serum, could present self antigens to autoreactive T cells¹³⁴. In another study, type I IFNs produced by pDCs, together with IL-6, were also shown to induce plasma cell differentiation¹³⁵.

Patients with SLE are treated with various medications, including non-steroidal anti-inflammatory drugs, glucocorticoids and immunosuppressants. Many of these therapies are associated with severe adverse effects. Therefore, there is a medical need for more-specific and safer therapies that target selective pathways involved in the pathogenesis of SLE. Recent data from a Phase I clinical trial showed that the administration of neutralizing antibodies specific for IFN α effectively inhibited the overexpression of IFN α -inducible genes in skin lesions of patients with SLE¹³⁶.

Multiple sclerosis. It is generally considered that both T_H1 and T_H17 cells are involved in the pathogenesis of multiple sclerosis and experimental autoimmune encephalomyelitis (EAE; an animal model of multiple sclerosis). Human recombinant IFN β therapy is widely prescribed for certain stages of multiple sclerosis, particularly relapsing–remitting multiple sclerosis¹³⁷. Two human recombinant IFN β forms are currently used: IFN β 1a, which is produced in Chinese hamster ovary cells; and IFN β 1b, which is produced in *E. coli*¹³⁸. The exact mechanism behind the beneficial effect of type I IFNs in multiple sclerosis and EAE is not known, but recombinant IFN β reduces the attack frequency and severity of multiple sclerosis^{138,139}. However, not all patients with relapsing–remitting multiple sclerosis respond to treatment, and the development of IFN β -specific autoantibodies may cause relapses and side effects.

IFN β -deficient mice are more susceptible to EAE development¹⁴⁰, and type I IFNs can attenuate EAE in mice¹⁴¹, suggesting that IFN β modulates both multiple sclerosis and EAE. In addition, the conditional deletion of *Ifnar1* showed that specific myeloid cells, such as monocytes and macrophages¹⁴² or DCs¹⁴³, modulate autoimmune inflammation of the central nervous system in an IFNAR-dependent manner. The expression of IFNAR by microglial cells (brain-endogenous macrophages) also modulates the severity of EAE¹⁴². IFN β reduces the ability of microglial cells to present antigens, and this in turn reduces the recruitment and effector functions of encephalitogenic T cells¹⁴⁴. Another mechanism that was proposed to explain the immunomodulatory effects of IFN β in patients with multiple sclerosis involves the induction of IL-10. Indeed, upregulation of IL-10 production is a hallmark of IFN β treatment in patients with multiple sclerosis¹⁴⁵.

Cancer. Type I IFNs have been extensively used for the treatment of several types of cancer, including haematological malignancies (for example, hairy cell leukaemia and some B or T cell lymphomas) and solid tumours (for example, melanoma, renal cell carcinoma and Kaposi's sarcoma)^{146,147}.

IFN α 1-producing tumour cells were shown to be less tumorigenic and less able to metastasize when transferred into mice, and this was attributed to the activation of several host antitumour mechanisms at the tumour site¹⁴⁸. These studies suggested that type I IFNs could act as an adjuvant in cancer vaccines¹³³.

Indeed, IFN α showed an effective adjuvant activity in patients with melanoma¹⁴⁹. In addition to their putative role as adjuvants in antitumour vaccines, type I IFNs inhibit angiogenesis and act on DCs to enhance the ability of these cells to cross-present apoptotic antigens to CD8⁺ T cells¹⁵⁰ to trigger a cytotoxic T lymphocyte response^{151,152}.

The main pitfall for the use of type I IFNs as an anti-tumour therapy is the frequent severe side effects, which decrease the enthusiasm for the application of type I IFNs in this clinical setting.

Concluding remarks

Our understanding of the functions of type I IFNs and the mechanisms that control these functions is continuously evolving. Since their discovery, type I IFNs

have been associated with host defence responses to viral infections, and it is only recently that their functions in bacterial infections and in immune-mediated and inflammatory disorders have been appreciated. The molecular mechanisms by which type I IFNs exert their immunomodulatory functions and the reasons why they restrain the development of some immunopathologies while increasing the severity of others are still largely unknown. One possible explanation for these opposing effects could be found in the differential regulation of STAT family members in different tissues or organs under different physiological and/or inflammatory conditions. A better understanding of the underlying pathophysiology is mandatory for the design of effective treatments with type I IFNs or with type I IFN-specific neutralizing antibodies.

- Jenner, E. Letter to the Editor. *Med. Phys. J.* **12**, 97–102 (1804).
- Salaman, R. N. Protective inoculation against a plant virus. *Nature* **131**, 468 (1933).
- White, P. B. Lysogenic strains of *V. cholerae* and the influence of lysozyme on cholera phage activity. *J. Pathol. Bacteriol.* **44**, 276–278 (1937).
- Hoskins, M. A protective action of neurotropic against viscerotropic yellow fever virus in *Macacus rhesus*. *Am. J. Trop. Med.* **15**, 675–680 (1935).
- Isaacs, A. & Lindenmann, J. Virus interference. I. The interferon. *Proc. R. Soc. Lond. B Biol. Sci.* **147**, 258–267 (1957).
- Pestka, S., Krause, C. D. & Walter, M. R. Interferons, interferon-like cytokines, and their receptors. *Immunol. Rev.* **202**, 8–32 (2004).
- Gad, H. H. *et al.* Interferon- λ is functionally an interferon but structurally related to the interleukin-10 family. *J. Biol. Chem.* **284**, 20869–20875 (2009).
- Hacker, H. *et al.* Specificity in Toll-like receptor signalling through distinct effector functions of TRAF3 and TRAF6. *Nature* **439**, 204–207 (2006). **This study showed that TRAF3 is essential for both type I IFN and IL-10 production, but dispensable for the production of pro-inflammatory cytokines.**
- Schafer, S. L., Lin, R., Moore, P. A., Hiscott, J. & Pitha, P. M. Regulation of type I interferon gene expression by interferon regulatory factor-3. *J. Biol. Chem.* **273**, 2714–2720 (1998).
- Monroe, K. M., McWhirter, S. M. & Vance, R. E. Induction of type I interferons by bacteria. *Cell. Microbiol.* **12**, 881–890 (2010).
- Trinchieri, G. Type I interferon: friend or foe? *J. Exp. Med.* **207**, 2053–2063 (2010).
- Akira, S. Toll-like receptors and innate immunity. *Adv. Immunol.* **78**, 1–56 (2001).
- Barber, G. N. Innate immune DNA sensing pathways: STING, AIMII and the regulation of interferon production and inflammatory responses. *Curr. Opin. Immunol.* **23**, 10–20 (2011).
- Kawai, T. & Akira, S. The roles of TLRs, RLRs and NLRs in pathogen recognition. *Int. Immunol.* **21**, 317–337 (2009).
- Ishikawa, H., Ma, Z. & Barber, G. N. STING regulates intracellular DNA-mediated, type I interferon-dependent innate immunity. *Nature* **461**, 788–792 (2009). **A description of STING as an important mediator in the host defence response against DNA-containing pathogens and in the adjuvant activity of DNA-based vaccines.**
- Takaoka, A. *et al.* DAI (DLM-1/ZBP1) is a cytosolic DNA sensor and an activator of innate immune response. *Nature* **448**, 501–505 (2007). **The first identification of DAI as a new cytosolic DNA sensor that enhances the induction of type I IFN genes and other genes involved in innate immunity.**
- Kaiser, W. J., Upton, J. W. & Mocarski, E. S. Receptor-interacting protein homotypic interaction motif-dependent control of NF- κ B activation via the DNA-dependent activator of IFN regulatory factors. *J. Immunol.* **181**, 6427–6434 (2008).
- Ishii, K. J. *et al.* TANK-binding kinase-1 delineates innate and adaptive immune responses to DNA vaccines. *Nature* **451**, 725–729 (2008).
- Chiu, Y. H., Macmillan, J. B. & Chen, Z. J. RNA polymerase III detects cytosolic DNA and induces type I interferons through the RIG-I pathway. *Cell* **138**, 576–591 (2009).
- Honda, K. & Taniguchi, T. IRFs: master regulators of signalling by Toll-like receptors and cytosolic pattern-recognition receptors. *Nature Rev. Immunol.* **6**, 644–658 (2006).
- Platanias, L. C. Mechanisms of type-I- and type-II-interferon-mediated signalling. *Nature Rev. Immunol.* **5**, 375–386 (2005). **An excellent review of the signalling mechanisms of type I IFNs, including the classical JAK-STAT pathways, as well as the MAPK and mTOR pathways.**
- Fasler-Kan, E., Pansky, A., Wiederkehr, M., Battegay, M. & Heim, M. H. Interferon- α activates signal transducers and activators of transcription 5 and 6 in Daudi cells. *Eur. J. Biochem.* **254**, 514–519 (1998).
- Matikainen, S. *et al.* Interferon- α activates multiple STAT proteins and upregulates proliferation-associated *IL-2R α* , *c-myc*, and *pim-1* genes in human T cells. *Blood* **93**, 1980–1991 (1999).
- Aaronson, D. S. & Horvath, C. M. A road map for those who don't know JAK-STAT. *Science* **296**, 1653–1655 (2002).
- Platanias, L. C. & Fish, E. N. Signaling pathways activated by interferons. *Exp. Hematol.* **27**, 1583–1592 (1999).
- Sato, T., Selli, C., Young, N. S. & Maciejewski, J. P. Inhibition of interferon regulatory factor-1 expression results in predominance of cell growth stimulatory effects of interferon- γ due to phosphorylation of Stat1 and Stat3. *Blood* **90**, 4749–4758 (1997).
- Heinrich, P. C., Behrmann, I., Muller-Newen, G., Schaper, F. & Graeve, L. Interleukin-6-type cytokine signalling through the gp130/Jak/STAT pathway. *Biochem. J.* **334**, 297–314 (1998).
- Baker, S. J., Rane, S. G. & Reddy, E. P. Hematopoietic cytokine receptor signaling. *Oncogene* **26**, 6724–6737 (2007).
- Regis, G., Pensa, S., Boselli, D., Novelli, F. & Poli, V. Ups and downs: the STAT1:STAT3 seesaw of interferon and gp130 receptor signalling. *Semin. Cell Dev. Biol.* **19**, 351–359 (2008).
- Stephanou, A., Brar, B. K., Knight, R. A. & Latchman, D. S. Opposing actions of STAT-1 and STAT-3 on the Bcl-2 and Bcl-x promoters. *Cell Death Differ.* **7**, 329–330 (2000).
- Azare, J. *et al.* Constitutively activated Stat3 induces tumorigenesis and enhances cell motility of prostate epithelial cells through integrin β 6. *Mol. Cell. Biol.* **27**, 4444–4453 (2007).
- Dechow, T. N. *et al.* Requirement of matrix metalloproteinase-9 for the transformation of human mammary epithelial cells by Stat3-C. *Proc. Natl Acad. Sci. USA* **101**, 10602–10607 (2004).
- Dustin, M. L., Singer, K. H., Tuck, D. T. & Springer, T. A. Adhesion of T lymphoblasts to epidermal keratinocytes is regulated by interferon γ and is mediated by intercellular adhesion molecule 1 (ICAM-1). *J. Exp. Med.* **167**, 1323–1340 (1988).
- Yoshimura, A. Signal transduction of inflammatory cytokines and tumor development. *Cancer Sci.* **97**, 439–447 (2006).
- Bruet, M., Marques, L., Sebastian, C., Lloberas, J. & Celada, A. Regulation of murine Tap1 and Lmp2 genes in macrophages by interferon γ is mediated by STAT1 and IRF-1. *Genes Immun.* **5**, 26–35 (2004).
- Marques, L., Bruet, M., Lloberas, J. & Celada, A. STAT1 regulates lipopolysaccharide- and TNF- α -dependent expression of transporter associated with antigen processing 1 and low molecular mass polypeptide 2 genes in macrophages by distinct mechanisms. *J. Immunol.* **173**, 1103–1110 (2004).
- Ito, S. *et al.* Interleukin-10 inhibits expression of both interferon α - and interferon γ -induced genes by suppressing tyrosine phosphorylation of STAT1. *Blood* **93**, 1456–1463 (1999).
- Yasukawa, H. *et al.* IL-6 induces an anti-inflammatory response in the absence of SOCS3 in macrophages. *Nature Immunol.* **4**, 551–556 (2003).
- El Kasmi, K. C. *et al.* General nature of the STAT3-activated anti-inflammatory response. *J. Immunol.* **177**, 7880–7888 (2006).
- Tanabe, Y. *et al.* Cutting edge: role of STAT1, STAT3, and STAT5 in IFN- α responses in T lymphocytes. *J. Immunol.* **174**, 609–613 (2005).
- Qing, Y. & Stark, G. R. Alternative activation of STAT1 and STAT3 in response to interferon- γ . *J. Biol. Chem.* **279**, 41679–41685 (2004).
- Chang, E. Y., Guo, B., Doyle, S. E. & Cheng, G. Cutting edge: involvement of the type I IFN production and signaling pathway in lipopolysaccharide-induced IL-10 production. *J. Immunol.* **178**, 6705–6709 (2007).
- Wang, X., Chen, M., Wandinger, K. P., Williams, G. & Dhib-Jalbut, S. IFN- β 1b inhibits IL-12 production in peripheral blood mononuclear cells in an IL-10-dependent mechanism: relevance to IFN- β 1b therapeutic effects in multiple sclerosis. *J. Immunol.* **165**, 548–557 (2000).
- Ziegler-Heitbrock, L. *et al.* IFN- α induces the human IL-10 gene by recruiting both IFN regulatory factor 1 and Stat3. *J. Immunol.* **171**, 285–290 (2003).
- Wang, H. *et al.* The role of glycogen synthase kinase 3 in regulating IFN- β -mediated IL-10 production. *J. Immunol.* **186**, 675–684 (2011).
- Uddin, S. *et al.* Activation of the p38 mitogen-activated protein kinase by type I interferons. *J. Biol. Chem.* **274**, 30127–30131 (1999).
- Uddin, S. *et al.* The Rac1/p38 mitogen-activated protein kinase pathway is required for interferon α -dependent transcriptional activation but not serine phosphorylation of Stat proteins. *J. Biol. Chem.* **275**, 27634–27640 (2000).
- Li, Y. *et al.* Role of p38 α Map kinase in type I interferon signaling. *J. Biol. Chem.* **279**, 970–979 (2004).
- Mayer, I. A. *et al.* The p38 MAPK pathway mediates the growth inhibitory effects of interferon- α in BCR-ABL-expressing cells. *J. Biol. Chem.* **276**, 28570–28577 (2001).

50. Ishida, H. *et al.* Involvement of p38 signaling pathway in interferon- α -mediated antiviral activity toward hepatitis C virus. *Biochem. Biophys. Res. Commun.* **321**, 722–727 (2004).
51. David, M. *et al.* Requirement for MAP kinase (ERK2) activity in interferon α - and interferon β -stimulated gene expression through STAT proteins. *Science* **269**, 1721–1723 (1995).
52. Wang, F. *et al.* Disruption of Erk-dependent type I interferon induction breaks the myxoma virus species barrier. *Nature Immunol.* **5**, 1266–1274 (2004).
53. Uddin, S. *et al.* Interferon- α engages the insulin receptor substrate-1 to associate with the phosphatidylinositol 3'-kinase. *J. Biol. Chem.* **270**, 15938–15941 (1995).
54. Kaur, S. *et al.* Role of the Akt pathway in mRNA translation of interferon-stimulated genes. *Proc. Natl Acad. Sci. USA* **105**, 4808–4813 (2008).
55. Kaur, S. *et al.* Dual regulatory roles of phosphatidylinositol 3-kinase in IFN signaling. *J. Immunol.* **181**, 7316–7323 (2008).
56. Lekmine, F. *et al.* Activation of the p70 S6 kinase and phosphorylation of the 4E-BP1 repressor of mRNA translation by type I interferons. *J. Biol. Chem.* **278**, 27772–27780 (2003).
57. Lekmine, F. *et al.* Interferon- γ engages the p70 S6 kinase to regulate phosphorylation of the 40S S6 ribosomal protein. *Exp. Cell Res.* **295**, 173–182 (2004).
58. Matsumoto, A. *et al.* Interferon- α -induced mTOR activation is an anti-hepatitis C virus signal via the phosphatidylinositol 3-kinase–Akt-independent pathway. *J. Gastroenterol.* **44**, 856–863 (2009).
59. Round, J. L. & Mazmanian, S. K. The gut microbiota shapes intestinal immune responses during health and disease. *Nature Rev. Immunol.* **9**, 313–323 (2009).
60. Katakura, K. *et al.* Toll-like receptor 9-induced type I IFN protects mice from experimental colitis. *J. Clin. Invest.* **115**, 695–702 (2005).
This study demonstrated that the protective effect of TLR9 signalling in colonic injury is mediated by type I IFNs.
61. Rachmilewitz, D. *et al.* Toll-like receptor 9 signaling mediates the anti-inflammatory effects of probiotics in murine experimental colitis. *Gastroenterology* **126**, 520–528 (2004).
62. Vijay-Kumar, M. *et al.* Activation of Toll-like receptor 3 protects against DSS-induced acute colitis. *Inflamm. Bowel Dis.* **13**, 856–864 (2007).
63. McFarland, A. P. *et al.* Localized delivery of interferon- β by *Lactobacillus* exacerbates experimental colitis. *PLoS ONE* **6**, e16967 (2011).
64. Abe, K. *et al.* Conventional dendritic cells regulate the outcome of colonic inflammation independently of T cells. *Proc. Natl Acad. Sci. USA* **104**, 17022–17027 (2007).
65. Hall, J. A. *et al.* Commensal DNA limits regulatory T cell conversion and is a natural adjuvant of intestinal immune responses. *Immunity* **29**, 637–649 (2008).
66. Hofmann, C. *et al.* T cell-dependent protective effects of CpG motifs of bacterial DNA in experimental colitis are mediated by CD11c⁺ dendritic cells. *Gut* **59**, 1347–1354 (2010).
67. Bleich, A. *et al.* CpG motifs of bacterial DNA exert protective effects in mouse models of IBD by antigen-independent tolerance induction. *Gastroenterology* **136**, 278–287 (2009).
68. Bilsborough, J., George, T. C., Norment, A. & Viney, J. L. Mucosal CD8a⁺ DC, with a plasmacytoid phenotype, induce differentiation and support function of T cells with regulatory properties. *Immunology* **108**, 481–492 (2003).
69. Levings, M. K. *et al.* IFN- α and IL-10 induce the differentiation of human type 1 T regulatory cells. *J. Immunol.* **166**, 5530–5539 (2001).
70. Szabo, S. J. *et al.* Distinct effects of Tbet in T_H1 lineage commitment and IFN- γ production in CD4 and CD8 T cells. *Science* **295**, 338–342 (2002).
71. Cho, S. S. *et al.* Activation of STAT4 by IL-12 and IFN- α : evidence for the involvement of ligand-induced tyrosine and serine phosphorylation. *J. Immunol.* **157**, 4781–4789 (1996).
72. Rogge, L. *et al.* The role of Stat4 in species-specific regulation of Th cell development by type I IFNs. *J. Immunol.* **161**, 6567–6574 (1998).
73. Cousens, L. P. *et al.* Two roads diverged: interferon α/β - and interleukin 12-mediated pathways in promoting T cell interferon γ responses during viral infection. *J. Exp. Med.* **189**, 1315–1328 (1999).
74. Nguyen, K. B. *et al.* Critical role for STAT4 activation by type 1 interferons in the interferon- γ response to viral infection. *Science* **297**, 2063–2066 (2002).
75. Berenson, L. S., Farrar, J. D., Murphy, T. L. & Murphy, K. M. Frontline: absence of functional STAT4 activation despite detectable tyrosine phosphorylation induced by murine IFN- α . *Eur. J. Immunol.* **34**, 2365–2374 (2004).
76. Farrar, J. D., Smith, J. D., Murphy, T. L. & Murphy, K. M. Recruitment of Stat4 to the human interferon- α/β receptor requires activated Stat2. *J. Biol. Chem.* **275**, 2693–2697 (2000).
77. Berenson, L. S., Gavioli, M., Farrar, J. D., Murphy, T. L. & Murphy, K. M. Distinct characteristics of murine STAT4 activation in response to IL-12 and IFN- α . *J. Immunol.* **177**, 5195–5203 (2006).
78. Huber, J. P. & Farrar, J. D. Regulation of effector and memory T-cell functions by type I interferon. *Immunology* **132**, 466–474 (2011).
79. Matikainen, S. *et al.* IFN- α and IL-18 synergistically enhance IFN- γ production in human NK cells: differential regulation of Stat4 activation and IFN- γ gene expression by IFN- α and IL-12. *Eur. J. Immunol.* **31**, 2236–2245 (2001).
80. Strengell, M., Julkunen, I. & Matikainen, S. IFN- α regulates IL-21 and IL-21R expression in human NK and T cells. *J. Leukoc. Biol.* **76**, 416–422 (2004).
81. Harrington, L. E. *et al.* Interleukin 17-producing CD4⁺ effector T cells develop via a lineage distinct from the T helper type 1 and 2 lineages. *Nature Immunol.* **6**, 1125–1132 (2005).
82. Moschen, A. R., Geiger, S., Krehan, I., Kaser, A. & Tilg, H. Interferon- α controls IL-17 expression *in vitro* and *in vivo*. *Immunobiology* **213**, 779–787 (2008).
83. Qiu, H. *et al.* Type I IFNs enhance susceptibility to *Chlamydia muridarum* lung infection by enhancing apoptosis of local macrophages. *J. Immunol.* **181**, 2092–2102 (2008).
84. Karaghiosoff, M. *et al.* Central role for type I interferons and Tyk2 in lipopolysaccharide-induced endotoxin shock. *Nature Immunol.* **4**, 471–477 (2003).
85. Vandenbark, A. A. *et al.* Interferon- β -1a treatment increases CD56^{bright} natural killer cells and CD4⁺CD25⁺Foxp3 expression in subjects with multiple sclerosis. *J. Neuroimmunol.* **215**, 125–128 (2009).
86. Zhou, X. *et al.* Instability of the transcription factor Foxp3 leads to the generation of pathogenic memory T cells *in vivo*. *Nature Immunol.* **10**, 1000–1007 (2009).
87. Oldenhove, G. *et al.* Decrease of Foxp3⁺ Treg cell number and acquisition of effector cell phenotype during lethal infection. *Immunity* **31**, 772–786 (2009).
88. Dominguez-Villar, M., Baecher-Allan, C. M. & Hafler, D. A. Identification of T helper type 1-like, Foxp3⁺ regulatory T cells in human autoimmune disease. *Nature Med.* **17**, 673–675 (2011).
89. Schroder, K. & Tschopp, J. The inflammasomes. *Cell* **140**, 821–832 (2010).
90. Guarda, G. *et al.* Type I interferon inhibits interleukin-1 production and inflammasome activation. *Immunity* **34**, 213–223 (2011).
An important study showing that type I IFN signalling inhibits NLRP1 and NLRP3 inflammasomes in a two-step process through STAT1 and IL-10.
91. Henry, T., Brotcke, A., Weiss, D. S., Thompson, L. J. & Monack, D. M. Type I interferon signaling is required for activation of the inflammasome during *Francisella* infection. *J. Exp. Med.* **204**, 987–994 (2007).
92. Fernandes-Alnemri, T. *et al.* The AIM2 inflammasome is critical for innate immunity to *Francisella tularensis*. *Nature Immunol.* **11**, 385–393 (2010).
93. Fujita, T. *et al.* Induction of the transcription factor IRF-1 and interferon- β mRNAs by cytokines and activators of second-messenger pathways. *Proc. Natl Acad. Sci. USA* **86**, 9936–9940 (1989).
94. Rivieccio, M. A. *et al.* The cytokine IL-1 β activates IFN response factor 3 in human fetal astrocytes in culture. *J. Immunol.* **174**, 3719–3726 (2005).
95. Gonzalez-Navajas, J. M. *et al.* Interleukin 1 receptor signaling regulates DUBA expression and facilitates Toll-like receptor 9-driven anti-inflammatory cytokine production. *J. Exp. Med.* **207**, 2799–2807 (2010).
96. Tseng, P. H. *et al.* Different modes of ubiquitination of the adaptor TRAF3 selectively activate the expression of type I interferons and proinflammatory cytokines. *Nature Immunol.* **11**, 70–75 (2010).
An elegant study showing the different effects of K63- and K48-linked polyubiquitylation on the biological functions of TRAF3.
97. Kayagaki, N. *et al.* DUBA: a deubiquitinase that regulates type I interferon production. *Science* **318**, 1628–1632 (2007).
The first report that DUBA binds to TRAF3 and selectively cleaves the K63-linked polyubiquitin chains, thereby dampening type I IFN production.
98. Lebeis, S. L., Powell, K. R., Merlin, D., Sherman, M. A. & Kalman, D. Interleukin-1 receptor signaling protects mice from lethal intestinal damage caused by the attaching and effacing pathogen *Citrobacter rodentium*. *Infect. Immun.* **77**, 604–614 (2009).
99. Kojouharoff, G. *et al.* Neutralization of tumour necrosis factor (TNF) but not of IL-1 reduces inflammation in chronic dextran sulphate sodium-induced colitis in mice. *Clin. Exp. Immunol.* **107**, 353–358 (1997).
100. Auerbuch, V., Brockstedt, D. G., Meyer-Morse, N., O'Riordan, M. & Portnoy, D. A. Mice lacking the type I interferon receptor are resistant to *Listeria monocytogenes*. *J. Exp. Med.* **200**, 527–533 (2004).
101. Carrero, J. A., Calderon, B. & Unanue, E. R. Type I interferon sensitizes lymphocytes to apoptosis and reduces resistance to *Listeria* infection. *J. Exp. Med.* **200**, 535–540 (2004).
102. Stanley, S. A., Johndrow, J. E., Manzanillo, P. & Cox, J. S. The type I IFN response to infection with *Mycobacterium tuberculosis* requires ESX-1-mediated secretion and contributes to pathogenesis. *J. Immunol.* **178**, 3143–3152 (2007).
103. Nagarajan, U. M. *et al.* Type I interferon signaling exacerbates *Chlamydia muridarum* genital infection in a murine model. *Infect. Immun.* **76**, 4642–4648 (2008).
104. Henry, T. *et al.* Type I IFN signaling constrains IL-17A/F secretion by $\gamma\delta$ T cells during bacterial infections. *J. Immunol.* **184**, 3755–3767 (2010).
105. Carrero, J. A. & Unanue, E. R. Lymphocyte apoptosis as an immune subversion strategy of microbial pathogens. *Trends Immunol.* **27**, 497–503 (2006).
106. Mancuso, G. *et al.* Type I IFN signaling is crucial for host resistance against different species of pathogenic bacteria. *J. Immunol.* **178**, 3126–3133 (2007).
107. Coers, J., Vance, R. E., Fontana, M. F. & Dietrich, W. F. Restriction of *Legionella pneumophila* growth in macrophages requires the concerted action of cytokine and Naip5/Ipaf signalling pathways. *Cell. Microbiol.* **9**, 2344–2357 (2007).
108. Burger, D. & Travis, S. Conventional medical management of inflammatory bowel disease. *Gastroenterology* **140**, 1827–1837 (2011).
109. Musch, E., Andus, T., Malek, M., Chrissafidou, A. & Schulz, M. Successful treatment of steroid refractory active ulcerative colitis with natural interferon- β — an open long-term trial. *Z. Gastroenterol.* **45**, 1255–1240 (2007).
110. Nikolaus, S. *et al.* Interferon β -1a in ulcerative colitis: a placebo controlled, randomised, dose escalating study. *Gut* **52**, 1286–1290 (2003).
111. Madsen, S. M. *et al.* An open-labeled, randomized study comparing systemic interferon- α 2A and prednisolone enemas in the treatment of left-sided ulcerative colitis. *Am. J. Gastroenterol.* **96**, 1807–1815 (2001).
112. Pena-Rossi, C. *et al.* Interferon β -1a for the maintenance of remission in patients with Crohn's disease: results of a phase II dose-finding study. *BMC Gastroenterol.* **9**, 22 (2009).
113. Pena-Rossi, C. *et al.* Clinical trial: a multicentre, randomized, double-blind, placebo-controlled, dose-finding, phase II study of subcutaneous interferon- β -1a in moderately active ulcerative colitis. *Aliment. Pharmacol. Ther.* **28**, 758–767 (2008).
114. Gasche, C. *et al.* Prospective evaluation of interferon- α in treatment of chronic active Crohn's disease. *Dig. Dis. Sci.* **40**, 800–804 (1995).
115. Musch, E. *et al.* Interferon- β -1a for the treatment of steroid-refractory ulcerative colitis: a randomized, double-blind, placebo-controlled trial. *Clin. Gastroenterol. Hepatol.* **3**, 581–586 (2005).
116. Mitoro, A. *et al.* Exacerbation of ulcerative colitis during α -interferon therapy for chronic hepatitis C. *Intern. Med.* **32**, 327–331 (1993).
117. Watanabe, T. *et al.* A case of exacerbation of ulcerative colitis induced by combination therapy with PEG-interferon α 2b and ribavirin. *Gut* **55**, 1682–1683 (2006).
118. Schott, E. *et al.* Development of ulcerative colitis in a patient with multiple sclerosis following treatment with interferon β 1a. *World J. Gastroenterol.* **13**, 3638–3640 (2007).

119. Freeman, H. J., Chopra, A., Clandinin, M. T. & Thomson, A. B. Recent advances in celiac disease. *World J. Gastroenterol.* **17**, 2259–2272 (2011).
120. MacDonald, T. T. & Spencer, J. Evidence that activated mucosal T cells play a role in the pathogenesis of enteropathy in human small intestine. *J. Exp. Med.* **167**, 1341–1349 (1988).
121. Neurath, M. F., Finotto, S. & Glimcher, L. H. The role of Th1/Th2 polarization in mucosal immunity. *Nature Med.* **8**, 567–573 (2002).
122. Monteleone, G., Pender, S. L., Wathen, N. C. & MacDonald, T. T. Interferon- α drives T cell-mediated immunopathology in the intestine. *Eur. J. Immunol.* **31**, 2247–2255 (2001).
123. Di Sabatino, A. *et al.* Evidence for the role of interferon- α production by dendritic cells in the Th1 response in celiac disease. *Gastroenterology* **133**, 1175–1187 (2007).
124. Molberg, O. *et al.* Tissue transglutaminase selectively modifies gliadin peptides that are recognized by gut-derived T cells in celiac disease. *Nature Med.* **4**, 713–717 (1998).
125. Molberg, O. *et al.* Gliadin specific, HLA DQ2-restricted T cells are commonly found in small intestinal biopsies from coeliac disease patients, but not from controls. *Scand. J. Immunol.* **46**, 103–109 (1997).
126. Kagnoff, M. F. Celiac disease: pathogenesis of a model immunogenetic disease. *J. Clin. Invest.* **117**, 41–49 (2007).
127. Ting, J. P. & Trowsdale, J. Genetic control of MHC class II expression. *Cell* **109**, S21–S33 (2002).
128. Cammarota, G., Cuoco, L., Cianci, R., Pandolfi, F. & Gasbarrini, G. Onset of coeliac disease during treatment with interferon for chronic hepatitis C. *Lancet* **356**, 1494–1495 (2000).
129. Zheng, Y. *et al.* Interleukin-22, a T_H17 cytokine, mediates IL-23-induced dermal inflammation and acanthosis. *Nature* **445**, 648–651 (2007).
130. Nestle, F. O. *et al.* Plasmacytoid dendritic cells initiate psoriasis through interferon- α production. *J. Exp. Med.* **202**, 135–143 (2005).
131. Dzionek, A. *et al.* BDCA-2, a novel plasmacytoid dendritic cell-specific type II C-type lectin, mediates antigen capture and is a potent inhibitor of interferon α/β induction. *J. Exp. Med.* **194**, 1823–1834 (2001).
132. Hua, J., Kirou, K., Lee, C. & Crow, M. K. Functional assay of type I interferon in systemic lupus erythematosus plasma and association with anti-RNA binding protein autoantibodies. *Arthritis Rheum.* **54**, 1906–1916 (2006).
133. Kirou, K. A. *et al.* Activation of the interferon- α pathway identifies a subgroup of systemic lupus erythematosus patients with distinct serologic features and active disease. *Arthritis Rheum.* **52**, 1491–1503 (2005).
134. Blanco, P., Palucka, A. K., Gill, M., Pascual, V. & Banchereau, J. Induction of dendritic cell differentiation by IFN- α in systemic lupus erythematosus. *Science* **294**, 1540–1543 (2001).
135. Jego, G. *et al.* Plasmacytoid dendritic cells induce plasma cell differentiation through type I interferon and interleukin 6. *Immunity* **19**, 225–234 (2003).
136. Yao, Y. *et al.* Neutralization of interferon- α/β -inducible genes and downstream effect in a phase I trial of an anti-interferon- α monoclonal antibody in systemic lupus erythematosus. *Arthritis Rheum.* **60**, 1785–1796 (2009).
137. Ann Marrie, R. & Rudick, R. A. Drug insight: interferon treatment in multiple sclerosis. *Nature Clin. Pract. Neurol.* **2**, 34–44 (2006).
138. Bendtzen, K. Critical review: assessment of interferon- β immunogenicity in multiple sclerosis. *J. Interferon Cytokine Res.* **30**, 759–766 (2010).
139. The IFNB Multiple Sclerosis Study Group. Interferon β -1b is effective in relapsing-remitting multiple sclerosis. I. Clinical results of a multicenter, randomized, double-blind, placebo-controlled trial. *Neurology* **43**, 655–661 (1993).
140. Teige, I. *et al.* IFN- β gene deletion leads to augmented and chronic demyelinating experimental autoimmune encephalomyelitis. *J. Immunol.* **170**, 4776–4784 (2003).
141. Brod, S. A. & Burns, D. K. Suppression of relapsing experimental autoimmune encephalomyelitis in the SJL/J mouse by oral administration of type I interferons. *Neurology* **44**, 1144–1148 (1994).
142. Prinz, M. *et al.* Distinct and nonredundant *in vivo* functions of IFNAR on myeloid cells limit autoimmunity in the central nervous system. *Immunity* **28**, 675–686 (2008).
143. Guo, B., Chang, E. Y. & Cheng, G. The type I IFN induction pathway constrains Th17-mediated autoimmune inflammation in mice. *J. Clin. Invest.* **118**, 1680–1690 (2008).
144. Teige, I., Liu, Y. & Issazadeh-Navikas, S. IFN- β inhibits T cell activation capacity of central nervous system APCs. *J. Immunol.* **177**, 3542–3553 (2006).
145. Rudick, R. A. *et al.* Interferon β induces interleukin-10 expression: relevance to multiple sclerosis. *Ann. Neurol.* **40**, 618–627 (1996).
146. Ferrantini, M., Capone, I. & Belardelli, F. Interferon- α and cancer: mechanisms of action and new perspectives of clinical use. *Biochimie* **89**, 884–893 (2007).
147. Rizza, P., Moretti, F. & Belardelli, F. Recent advances on the immunomodulatory effects of IFN- α : implications for cancer immunotherapy and autoimmunity. *Autoimmunity* **43**, 204–209 (2010).
148. Ferrantini, M. & Belardelli, F. Gene therapy of cancer with interferon: lessons from tumor models and perspectives for clinical applications. *Semin. Cancer Biol.* **10**, 145–157 (2000).
149. Di Pucchio, T. *et al.* Immunization of stage IV melanoma patients with Melan-A/MART-1 and gp100 peptides plus IFN- α results in the activation of specific CD8⁺ T cells and monocyte/dendritic cell precursors. *Cancer Res.* **66**, 4943–4951 (2006).
150. Lapenta, C. *et al.* IFN- α -conditioned dendritic cells are highly efficient in inducing cross-priming CD8⁺ T cells against exogenous viral antigens. *Eur. J. Immunol.* **36**, 2046–2060 (2006).
151. Lapenta, C. *et al.* Potent immune response against HIV-1 and protection from virus challenge in hu-PBL-SCID mice immunized with inactivated virus-pulsed dendritic cells generated in the presence of IFN- α . *J. Exp. Med.* **198**, 361–367 (2003).
152. Santodonato, L. *et al.* Monocyte-derived dendritic cells generated after a short-term culture with IFN- α and granulocyte-macrophage colony-stimulating factor stimulate a potent Epstein-Barr virus-specific CD8⁺ T cell response. *J. Immunol.* **170**, 5195–5202 (2003).

Acknowledgements

We thank S. Herdman for his editorial assistance. This work was supported by US National Institutes of Health grants AI068685, AI077989, AI095623 and DK35108; grants from the Crohn's and Colitis Foundation of America; and grant CP10/00417 from Instituto de Salud Carlos III, Madrid, Spain.

Competing interests statement

The authors declare no competing financial interests.